

Constituents of Emerging Concern

Current Regulatory Framework and Results on Russian River Watershed Pilot Study

Jeremiah Puget – Regional Water Board

Dr. Alvina Mehinto – Southern CA Coastal Water Research Project

Dr. Rebecca Sutton – San Francisco Estuary Institute

Jennifer Sun – San Francisco Estuary Institute

Item No. 7

North Coast Regional Water Quality Control Board


February 8, 2018



Presentation Outline

- Background on Statewide Efforts *(15 min.)*
- Russian River Pilot Study Results
 - ❖ Water & Sediment - Analytical & BioAnalytical Screens
 - Dr. Alvina Mehinto – SCCWRP *(15 min.)*
 - ❖ Fish Tissue
 - Dr. Rebecca Sutton- SFEI *(15 min.)*
 - ❖ Pesticides
 - Jennifer Sun – SFEI *(15 min.)*
- Next Steps
- Questions and Comments

Constituents of Emerging Concern



Challenges to Current Monitoring

- Too many chemicals to monitor
 - Over 100,000 known chemicals
 - More discovered every year
- No standardized analytical methods for unexpected and/or unknown chemicals incl. metabolites, byproducts
- Relevant toxicity data often unavailable
 - Chronic sub-lethal toxicity is of concern
 - Toxicity potential of chemical mixtures understudied

Pathways to the Environment

Treated Wastewater

- Permitted Discharges
- Recycled Water
- Biosolids



Septic Tanks Landfills




Agricultural Runoff Industrial Discharges Storm Water Runoff



Regulatory Framework for CECs

- Recycled Water Policy (2009)
 - CEC Expert Panel (2010)
 - *Monitoring Strategies for Chemicals of Emerging Concern in Recycled Water (2010)*
 - *Monitoring Strategies for Chemicals of Emerging Concern in California's Aquatic Ecosystems (2012)*
- Policy Amendment (2013)
 - Included monitoring and reporting of recycled water used for groundwater recharge projects
- Current Policy Amendment (2018/2019)
 - *Updated CEC Panel Recommendations for Recycled Water (Draft report is currently available for public review)*

Regulatory Framework for CECs




State Water Board Role

- Identify and improve the knowledge base
- Work with DWQ, DDW, Regions, and Expert Panel to develop and implement monitoring strategies for recycled water and other types of discharges
- Track and help evaluate effectiveness of regulatory interventions
- Direct pilot monitoring in ambient recommended by expert panel



CEC Initiative




Origin of the Ecosystem Panel

- State of knowledge regarding CECs is incomplete
- Regulatory requirements need to be based on best available peer-reviewed science
- Experts needed to guide future monitoring activities
- All members of Recycled Water Panel retained, with the addition of experts in marine resources & antibiotic resistance

Is there a better way to monitor CECs?

- Adaptive management
 - Collect and *interpret* data
 - *Adjust* target parameters, monitoring effort
 - Test promising *new* technologies



Is there a better way to monitor CECs?

- New monitoring tools
 - *bioanalytical tools* to screen for toxicants by mode of action
 - *non-targeted analysis* to identify toxicants that elude targeted methods
- Develop monitoring thresholds
 - *Monitoring Trigger Levels* (MTLs)
 - Measured environmental concentrations (MEC)
 - Predicted environmental concentrations (PEC)
- Research initiatives
 1. Developing of *bioanalytical screening tools*;
 2. Filling *data gaps* on CEC sources, fate, occurrence and toxicity; and
 3. Assessing the *relative risk* of CECs and other monitored chemicals.

CEC Monitoring Methods

Biological



Bioassessment


Chemical




Targeted



Bioanalytical




Non-targeted



?

Russian River CEC Pilot Study




Russian River CEC Pilot Study


- Are CECs in WWTP effluent and storm water runoff present?
- What is the relative contribution of treated wastewater effluent and storm water runoff to CEC loading into the watershed?
- Do bioanalytical tools effectively screen for the occurrence of CECs?
- What is the extent and magnitude of CECs are in the water column, sediments and fish tissue?
- Which pesticides applied in the Russian River watershed are of highest priority for monitoring

Tools for Russian River CEC Study

**Targeted
Chemistry**





Bioanalytical




BioAssessment



North Coast Region - Russian River CEC Monitoring Sites

	Sediment		Water
	Fish Tissue		Region 1 Boundary
	Urban Areas		




Screening for CECs in Water and Sediment from the Russian River Watershed

Dr. Alvina Mehinto, Dr. Keith Maruya
Southern California Coastal Water Research Project




Effect-Based Monitoring




- Framework currently considered by the State Water Board
- New tools proposed to:
 - *Streamline existing monitoring approaches*
 - *Enhance capabilities to identify new and/or unknown contaminants*
 - *Identify ecologically relevant impacts*

What Are Cell Assays?

- Cells engineered to respond to specific classes of CECs
- Light intensity is proportional to the concentration of bioactive chemicals
- Results expressed relative to a known/reference chemical
 - *Bioanalytical equivalent concentration (BEQ, ng/L)*




What Are Cell Assays?



Cell culture




Sample extraction



Add cells + samples
Incubate plate



Add substrate
Then incubate



Measure fluorescence

Advantages of Cell Assays

- Rapid method to screen for hundreds of contaminants simultaneously in one assay
- Integrated measure of known and unknown chemicals acting via a common mode of action
 - *Potential for linkage to toxicity*
- Technology adopted by pharmaceutical, cosmetic and industrial companies to develop their products

Objectives and Study Design

What is the extent and magnitude of **endocrine active CECs** in water and sediment in the Russian River Watershed?



- Water, sediment and effluent samples collected
- Sample analyses:
 - *Cell assay bioscreening (estrogen and glucocorticoid receptor)*
 - *Targeted analyses of known CECs*


Estrogenic Screen of Water Samples

	Effluent #1	Effluent #2	Mirabel	Piner Creek	Santa Rosa Crk	El Roble
ER Bioscreen (ng E2 equiv/L)	<0.5	1.9	<0.5	<0.5	<0.5	<0.5
Targeted chemical analyses (ng/L)						
17b-estradiol (E2)	<0.5	0.6	<0.5	<0.5	<0.5	<0.5
estrone	<0.5	11.0	0.5	0.6	<0.5	<0.5
bisphenol A	<10	12.0	<10	55.0	16	<10
4-nonylphenol	60.8	247	25.4	53.3	62	63
Chem. equiv. (ng/L)	<0.5	1.6	<0.5	<0.5	<0.5	<0.5

Estrogenic Screen of Sediment Samples

	Lytton Spring	Mirabel	Piner Creek	Santa Rosa Crk	El Roble
ER Bioscreen (ng E2 equiv./g)	<0.01	<0.01	0.09	<0.01	<0.01
Targeted chemical analyses (ng/g)					
17b-estradiol (E2)	<0.12	<0.12	0.23	<0.12	<0.12
estrone	<0.12	0.14	1.3	0.4	0.28
bisphenol A	1.4	1.9	15	4.6	<1.0
4-nonylphenol	20	34	29	18	18
bifenthrin	<0.2	<0.2	130	1.96	<0.2
Chem. equiv. (ng/L)	<0.1	<0.1	0.36	<0.1	<0.1

Linking Bioactivity to Toxicity



- Understanding cell assay effect thresholds is key
- Fish studies have shown that exposure to 2 – 4 ng E2/L had no effect on growth and survival
 - *Effluent BEQ of 1.9 ng E2/L (without dilution) = low concern*
 - *River water BEQ < 0.5 ng E2/L = no concern*

Conclusions

- CECs present low to moderate concern in the Russian river
 - *Water concentrations of pharmaceuticals below MTLs*


Analyte	Max. measured conc. (ng/L)	Monitoring trigger level (ng/L)
Diclofenac	< 10	100
Estrone	0.56	6
Ibuprofen	< 10	100

- *Some pesticide concentrations in sediment were > MTLs*

Analyte	Max. measured conc. (ng/g)	Monitoring trigger level (ng/g)
Bifenthrin	130	0.052
Fipronil	3.4	0.09
Permethrin	4.9	0.073

Conclusions


- CECs present low to moderate concern in the Russian river
- Cell assays provided a reliable and integrated measure of estrogenic chemicals
- Routine application of cell assays could provide a cost-effective strategy to prioritize sites requiring more chemical and toxicity testing



CECs in Sport Fish


R1 CEC Pilot Monitoring

Rebecca Sutton, Thomas Jabusch, Jay Davis
San Francisco Estuary Institute




Study Objectives

MQ3. What is the extent and magnitude of PBDE and PFOS contamination in fish tissue in the Russian River Watershed?




Polybrominated diphenyl ethers
(PBDEs)



Study Objectives

MQ3. What is the extent and magnitude of PBDE and PFOS contamination in fish tissue in the Russian River Watershed?



Perfluorooctane Sulfonate
(PFOS)



Study Design

6 popular fishing sites

Sacramento Pikeminnow (5)

Sacramento Sucker (5)

Redear Sunfish (1)

Smallmouth Bass (1)


Largemouth Bass (1)

PBDEs (13 analytes)

PBDE 15, 28, 33, 47, 49, 66, 75, 99, 100, 153, 154, 155, 183

PFASs (13 analytes)



PFBA, PFBS, PFPA, PFHx, PFHxS, PFHpA, PFOA, **PFOS**, PFOSA, PFNA, PFDA, PFUA, PFDoA




Safe to Eat Thresholds

California: Advisory Tissue Levels				
	3 servings/week	2 servings/week	1 serving/week	No Consumption
PBDEs	< 100 ppb	100-210 ppb	210-630 ppb	> 630 ppb
Minnesota: Meal Advice Categories				
	Unrestricted	1 meal/week	1 meal/month	DO NOT EAT
PFOS	≤ 40 ppb	> 40-200 ppb	> 200-800 ppb	> 800 ppb
Michigan: Fish Consumption Screening Values				
	16 meals/month	12 meals/month	8 meals/month	4 meals/month
PFOS	≤ 9 ppb	> 9-13 ppb	> 13-19 ppb	> 19-38 ppb

PBDE Results





PBDEs by Species




PBDEs by Site

Sum of PBDEs (ppb)




To Ocean

PFOS & Other PFASs Results




Lowest Minnesota Meal Advice Threshold

PFOS by Species



PFOS by Site



Conclusions


- Fish tissue findings suggest minimal concern
 - Levels of PBDEs and PFOS generally below available consumption thresholds
 - For PFOS, potential for impacts further up the food chain
- Periodic monitoring (e.g., every 5-10 years) is recommended

Current Use Pesticides

R1 CEC Pilot Monitoring

Jennifer Sun, Rebecca Sutton, Diana Lin

San Francisco Estuary Institute



Study Objectives

MQ4. Which pesticides applied in the Russian River watershed are of highest priority for monitoring?

MQ5. What is the extent and magnitude of pesticide contamination in Russian River water and sediment?




Pesticide Prioritization

DPR Surface Water Monitoring Program modeling tool

The screenshot shows a software window titled "Pesticide Prioritization for Surface Water Monitoring...". It features a "Help" menu and three tabs: "Configuration", "Advanced Options", and "Watershed". The "Configuration" tab is active and contains several sections:

- Use patterns:** Includes checkboxes for "Agricultural use" (checked), "Urban use", and "'Rights of way' (site_code=40)". Below these is a text input field for "Or, user-specified site_code(s)=" with the value "site codes delimited by comma".
- PUR data:** Includes a section "Based on PUR data from" with input fields for "2012" and "2014", and a "Check data" button.
- Toxicity data:** Includes radio buttons for "Acute" (selected), "Chronic", and "Both". It also has checkboxes for "USEPA Aquatic Life Benchmarks" (checked), "Supplemented by Benchmark Equivalent (based on FOOTPRINT PPDB)" (checked), "USEPA Drinking Water Standard", and "USEPA Human Health Benchmark".

A note at the bottom of the toxicity data section reads: "Note: if multiple toxicity databases are selected, the lowest toxicity value for each pesticide will be used for prioritization". A "Prioritize..." button is located at the bottom right of the window.



Pesticide Prioritization

DPR Pesticide Use Database
(2012-2014 data, monthly)

+

USEPA Aquatic Life Benchmarks
or DPR equivalents (acute or
chronic)

+

Physical-chemical properties



Prioritized Pesticide list

Pesticide Prioritization for Surface Water Monitoring...

Help

Configuration | Advanced Options | Watershed

Use patterns

Agricultural use Urban use "Rights of way" (site_code=40)

Or, user-specified site_code(s)=

PUR data

Based on PUR data from to

Toxicity data

Acute Chronic Both

USEPA Aquatic Life Benchmarks

Supplemented by Benchmark Equivalent (based on FOOTPRINT PPDB)

USEPA Drinking Water Standard

USEPA Human Health Benchmark


Note: if multiple toxicity databases are selected, the lowest toxicity value for each pesticide will be used for prioritization

2. Use Maps (Site Selection)

1. Prioritization List (Analytical Lab Selection)

Chemical Name	Water Toxicity Benchmark (ug/L)	DPR Use Score	DPR Toxicity Score	DPR Final Score	Timing
Ethylene thiourea (MANCOZEB degradate)	2	5	4	20	Spring
PENDIMETHALIN	5.2	4	4	16	Spring
CYPRODINIL	8	4	4	16	Fall
OXYFLUORFEN	0.29	3	5	15	Spring
THPA; 482-HA; APF (FLUMIOXAZIN degradates)	0.49	3	5	15	Spring
CHLORPYRIFOS	0.04	2	6	12	Summer
IMIDACLOPRID	1.05	3	4	12	Fall
PYRACLOSTROBIN	1.5	3	4	12	Fall
SIMAZINE	2.24	3	4	12	Spring
TRIFLOXYSTROBIN	2.76	3	4	12	Fall
DIFENOCONAZOLE	5.6	3	4	12	Fall
QUINOXYFEN	7	3	4	12	Fall

Imidacloprid: Total Use 2012-2014



Study Design

5 co-located sites

USGS - CWSC

Sediment

September 2016

118 pesticides

Water

Oct 2016 (“first fall flush”)


162 pesticides (dissolved)


131 pesticides (particulate)



Pesticides in sediment were low


- No exceedances of USGS benchmarks
- Largest number of detections at the mixed use ag- and urban site
- Six pesticides detected
 - Fungicides: boscalid, iprodione
 - Legacy insecticides: DDT, DDD, DDE
 - **Pyrethroid insecticide: bifenthrin**






Stormwater runoff may not have been captured at northern sites



Potter Valley




Trenton Road




Sediment Sampling Event
Water Sampling Event




Two urban insecticides exceeded chronic invertebrate thresholds



Fipronil Sulfone




Fipronil Sulfone




Fipronil



Imidacloprid



*polystyrene insulation,
vinyl siding, adhesives, sealants,
textiles for outdoor use, pressure-
treated wood decking*



Conclusions

- Pesticides from **agricultural runoff are not likely a major concern** during the fall, based on this study
 - Pesticide use varies seasonally – this study did not characterize risks from **spring runoff**
 - Pesticide concentrations may be higher nearer to sources
- Some **urban insecticides** currently exceed or are approaching levels of concern
 - **Imidacloprid** exceeded a USEPA chronic invertebrate benchmark
 - **Fipronil degradates** are approaching or exceed chronic invertebrate threshold
 - **Bifenthrin** is approaching a USGS sediment benchmark

Recommended for monitoring in receiving waters by
California Statewide CEC Expert Panel

Pesticide Monitoring Partners

- **USGS** National Water Quality Assessment: Stream Quality Assessment Project
 - 2017 spring monitoring
 - Trenton Road and Riverfront/Pull-Out sites
- **DPR, SWRCB, CASQA:** statewide framework for urban pesticide monitoring



Lessons Learned

- ✓ BioAnalytical tools show promise
- ✓ Initial screening results for water and fish tissue suggest minimal concern for impacts; however, keep an eye on PFOS
- ✓ Urban use insecticides warrant a closer look
- ✓ Continue implementing improved monitoring strategies

Next Steps

What can be done?

- ✓ Prudent usage of products or use alternative products
- ✓ Proper disposal (*Medicines collected regionally*)
- ✓ Improve treatment technologies
- ✓ Implement expert panel recommended monitoring strategies
- ✓ Efficient and proper use of recycled water
- ✓ Implement the Recycled Water Policy



I WANT

YOU

**To
Properly
Dispose
Unwanted
Medication**

Partnerships

- Other agencies
 - Municipalities
 - Advocates
 - Academia
 - Public
- ✓ Together we can assess conditions and minimize harmful effects



MCHUMOR

by T. McCracken



"Hmphh. Happy as clams, indeed.
They're just all on Prozac."

To be
continued....

Questions?