

Appendix 5.J
Effects on Natural Communities, Wildlife, and Plants

Appendix 5.J

Effects on Natural Communities, Wildlife, and Plants

Contents

1
2
3
4
5
6
7
8
9

	Page
Appendix 5.J Effects on Natural Communities, Wildlife, and Plants	5.J-1
5.J.1 Reference	5.J-69

1 **Tables**

2		Page
3	5.J-1. Quantitative Effects Analysis Methods and Assumptions.....	5.J-2
4	5.J-2. Covered Activities, Effect Types, and Associated Conservation Measures.....	5.J-8
5	5.J-3. Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat	5.J-9
6	5.J-4. Indirect Effect Distances from Covered Activity, Wildlife	5.J-14
7	5.J-5. Indirect Effect Distances from Covered Activity, Plants	5.J-16
8	5.J-6. Near Term Natural Communities Loss by Covered Activity	5.J-17
9	5.J-7. Early Long-Term Natural Communities Loss by Covered Activity	5.J-20
10	5.J-8. Late Long-Term Natural Communities Loss by Covered Activity	5.J-23
11	5.J-9. Near Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity	5.J-26
12	5.J-10. Early Long-Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity	5.J-36
13	5.J-11. Late Long-Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity	5.J-46
14	5.J-12. Near Term Plant Modeled Habitat Loss and Conversion by Covered Activity	5.J-56
15	5.J-13. Early Long-Term Plant Modeled Habitat Loss and Conversion by Covered Activity	5.J-60
16	5.J-14. Late Long-Term Plant Modeled Habitat Loss and Conversion by Covered Activity	5.J-64
17		

1 **Figures**

2 **Figures appear at the end of Appendix**

3

4 5.J-1 Difference between Proposed and Existing Yolo Bypass Flows: Scenario 1

5 5.J-2 Difference between Proposed and Existing Yolo Bypass Flows: Scenario 2

6 5.J-3 Difference between Proposed and Existing Yolo Bypass Flows: Scenario 3

7 5.J-4 Difference between Proposed and Existing Yolo Bypass Flows: Scenario 4

8 5.J-5 Difference between Proposed and Existing Yolo Bypass Flows: Scenario 5

9 5.J-6 Difference between Proposed and Existing Yolo Bypass Flows: Scenario 6

10 5.J-7 Difference between Proposed and Existing Yolo Bypass Flows: Scenario 7

11

12

13

14 **Attachments**

15 **Attachment 5J.A Construction-Related Nitrogen Deposition on BDCP Natural**
16 **Communities**

17 **Attachment 5J.B Natural Community Restoration and Protection Contributing to**
18 **Covered Species Conservation**

19 **Attachment 5J.C Analysis of Potential Bird Collisions at Proposed BDCP Powerlines**

20 **Attachment 5J.D Indirect Effects of the Construction of the BDCP Conveyance Facility on**
21 **Sandhill Crane**

22 **Attachment 5J.E Estimation of BDCP Impact on Giant Garter Snake Summer Foraging**
23 **Habitat (Acreage of Rice) in the Yolo Bypass**

24

Appendix 5.J

Effects on Natural Communities, Wildlife, and Plants

The tables in this appendix support the effects analysis for natural communities and covered wildlife and plant species. The calculations in the tables are presented in summary form in Chapter 5, *Effects Analysis*. The methods are further discussed in Section 5.2, *Methods*.

- Table 5.J-1 presents the methods used and assumptions applied to arrive at quantitative estimates of natural community and species habitat effects, as presented in Section 5.4, *Effects on Natural Communities*, and Section 5.6, *Effects on Wildlife and Plants*.
- Table 5.J-2 indicates the types of effects expected to result from each covered activity, these types of effects correspond with subsections of the effects analysis for each covered wildlife and plant species in Chapter 5. Table 5.J-2 also indicates which conservation measures would be involved for each covered activity.
- Table 5.J-3 provides the key assumptions related to effects of tidal restoration on covered species, based on the species' habitat requirements and the characteristics of the tidal natural community expected to be restored in a particular area (e.g., high tidal marsh, middle tidal marsh, or subtidal). Modifications resulting from tidal restoration are expected to either result in habitat loss, conversion from one habitat type to another (e.g., from primary to secondary habitat), or no change, depending on the existing conditions relative to expected future conditions based on RMA and ESAPWA modeling (Appendix 3.B, *BDCP Tidal Habitat Evolution Assessment*).
- Table 5.J-4 and Table 5.J-5 provide the distances that were applied to arrive at indirect effect acreages for wildlife and plant species, respectively. These indirect effects are described in Section 5.6, *Effects on Wildlife and Plants*.
- Table 5.J-6, Table 5.J-7, and Table 5.J-8 provide the acres of effect expected to result from each covered activity type for natural communities, wildlife, and plants, respectively. These tables provide a breakdown of tidal natural community restoration by elevation zone (e.g., high tidal brackish marsh, middle tidal brackish marsh, low tidal brackish marsh, and subtidal). These tables provide the basis for the quantitative analyses provided in Section 5.4, *Effects on Natural Communities*, and Section 5.6, *Effects on Wildlife and Plants*, for permanent habitat loss and conversion, and for temporary habitat loss.

1 Table 5.J-1. Quantitative Effects Analysis Methods and Assumptions

Activity/Impact Mechanism	Method of Impact Estimation	Key Assumptions ¹ for Purposes of Analysis
CM1 Water Facilities and Operation		
Conveyance facilities construction/permanent removal of natural communities and habitat	<ul style="list-style-type: none"> • GIS layer for construction footprint was overlain on natural community and habitat GIS layers. 	<ul style="list-style-type: none"> • Construction of the forebay, intakes, permanent access roads, and shafts result in permanent removal of natural communities and species habitats located within construction footprint.
Reusable tunnel material/permanent removal of natural communities and habitat	<ul style="list-style-type: none"> • GIS layer for footprint of reusable tunnel material areas was overlain on natural community and habitat GIS layers. • Where AMMs in Appendix 3.C, <i>Avoidance and Minimization Measures</i>, require minimization of the reusable tunnel material footprint or avoidance of a natural community or species habitat, this requirement was factored into the impact estimation for the natural community or species. 	<ul style="list-style-type: none"> • For the purposes of impact analysis, it is assumed reusable tunnel material areas will not be returned pre-project conditions unless required under an AMM. • The final footprint for the reusable tunnel material will be reduced and will meet avoidance and minimization requirements in the AMMs.
Conveyance facilities/temporary removal of natural communities and habitat	<ul style="list-style-type: none"> • GIS layer for footprint of staging areas, intake pipelines, and barge unloading facilities was overlain on natural community and habitat GIS layers. 	<ul style="list-style-type: none"> • Staging areas, intake pipelines, and barge unloading facilities result in temporary impacts on natural communities and species habitats located in the construction footprint of these features. • Affected areas will return to their pre-impact condition following completion of activities (restoration to occur within a year following completion of construction). • Subsurface segments of the tunnel/pipeline have no effects on biological resources.
Borrow/spoil area / temporary loss of natural communities and habitat	<ul style="list-style-type: none"> • GIS layer for footprint of borrow/spoils area was overlain on natural community and habitat GIS layers. • The affected areas will be restored to their former state over the term of the BDCP, but not within the time frame typically characterized as <i>temporary loss</i>. Characterizing this effect as <i>permanent loss</i> would not be accurate. 	<ul style="list-style-type: none"> • Borrow and spoil sites will be reclaimed to their former state over the term of the BDCP, except that cultivated lands will be reclaimed as grasslands if return to cultivated state is not feasible. • Borrow/spoil areas are areas that will initially be used for borrow, and will be used for spoils later. • Restoration to occur within a year following completion of construction.

Activity/Impact Mechanism	Method of Impact Estimation	Key Assumptions ¹ for Purposes of Analysis
CM2 Yolo Bypass Fisheries Enhancement		
Construction/permanent removal of natural communities and habitat	<ul style="list-style-type: none"> • GIS layer for footprint of activities resulting in permanent loss (see <i>Assumptions</i>) was overlain on natural community and habitat GIS layers. 	<ul style="list-style-type: none"> • Permanent loss of natural communities and habitat will result from Fremont Weir improvements, Putah Creek realignment activities, Lisbon Weir and fish crossing improvements, and Sacramento Weir improvements.
Construction/temporary removal of natural communities and habitat	<ul style="list-style-type: none"> • GIS layer for footprint of activities resulting in permanent loss (see <i>Assumption</i>) was overlain on natural community and habitat GIS layers. 	<ul style="list-style-type: none"> • Temporary loss of natural communities and habitat will result from construction areas associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon Weir and fish crossing improvements, and Sacramento Weir improvements.
Operation/periodic inundation from flooding in Yolo Bypass	<ul style="list-style-type: none"> • Compared inundation areas under existing and proposed Fremont Weir flows for seven proposed flow scenarios under the MIKE21 model (see Appendix 5.C, Attachment 5C.E, <i>BDCP Effects Analysis: 2D Hydrodynamic Modeling of the Fremont Weir Diversion Structure</i>, for description of the model) (1,000-cfs notch flow to 6,000-cfs notch flow, with two different baseline flow scenarios for 6,000 cfs). For each of these seven scenarios, the GIS footprint for the difference between existing and proposed flows were overlain on GIS layers for natural communities and modeled covered species habitat. Figures 5.J-1 through 5.J-7 show the footprint of the difference between existing and proposed conditions for each flow scenario. Results from all seven scenarios are presented in Chapter 5, <i>Effects Analysis</i>, for each natural community and covered species affected. 	<ul style="list-style-type: none"> • Fremont Weir will be operated as described in Conservation Measure 2.

Activity/Impact Mechanism	Method of Impact Estimation	Key Assumptions ¹ for Purposes of Analysis
CM4 Tidal Natural Communities Restoration		
Inundation/ permanent loss of natural communities and species habitat	<ul style="list-style-type: none"> • GIS layer for hypothetical tidal restoration footprint¹ (see Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>, for a description of the tidal restoration hypothetical), including only those areas below EHW elevation, was overlain on natural community and species modeled habitat GIS layers. • Exceptions: <ul style="list-style-type: none"> ○ Natural communities: The tidal perennial aquatic natural community was not treated as lost as a result of inundation. Tidal brackish emergent wetland and tidal freshwater emergent wetland natural communities were considered lost for those areas below MLLW + 1 foot. ○ Species: See Table 5.J-3. 	<ul style="list-style-type: none"> • BDCP is not responsible for the long-term effects of sea level rise on natural communities. • All tidally inundated areas below EHW elevation within the hypothetical footprint, based on tidal restoration model, will result in permanent natural community loss, except for to tidal perennial aquatic and tidal emergent wetland natural communities. • All tidally inundated areas below EHW elevation within the hypothetical footprint, based on tidal restoration model, will result in permanent habitat loss for all species except as described in Table 5.J-3.
Inundation/ permanent loss of tidal wetland natural communities and species habitat	<ul style="list-style-type: none"> • GIS data for tidal brackish emergent wetland and tidal freshwater emergent wetland communities was overlain on hypothetical tidal restoration footprint¹ (see Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>, for a description of the tidal restoration hypothetical), including only those areas below EHW elevation, to determine permanent loss and habitat conversion (e.g., conversion from primary habitat to secondary habitat). • See Table 5.J-3 for description of species methods. 	<ul style="list-style-type: none"> • BDCP is not responsible for the long-term effects of sea level rise on tidal wetland natural communities. • All existing tidal aquatic and tidal emergent wetland within the hypothetical footprint below MLLW + 1-foot elevation will be permanently lost. • See Table 5.J-3 for description of assumptions made in regard to permanent loss or conversion for individual covered species.
Inundation/ permanent loss of western pond turtle aquatic habitat	<ul style="list-style-type: none"> • National Hydrologic Database (NHD) GIS data was used to determine the relative percentage of suitable and unsuitable western pond turtle aquatic habitat in artificial water features such as agricultural ditches and canals. 	<ul style="list-style-type: none"> • The visual signature of emergent wetland in the aerial photo indicates perennial water. • The percent cover of suitable habitat within randomly selected grids within each ROA are representative of the entire ROA (see Appendix 2.A, <i>Covered Species Accounts</i>, Section 2.A.29, for a more specific description of this method). • Of all NHD stream miles within the Plan Area, 35% are suitable (Laura Patterson pers. comm. 2012)

¹ The spatial data received from ESAPWA was processed in two ways to prepare it for intersection with natural community and species models: 1.) existing tidal wetlands were removed using a “subtraction” tool in ArcGIS and 2.) the sea level rise accommodation area and upland polygons were removed.

Activity/Impact Mechanism	Method of Impact Estimation	Key Assumptions ¹ for Purposes of Analysis
Riparian restoration within ROAs, natural community permanent loss	<ul style="list-style-type: none"> • All natural community (cultivated land or grassland) loss was applied as permanent habitat loss for a species if cultivated lands or grasslands are major components of the species model and the species distribution overlaps geographically with the ROAs. • Permanent cultivated lands and grassland loss was applied to the foraging habitat value classes for the greater sandhill crane and Swainson’s hawk (Tables 5.6-5 and 5.6-6, respectively). 	<ul style="list-style-type: none"> • 971 acres of riparian restoration will occur as a component of tidal restoration, including 18 acres in Cache Slough ROA, 14 acres in West Delta ROA, and 939 acres in South Delta ROA. • All riparian restoration will occur on existing cultivated land, except for 11 acres in Cache Slough ROA (Conservation Zones 1 and 2) which will occur on existing grassland. • The loss of cultivated land natural community type (e.g., rice, corn, etc.) will occur in proportion to the existing distribution of types within each conservation zone.
CM5 Seasonally Inundated Floodplain Restoration		
Seasonal flooding—periodic inundation of natural communities and habitat	<ul style="list-style-type: none"> • Calculation of effects based on hypothetical floodplain restoration designs. • GIS layer for hypothetical floodplain restoration was overlain on natural community and species habitat layers. 	<ul style="list-style-type: none"> • All areas between setback levees will be subject to periodic inundation from seasonal flooding.
Levee construction—permanent removal of natural communities and habitat	<ul style="list-style-type: none"> • Calculation of effects based on hypothetical floodplain restoration designs. • GIS layer of hypothetical footprint for floodplain levees overlain on natural community species habitat models. 	<ul style="list-style-type: none"> • Floodplain restoration includes an average 1,500-foot setback to levees, with appropriate as-needed grading and lowering of the land elevation to achieve average inundation and intervals noted above. • Floodplain restoration will take place in areas with the greatest potential for restoration, primarily in Conservation Zone 7.
Levee construction—temporary removal of natural communities and habitat	<ul style="list-style-type: none"> • Calculation of effects based on hypothetical floodplain restoration designs. • GIS layer of hypothetical footprint for floodplain levees overlain on natural community species habitat models and buffered 100 feet on each side of the levee footprint. 	<ul style="list-style-type: none"> • Temporary work area of 100 feet on either side of the setback levee base.

Activity/Impact Mechanism	Method of Impact Estimation	Key Assumptions ¹ for Purposes of Analysis
CM7 Riparian Natural Community Restoration		
Permanent loss of natural communities and habitat	<ul style="list-style-type: none"> • The 3,992-acre permanent loss was applied to a species if cultivated lands or grassland are a major component of the species model and the species distribution overlaps geographically with the hypothetical floodplain restoration footprint. • Permanent cultivated lands and grassland natural communities loss was applied to the foraging habitat value classes for the greater sandhill crane and Swainson's hawk (see Tables 5.6-5 and 5.6-6, respectively). 	<ul style="list-style-type: none"> • Riparian restoration in seasonally inundated floodplain will convert up to 3,593 acres of cultivated lands in Conservation Zone 7 and 399 acres of grassland in Conservation Zone 7. • The loss of cultivated land natural community type (e.g., rice, corn, etc.) will occur in proportion to the existing distribution of types within each conservation zone.
CM8 Grassland Natural Community Restoration		
Permanent loss of natural communities and habitat	<ul style="list-style-type: none"> • The 2,000-acre permanent loss was applied to a species if cultivated lands are a major component of the species model and the species distribution overlaps geographically with the hypothetical floodplain restoration footprint. • Permanent cultivated lands natural community loss was applied to the foraging habitat value classes for the greater sandhill crane and Swainson's hawk (see Tables 5.6-5 and 5.6-6, respectively). 	<ul style="list-style-type: none"> • All grassland restoration will require the conversion of cultivated lands to grassland. • 70% of grassland restoration (1,400 acres) will occur in Conservation Zones 1, 8, and 11, 30% (600 acres) in Conservation Zones 2, 4, 5, and 7; restoration acres are thereafter split equally between conservation zones. • The loss of cultivated land natural community type (e.g., rice, corn, etc.) will occur in proportion to the existing distribution of types within each conservation zone.
CM10 Nontidal Marsh Restoration		
Permanent loss of natural communities and habitat	<ul style="list-style-type: none"> • Includes 1,200 acres of nontidal freshwater emergent wetland and nontidal freshwater perennial aquatic restoration and 500 acres of managed wetland restoration, plus additional restoration that may be necessary to meet giant garter snake objectives. • The 1,950-acre permanent loss was applied to a species if cultivated lands are a major component of the species model and the species distribution overlaps geographically with the hypothetical floodplain restoration footprint. • Permanent cultivated lands natural community loss was applied to the foraging habitat value classes for the greater sandhill crane and Swainson's hawk (see Tables 5.6-5 and 5.6-6, respectively). 	<ul style="list-style-type: none"> • All nontidal marsh restoration will require the conversion of cultivated lands to nontidal marsh • 600 acres of nontidal marsh restoration will occur in Conservation Zone 2, 675 acres in Conservation Zone 4 and 675 acres in Conservation Zone 5 • The loss of cultivated land natural community type (e.g., rice, corn, etc.) will occur in proportion to the existing distribution of types within each conservation zone.

Activity/Impact Mechanism	Method of Impact Estimation	Key Assumptions ¹ for Purposes of Analysis
CM11 Natural Community Enhancement and Management		
Construction/Permanent loss of natural communities and habitat	<ul style="list-style-type: none"> The 50-acre grassland loss was applied to permanent habitat loss for a species if grassland is a major component of the species model and the species distribution overlaps geographically with Conservation Zone 1, 4, 5,6, 7, 8, and 11. 	<ul style="list-style-type: none"> LLT Permanent loss was distributed by conservation zones accordingly: 15.5 acres in Conservation Zone 1, 1 acre in Conservation Zone 4, 1.5 acres in Conservation Zone 5, 1.5 acres in Conservation Zone 6, 6.5 acres in Conservation Zone 7, 7.5 acres in Conservation Zone 8, and 16.5 acres in Conservation Zone 11.
CM18 Conservation Hatcheries Facilities		
Construction/Permanent loss of natural communities and habitat	<ul style="list-style-type: none"> The 35-acre grassland loss was applied to permanent habitat loss for a species if grassland is a major component of the species model and the species distribution overlaps geographically with Conservation Zone 1. 	<ul style="list-style-type: none"> Permanent loss of 35 acres of grasslands will result from hatchery construction in Conservation Zone 1.
<p>¹ This table of impact analysis methods and key assumptions is not intended to be all inclusive of all covered activities. Rather, this table shows how effects were calculated for covered activities that have effects significant enough to be estimated. Minor activities described in Chapter 4, <i>Covered Activities and Associated Actions</i>, are covered under the BDCP even though they may not appear in this table. Also, the assumptions made are for the purposes of analysis only and reflect reasonable worst case assumptions for covered activities. Actual footprints of activities may be less than or greater than that assumed and would still fall within the limits of the permits because impacts are within the total range evaluated.</p> <p>cfs = cubic feet per second; GIS = geographic information systems; EHW = extremely high water; NHD = National Hydrology Dataset; MLLW = mean lower low water; ROA = restoration opportunity area.</p>		

1 **Table 5.J-2. Covered Activities, Effect Types, and Associated Conservation Measures**

Covered Activity	Effect Type								Relevant CM
	Permanent Loss/ Conversion	Periodic Inundation	Construction-Related Effects				Permanent Indirect (Adjacent to Activity)	Other Indirect	
			Temporary Loss	Long-Term Loss (Borrow and Spoil)	Injury or Mortality	Temporary Indirect (Adjacent to Activity)			
Conveyance Facility Construction and Operation									
Conveyance facility construction	X		X	X	X	X			CM1
Transmission line construction	X		X		X	X			CM1
Conveyance facility operation								X	CM1
Conveyance facility maintenance						X	X		CM1
Fremont Weir/Yolo Bypass Improvements									
Fisheries enhancement construction	X		X		X	X			CM2
Fisheries enhancement facility maintenance							X		CM2
Yolo Bypass operations		X							CM2
Tidal Restoration									
Grading, levee breaching, and resulting tidal inundation	X				X			X	CM4
Riparian restoration	X								CM4, CM7
Floodplain Restoration									
Levee construction	X		X		X	X			CM5
Restoration activities resulting in seasonal flooding		X			X				CM5
Riparian restoration	X								CM5, CM7
Nontidal Marsh Restoration									
Marsh restoration	X				X	X			CM10
Conservation Hatcheries Facilities									
Facilities construction	X				X	X			
Facilities operation and maintenance						X			
Natural Community and Habitat Enhancement and Management									
Enhancement and management			X		X	X			CM11

2

3

1 Table 5.J-3. Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat

Habitat	Suisun					Delta			
	High Tidal Marsh	Middle Tidal Marsh	Low Tidal Marsh	Intertidal Mudflat	Subtidal	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal
Mammals									
Salt marsh harvest mouse									
Tidal brackish emergent wetland primary	No loss	No loss	Conversion to low value	Loss	Loss	N/A	N/A	N/A	N/A
Tidal brackish emergent wetland secondary	No loss	No loss	No loss	Loss	Loss	N/A	N/A	N/A	N/A
Upland secondary	Conversion to primary	Conversion to primary	No loss	Loss	Loss	N/A	N/A	N/A	N/A
Managed wetland—wetland primary low, long-term conservation value	Conversion to higher value	Conversion to higher value	Conversion to secondary	Loss	Loss	N/A	N/A	N/A	N/A
Managed wetland—wetland secondary low, long-term conservation value	Conversion to higher value primary	Conversion to higher value primary	Conversion to higher value secondary	Loss	Loss	N/A	N/A	N/A	N/A
Managed wetland—upland low, long-term conservation value	Conservation to higher value primary	Conversion to higher value primary	Conservation to higher value secondary	Loss	Loss	N/A	N/A	N/A	N/A
Suisun shrew									
Primary habitat	No loss	No loss	Loss	Loss	Loss	N/A	N/A	N/A	N/A
Secondary habitat	Conversion to primary	Conversion to primary	No loss	Loss	Loss	N/A	N/A	N/A	N/A

Habitat	Suisun					Delta			
	High Tidal Marsh	Middle Tidal Marsh	Low Tidal Marsh	Intertidal Mudflat	Subtidal	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal
Birds									
California black rail									
Primary habitat	No loss	No loss	Conversion to secondary	Loss	Loss	Conversion to secondary	Loss	No loss	Loss
Secondary habitat	Conversion to primary	Conversion to primary	No loss	Loss	Loss	No loss	Loss	Conversion to primary	loss
California clapper rail									
Primary habitat	No loss	No loss	Conversion to secondary	Loss	loss	N/A	N/A	N/A	N/A
Secondary habitat	Conversion to primary	Conversion to primary	No loss	Loss	Loss	N/A	N/A	N/A	N/A
Greater sandhill crane									
Roosting and foraging habitat	N/A	N/A	N/A	N/A	N/A	Partial loss	N/A	Partial loss	Loss
Foraging habitat	N/A	N/A	N/A	N/A	N/A	Partial loss	N/A	Partial loss	Loss
Least Bell's vireo									
Nesting and migratory habitat	Loss	Loss	Loss	Loss	Loss	No loss	Loss	Loss	Loss
Suisun song sparrow									
Primary habitat	No loss	No loss	Conversion to secondary	Loss	Loss	N/A	N/A	N/A	N/A
Secondary habitat	Conversion to primary	Conversion to primary	No loss	Loss	Loss	N/A	N/A	N/A	N/A
Swainson's hawk									
Foraging habitat	Loss	Loss	Loss	Loss	Loss	Partial loss	Loss	Loss	Loss
Nesting habitat	N/A	N/A	N/A	N/A	N/A	No loss	N/A	N/A	N/A

Habitat	Suisun					Delta			
	High Tidal Marsh	Middle Tidal Marsh	Low Tidal Marsh	Intertidal Mudflat	Subtidal	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal
Tricolored blackbird									
Breeding habitat— ag foraging	Partial loss	N/A	N/A	N/A	N/A	Partial loss	N/A	Conversion to nonbreeding roosting habitat	loss
Breeding habitat— foraging	Loss	Conversion to nonbreeding roosting habitat (portion with bulrush)	Conversion to nonbreeding roosting habitat	Loss	Loss	Loss	Loss	Conversion to nonbreeding roosting habitat	Loss
Breeding habitat— nesting	Loss	N/A	N/A	N/A	Loss	Loss	Loss	Conversion to nonbreeding roosting habitat	Loss
Nonbreeding habitat— foraging ag	Loss	N/A	N/A	N/A	N/A	Loss	Loss	Conversion to nonbreeding roosting habitat	Loss
Nonbreeding habitat— roosting	Loss	Partial loss	No loss	Loss	Loss	Loss	Loss	No loss	Loss
Nonbreeding habitat— foraging	Loss	N/A	Conversion to nonbreeding roosting habitat	Loss	Loss	Loss	Loss	Conversion to nonbreeding roosting habitat	Loss
Western yellow-billed cuckoo									
Breeding habitat	Loss	Loss	Loss	Loss	Loss	No loss	Loss	Loss	Loss
Migratory habitat	Loss	Loss	Loss	Loss	Loss	No loss	Loss	Loss	Loss

Habitat	Suisun					Delta			
	High Tidal Marsh	Middle Tidal Marsh	Low Tidal Marsh	Intertidal Mudflat	Subtidal	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal
White-tailed kite									
Breeding habitat	N/A	N/A	Loss	Loss	Loss	No loss	N/A	Loss	loss
Foraging habitat	No loss	Partial loss	Loss	Loss	Loss	No loss	Loss	Loss	loss
Yellow-breasted chat									
Primary nesting and migratory habitat	Loss	Loss	Loss	Loss	Loss	No loss	Loss	Loss	Loss
Secondary nesting and migratory habitat	Loss	Loss	Loss	Loss	Loss	No loss	Loss	Loss	Loss
Suisun Marsh/upper Yolo Bypass nest and migratory habitat	Loss	Loss	Loss	Loss	Loss	No loss	Loss	Loss	Loss
Reptiles									
Giant garter snake									
Aquatic-tidal	N/A	N/A	N/A	N/A	N/A	Loss	Loss	No loss	No loss
Aquatic-nontidal	N/A	N/A	N/A	N/A	N/A	Loss	Loss	Partial loss	Partial loss
Western pond turtle									
Aquatic habitat	Loss	No loss	No loss	No Loss	No loss	No loss	No loss	No loss	No loss
Upland nesting and overwintering	Loss	Loss	Loss	Loss	Loss	Loss	Loss	Loss	Loss
Upland nesting and overwintering—NHD	Loss	Loss	Loss	Loss	Loss	Loss	Loss	Loss	Loss
Invertebrates									
Valley elderberry longhorn beetle									
Riparian vegetation	Loss	Loss	Loss	Loss	Loss	No loss	Loss	Loss	Loss

Habitat	Suisun					Delta			
	High Tidal Marsh	Middle Tidal Marsh	Low Tidal Marsh	Intertidal Mudflat	Subtidal	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal
Nonriparian channels and grasslands	Loss	Loss	Loss	Loss	Loss	Loss	Loss	Loss	Loss
Plants									
Delta button celery									
All	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Delta mudwort									
All	No loss	No loss	No loss	No loss	Partial, only Subtidal 2 and 3 is loss	No loss	No loss	No loss	Partial, only Subtidal 2 and 3 is loss
Mason's lilaeopsis									
All	No loss	No loss	No loss	No loss	Partial, only Subtidal 2 and 3 is loss	No loss	No loss	No loss	Partial, only Subtidal 2 and 3 is loss
Delta tule pea									
All	No loss	No loss	No loss	Loss	Loss	No loss	Loss	No loss	Loss
Suisun Marsh aster									
All	No loss	No loss	No loss	Loss	Loss	No loss	Loss	No loss	Loss
Side-flowering skullcap									
All	N/A	N/A	N/A	N/A	N/A	Loss	No loss	No loss	Partial, subtidal 2 and 3 are loss
Soft bird's-beak									
All	No loss	No loss	Loss	Loss	Loss	N/A	N/A	N/A	N/A
Suisun thistle									
All	No loss	No loss	Loss	Loss	Loss	N/A	N/A	N/A	N/A

1 Table 5.J-4. Indirect Effect Distances from Covered Activity, Wildlife

Covered Species and Habitat Type	Area of Effect Extending from Disturbance Locations into Modeled Species' Habitats				
	100 Feet	250 Feet	500 Feet	1,300 Feet	2,600 Feet
Mammals					
Riparian brush rabbit		X			
Riparian woodrat		X			
Salt marsh harvest mouse	X				
San Joaquin kit fox ¹		X			
Suisun shrew	X				
Birds					
California black rail			X		
California clapper rail			X		
Greater sandhill crane ²				X	
Least Bell's vireo			X		
Suisun song sparrow ³			X		
Swainson's hawk (foraging habitat) ⁴	X (0 feet)				
Swainson's hawk (nesting sites) ³			X (600 feet)		
Tricolored blackbird (nesting colonies) ³				X	
Tricolored blackbird (foraging habitat) ⁴	X (0 feet)				
Western burrowing owl ⁵			X		
Western yellow-billed cuckoo ^{3,6}			X		
White-tailed kite (nesting sites) ³			X (600 feet)		
White-tailed kite (foraging habitat) ⁴	X (0 feet)				
Yellow-breasted chat ²			X		
Reptiles					
Giant garter snake ⁷		X (200 feet)			
Western pond turtle ⁷		X (200 feet)			
Amphibians					
California red-legged frog ⁷			X		

Covered Species and Habitat Type	Area of Effect Extending from Disturbance Locations into Modeled Species' Habitats				
	100 Feet	250 Feet	500 Feet	1,300 Feet	2,600 Feet
California tiger salamander ⁷			X		
Invertebrates					
Valley elderberry longhorn beetle ⁸	X				
California linderiella ⁹		X			
Conservancy fairy shrimp ⁹		X			
Longhorn fairy shrimp ⁹		X			
Midvalley fairy shrimp ⁹		X			
Vernal pool fairy shrimp ⁹		X			
Vernal pool tadpole shrimp ⁹		X			
<p>¹ This distance applies to all occupied kit fox dens.</p> <p>² A detailed analysis of potential indirect effects on greater sandhill crane is provided in Attachment 5J.D, <i>Indirect Effects of the Construction of the BDCP Conveyance Facility on Sandhill Crane</i>.</p> <p>³ Many covered bird species are sensitive to noise, lighting, and line-of-sight disturbances during the nesting season. For example, construction activity that is within 1,300 feet of a marsh identified as potential tricolored blackbird nesting habitat can result in the loss of this habitat function due to human disturbances and avoidance of the site by tricolored blackbirds. Construction-related activities can also result in the abandonment of nesting sites by tricolored blackbirds, yellow-breasted chats, and other birds if appropriate distances from breeding sites are not maintained.</p> <p>⁴ For some species, habitat use in the immediate vicinity of construction activities is reduced due to long-term, but temporary, disturbances from excavation and related activities, noise, and human presence. For example, tricolored blackbirds, greater sandhill cranes, Swainson's hawks, and white-tailed kites may avoid suitable foraging habitat that is near construction activities.</p> <p>⁵ Buffer distances for burrowing owls are applicable to the breeding and non-breeding seasons.</p> <p>⁶ Yellow-billed cuckoo was detected at one location during 2009. While nesting was not confirmed, this disturbance distance applies to any site found to be occupied by this species.</p> <p>⁷ Habitat function and value for most covered species decreases with proximity to ground disturbances or sources of visual or noise disturbance. For reptiles and amphibians that use upland habitats for nesting or aestivation, ground disturbances distant from aquatic habitats may also have affects. A 500-foot buffer is generally sufficient to avoid direct disturbances to occupied wetland habitats (e.g., ponds, creeks, pools) and most adjacent upland sites; however, where aquatic habitats are found to be occupied by California red-legged frog, California tiger salamander, giant garter snake, or western pond turtle occur, care should be taken to determine the potential for movement corridors that might extend beyond the 500-foot buffer. Where aquatic habitats are found to be occupied by any of these species, the buffer will be expanded to incorporate additional features (e.g., watersheds, drainages, or other possible movement corridors) that have a greater likelihood of supporting occupied upland habitat.</p> <p>⁸ 100 feet is the standard distance recommended by the U.S. Fish and Wildlife Service to avoid direct and indirect effects on elderberry shrubs.</p> <p>⁹ Vernal pool invertebrates can be affected by construction-related runoff into vernal pool habitats. A distance of 250 feet is often used to avoid impacts when there may be a hydrologic connection to the pool; however, potential impacts on occupied pools that are subject to construction-related runoff regardless of the distance should be avoided.</p>					

1 Table 5.J-5. Indirect Effect Distances from Covered Activity, Plants

Covered Species and Habitat Type	Area of Effect Extending from Disturbance Locations into Modeled Species' Habitats				
	100 feet	250 feet	500 feet	1,300 feet	2,600 feet
Plants					
Brittlescale		X			
Heartscale		X			
San Joaquin spearscale		X			
Carquinez goldenbush		X			
Delta button celery		X			
Delta mudwort		X			
Mason's lilaepsis		X			
Delta tule pea		X			
Suisun Marsh aster		X			
Slough thistle		X			
Soft bird's-beak		X			
Suisun thistle		X			
Vernal Pool Plants					
Alkali milk-vetch		X			
Legenere		X			
Heckard's peppergrass		X			
Boggs Lake hedge-hyssop		X			
Dwarf downingia		X			

1 **Table 5.J-6. Near Term Natural Communities Loss by Covered Activity**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}															
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM4 Tidal Natural Communities Restoration													Plan Area Total
		Construction and Inundation ⁸ in Suisun Marsh							Construction and Inundation ⁸ in the Delta						
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	
Tidal Perennial Aquatic	86,263	0	0	1	0	0	0	0	1	1	11	0.00	0.00	0.00	14
Tidal Mudflat ¹⁴	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Tidal Brackish Emergent Wetland ¹⁵	8,501	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Tidal Freshwater Emergent Wetland ¹⁵	8,856	0	0	0	0	0	0	0	1	0	0	0	0	0	1
Valley/Foothill Riparian	17,644	0	0	0	0	0	0	0	0	0	286	6	4	2	298
Grassland	76,315	0	0	1	0	0	0	0	39	3	345	37	16	6	448
Alkali Seasonal Wetland Complex	3,723	0	0	0	0	0	0	0	3	0	10	0	0	0	13
Vernal Pool Complex	11,284	0	0	1	0	0	0	0	24	0	3	0	0	0	28 ^{18,19}
Other Natural Seasonal Wetland	276	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nontidal Freshwater Perennial Emergent Wetland	1,385	0	0	0	0	0	0	0	0	0	38	0	1	0	40
Nontidal Perennial Aquatic	5,489	0	0	0	0	0	0	0	5	2	19	8	0	0	34
Managed Wetland	70,698	88	0	1,569	1,099	1,183	42	0	223	141	1,339	26	7	2	5,718
Inland Dune Scrub	19	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cultivated Lands	481,909	0	0	0	0	0	0	0	482	13	3,494	1,386	432	71	5,878
Total	772,364	88	0	1,572	1,099	1,183	42	0	778	159	5,544	1,465	460	81	12,471

2
3

1 Table 5.J-6. Near Term Natural Communities Loss by Covered Activity (cont'd)

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline/Conveyance Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent ^{4,20}	Permanent Reusable Tunnel Material ⁵	Temporary (Borrow and Spoil) ^{4,6,20}	Temporary ⁴	Permanent ⁷	Temporary ⁷	Permanent ¹¹	Temporary ¹¹	Permanent	Permanent ¹²		Permanent ¹³	Permanent ¹³	Permanent	Permanent ¹³	Permanent ¹⁶	Temporary (Borrow and Spoil) ¹⁶
Tidal Perennial Aquatic	86,263	178	0	0	2,101	8	11	0	0	0	0	0	0	0	200	0	2,112	
Tidal Mudflat ¹⁴	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Tidal Brackish Emergent Wetland ¹⁵	8,501	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Tidal Freshwater Emergent Wetland ¹⁵	8,856	5	1	0	10	6	0	0	0	0	0	0	0	0	12	0	10	
Valley/Foothill Riparian	17,644	16	18	1	29	89	88	0	0	0	0	0	0	0	420	1	116	
Grassland	76,315	211	249	0	158	388	239	0	0	4	0	0	0	13	35	1,349	397	
Alkali Seasonal Wetland Complex	3,723	0	0	0	0 ¹⁷	45	0	0	0	0	0	0	0	0	58	0	0	
Vernal Pool Complex	11,284	15 ¹⁸	0	0	0 ^{17,18,19}	0	0	0	0	0	0	0	0	0	43	0	0	
Other Natural Seasonal Wetland	276	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Nontidal Freshwater Perennial Emergent Wetland	1,385	1	1	0	5	25	1	0	0	0	0	0	0	0	67	0	6	
Nontidal Perennial Aquatic	5,489	2	55	0	7	24	12	0	0	0	0	0	0	0	115	0	18	
Managed Wetland	70,698	7	0	0	28	24	44	0	0	0	0	0	0	0	5,750	0	72	
Inland Dune Scrub	19	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Cultivated Lands	481,909	1,448	3,140	199	1,196	629	363	0	0	10	0	1,140	700	0	12,945	199	1,559	
Total	772,364	1,885	3,465	200	3,533	1,238	757	0	0	14	0	1,140	700	13	35	20,961	200	4,290

N/A = Not available.

¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, *Covered Activities and Associated Federal Actions*) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Nonfederal Actions.

² Existing habitat and habitat loss are estimated using natural community models created from detailed vegetation mapping. See Chapter 2, Section 2.3 for a complete description of mapping methods. Effects on natural communities will be tracked during implementation through on-the-ground surveys performed by qualified biologists.

³ See Table 5.J.1, *Quantitative Effects Analysis Methods and Assumptions*, in Appendix 5.J, *Effects on Natural Communities, Wildlife, and Plants*, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, *Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat*, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.

⁴ Permanent and temporary effects assessed under CM1 Water Facilities and Operation are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, *Covered Activities and Associated Federal Actions*, Section 4.1.3.1, Tunnel/Pipeline Facility Construction and Operations, for a complete description of all activities assessed under CM1.

⁵ This represents the maximum area potentially necessary for storing reusable tunnel material. This material will likely be moved to other sites for use in levee build-up and restoration, and the affected area will likely be restored. While this effect is categorized as permanent, because there is no assurance that the material will eventually be moved, the effect will likely be temporary. Furthermore, the amount of storage area needed for reusable tunnel material is flexible (based on height of storage piles and other factors) and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.

⁶ Borrow/Spoil Area: Borrow: a location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil. While these impacts are considered "temporary", because affected lands will be restored when conveyance facility construction is complete, for the purposes of determining net effects, impacts are considered "permanent".

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline/Conveyance Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction				
		Permanent ^{4,20}	Permanent Reusable Tunnel Material ⁵	Temporary (Borrow and Spoil) ^{4,6,20}	Temporary ⁴	Permanent ⁷	Temporary ⁷	Permanent ¹¹	Temporary ¹¹	Permanent	Permanent ¹²	Permanent ¹³	Permanent ¹³	Permanent	Permanent ¹³	Permanent ¹³	Permanent ¹⁶	Temporary (Borrow and Spoil) ¹⁶	Temporary ¹⁶
<p>⁷ Permanent and temporary effects assessed under CM2 Yolo Bypass Fisheries Enhancement include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.</p> <p>⁸ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>⁹ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>¹⁰ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>¹¹ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>¹² Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>¹³ Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>¹⁴ Tidal mudflat features were not mapped in the BDCP vegetation layer.</p> <p>¹⁵ Effects on tidal wetland communities are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below MLLW in Suisun and MLLW + 1 ft. in the rest of the Delta. See Table 5.J-1 for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>¹⁶ Totals may not sum due to rounding.</p> <p>¹⁷ Loss reduced to zero. Although the temporary powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools be avoided during temporary powerline installation.</p> <p>¹⁸ Of the 11,284 acres of vernal pool complex natural community, 2,576 acres are considered “degraded”. Of the original (some impacts subsequently reduced, see footnotes 17 and 19) 15 acres of permanent loss (CM1), 0 acres of temporary loss (CM1), and 28 acres of permanent loss (CM4), 7 acres, 2 acres, and 370 acres of loss are to degraded vernal pool complex, respectively.</p> <p>¹⁹ Total permanent loss reduced from 201 acres (CM4) to 28 acres. This reduction is based on a 10-acre cap for total loss of wetted acres, assuming 15% density of vernal pools in the area affected. Acreage of vernal pool complex loss may be higher if actual vernal pool density is lower. The maximum acreage loss is based on loss of wetted acres and not total vernal pool complex acreage.</p> <p>²⁰ Current proposed transmission line alignment extends outside the Plan Area, although final alignment is unknown. Acreage loss associated with transmission line construction outside the Plan Area is included in this column. Plan Area will be adjusted if needed for final plan when transmission line alignment is further designed.</p>																			

1
2

1 **Table 5.J-7. Early Long-Term Natural Communities Loss by Covered Activity**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}															
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM4 Tidal Natural Communities Restoration													Plan Area Total
		Construction and Inundation ⁸ in Suisun Marsh							Construction and Inundation ⁸ in the Delta						
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	
Tidal Perennial Aquatic	86,263	0	0	1	0	0	0	0	1	1	13	0	0	0	16
Tidal Mudflat ¹⁴	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Tidal Brackish Emergent Wetland ¹⁵	8,501	0	0	0	1	0	0	0	0	0	0	0	0	0	1
Tidal Freshwater Emergent Wetland ¹⁵	8,856	0	0	0	0	0	0	0	1	0	0	0	0	0	1
Valley/Foothill Riparian	17,644	0	0	0	0	0	0	0	0	0	389	7	5	2	403
Grassland	76,315	0	0	1	0	0	0	0	32	3	632	39	17	6	732
Alkali Seasonal Wetland Complex	3,723	0	0	0	0	0	0	0	0	0	13	0	0	0	13
Vernal Pool Complex	11,284	0	0	0	0	0	0	0	26	0	26	0	0	0	53 ^{18,19}
Other Natural Seasonal Wetland	276	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nontidal Freshwater Perennial Emergent Wetland	1,385	0	0	0	0	0	0	0	0	0	50	0	1	0	51
Nontidal Perennial Aquatic	5,489	0	0	0	0	0	0	0	6	0	51	10	1	0	68
Managed Wetland	70,698	56	112	1,783	1,628	1,765	69	0	232	0	1,479	161	14	2	7,301
Inland Dune Scrub	19	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cultivated Lands	481,909	0	0	0	0	0	0	0	823	207	7,104	2,365	829	95	11,423
Total	772,364	57	113	1,786	1,630	1,765	69	0	1,121	211	9,757	2,584	866	105	20,062

2
3

1 **Table 5.J-7. Early Long-Term Natural Communities Loss by Covered Activity (cont'd)**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline/Conveyance Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent ^{4,20}	Permanent Reusable Tunnel Material ⁵	Temporary (Borrow and Spoil) ^{6,6,20}	Temporary ⁴	Permanent ⁷	Temporary ⁷	Permanent ¹¹	Temporary ¹¹	Permanent	Permanent ¹²		Permanent ¹³	Permanent ¹³	Permanent	Permanent ¹³	Permanent ¹⁶	Temporary (Borrow and Spoil) ¹⁶
Tidal Perennial Aquatic	86,263	178	0	0	2,101	8	11	0	0	0	0	0	0	0	0	202	0	2,112
Tidal Mudflat ¹⁴	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Tidal Brackish Emergent Wetland ¹⁵	8,501	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
Tidal Freshwater Emergent Wetland ¹⁵	8,856	5	1	0	10	6	0	0	0	0	0	0	0	0	0	12	0	10
Valley/Foothill Riparian	17,644	16	18	1	29	89	88	6	6	0	0	0	0	0	0	532	1	123
Grassland	76,315	211	249	0	158	388	239	11	12	7	0	0	0	20	35	1,653	0	409
Alkali Seasonal Wetland Complex	3,723	0	0	0	0 ¹⁷	45	0	0	0	0	0	0	0	0	0	59	0	0
Vernal Pool Complex	11,284	15 ¹⁸	0	0	0 ^{17,18,19}	0	0	0	0	0	0	0	0	0	0	68	0	0
Other Natural Seasonal Wetland	276	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nontidal Freshwater Perennial Emergent Wetland	1,385	1	1	0	5	25	1	0	0	0	0	0	0	0	0	78	0	6
Nontidal Perennial Aquatic	5,489	2	55	0	7	24	12	19	8	0	0	0	0	0	0	168	0	27
Managed Wetland	70,698	7	0	0	28	24	44	0	0	0	0	0	0	0	0	7,332	0	72
Inland Dune Scrub	19	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cultivated Lands	481,909	1,448	3,140	199	1,196	629	363	252	153	19	0	1,480	1,000	0	0	19,392	199	1,711
Total	772,364	1,885	3,465	200	3,533	1,238	757	288	180	26	0	1,480	1,000	20	35	29,498	200	4,470

N/A = Not available.

¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Nonfederal Actions.

² Existing habitat and habitat loss are estimated using natural community models created from detailed vegetation mapping, See Chapter 2, Existing Ecological Conditions, Section 2.3, for a complete description of mapping methods. Effects on natural communities will be tracked during implementation through on-the-ground surveys performed by qualified biologists.

³ See Table 5.J.1, *Quantitative Effects Analysis Methods and Assumptions*, in Appendix 5.J, *Effects on Natural Communities, Wildlife, and Plants*, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, *Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat*, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.

⁴ Permanent and temporary effects assessed under CM1 Water Facilities and Operation are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Covered Activities and Associated Federal Actions, Section 4.1.3.1, Tunnel/Pipeline Facility Construction and Operations, for a complete description of all activities assessed under CM1.

⁵ This represents the maximum area potentially necessary for storing reusable tunnel material. This material will likely be moved to other sites for use in levee build-up and restoration, and the affected area will likely be restored. While this effect is categorized as

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline/Conveyance Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent ^{4,20}	Permanent Reusable Tunnel Material ⁵	Temporary (Borrow and Spoil) ^{4,6,20}	Temporary ⁴	Permanent ⁷	Temporary ⁷	Permanent ¹¹	Temporary ¹¹	Permanent	Permanent ¹²	Permanent ¹³	Permanent ¹³	Permanent	Permanent ¹³	Permanent ¹³	Permanent ¹⁶	Temporary (Borrow and Spoil) ¹⁶
<p>permanent, because there is no assurance that the material will eventually be moved, the effect will likely be temporary. Furthermore, the amount of storage area needed for reusable tunnel material is flexible (based on height of storage piles and other factors) and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.</p> <p>⁶ Borrow/Spoil Area: Borrow: a location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil. While these impacts are considered “temporary”, because affected lands will be restored when conveyance facility construction is complete, for the purposes of determining net effects, impacts are considered “permanent”.</p> <p>⁷ Permanent and temporary effects assessed under CM2 Yolo Bypass Fisheries Enhancement include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.</p> <p>⁸ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>⁹ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>¹⁰ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>¹¹ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>¹² Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>¹³ Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>¹⁴ Tidal mudflat features were not mapped in the BDCP vegetation layer.</p> <p>¹⁵ Effects on tidal wetland communities are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below MLLW in Suisun and MLLW + 1 ft. in the rest of the Delta. See Table 5.J-1 for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>¹⁶ Totals may not sum due to rounding.</p> <p>¹⁷ Loss reduced to zero. Although the temporary powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools be avoided during temporary powerline installation.</p> <p>¹⁸ Of the 11,284 acres of vernal pool complex natural community, 2,576 acres are considered “degraded”. Of the original (some impacts subsequently reduced, see footnotes 17 and 19) 15 acres of permanent loss (CM1), 0 acres of temporary loss (CM1), and 28 acres of permanent loss (CM4), 7 acres, 2 acres, and 370 acres of loss are to degraded vernal pool complex, respectively.</p> <p>¹⁹ Total permanent loss reduced from 201 acres (CM4) to 28 acres. This reduction is based on a 10-acre cap for total loss of wetted acres, assuming 15% density of vernal pools in the area affected. Acreage of vernal pool complex loss may be higher if actual vernal pool density is lower. The maximum acreage loss is based on loss of wetted acres and not total vernal pool complex acreage.</p> <p>²⁰ Current proposed transmission line alignment extends outside the Plan Area, although final alignment is unknown. Acreage loss associated with transmission line construction outside the Plan Area is included in this column. Plan Area will be adjusted if needed for final plan when transmission line alignment is further designed.</p>																		

1
2

1 **Table 5.J-8. Late Long-Term Natural Communities Loss by Covered Activity**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}															
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM4 Tidal Natural Communities Restoration													Plan Area Total
		Construction and Inundation ⁸ in Suisun Marsh						Construction and Inundation ⁸ in the Delta						Permanent ^{8,9,10}	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2		
Tidal Perennial Aquatic	86,263	0	0	1	0	0	0	0	1	1	14	0	0	0	18
Tidal Mudflat ¹⁴	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
Tidal Brackish Emergent Wetland ¹⁵	8,501	0	0	0	1	0	0	0	0	0	0	0	0	0	1
Tidal Freshwater Emergent Wetland ¹⁵	8,856	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Valley/Foothill Riparian	17,644	0	0	0	0	0	0	0	0	0	490	25	28	8	552
Grassland	76,315	1	0	1	0	1	0	0	74	3	881	61	65	35	1,122
Alkali Seasonal Wetland Complex	3,723	0	0	0	0	0	0	0	2	0	25	0	0	0	27
Vernal Pool Complex	11,284 ¹⁸	0	0	1	0	0	0	0	9	0	41	1	0	0	52 ^{18,19}
Other Natural Seasonal Wetland	276	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nontidal Freshwater Perennial Emergent Wetland	1,385	0	0	0	0	0	0	0	7	0	81	7	3	2	99
Nontidal Perennial Aquatic	5,489	0	0	0	0	0	0	0	25	0	94	23	38	10	189
Managed Wetland	70,698	68	71	2,499	1,756	6,493	644	1	137	0	1,882	157	30	9	13,746
Inland Dune Scrub	19	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cultivated Lands	481,909	2	0	0	0	0	0	0	1,437	1	18,707	7,316	8,982	3,120	39,565
Total	772,364	71	72	2,503	1,757	6,494	644	1	1,693	4	22,214	7,589	9,146	3,184	55,373

2
3

1 **Table 5.J-8. Late Long-Term Natural Communities Loss by Covered Activity (cont'd)**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline/Conveyance Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction				
		Permanent ^{4,20}	Permanent Reusable Tunnel Material ⁵	Temporary (Borrow and Spoil) ^{6,6,20}	Temporary ⁴	Permanent ⁷	Temporary ⁷	Permanent ¹¹	Temporary ¹¹	Permanent	Permanent ¹²		Permanent ¹³	Permanent ¹³	Permanent	Permanent ¹³	Permanent ¹⁶	Temporary (Borrow and Spoil) ¹⁶	Temporary ¹⁶
Tidal Perennial Aquatic	86,263	178	0	0	2,101	8	11	2	5	0	0	0	0	0	0	207	0	2,116	
Tidal Mudflat ¹⁴	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Tidal Brackish Emergent Wetland ¹⁵	8,501	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	
Tidal Freshwater Emergent Wetland ¹⁵	8,856	5	1	0	10	6	0	1	1	0	0	0	0	0	0	13	0	11	
Valley/Foothill Riparian	17,644	16	18	1	29	89	88	43	35	0	0	0	0	0	0	717	1	151	
Grassland	76,315	211	249	0	158	388	239	51	34	11	399	0	0	50	35	2,517	0	431	
Alkali Seasonal Wetland Complex	3,723	0	0	0	0 ¹⁷	45	0	0	0	0	0	0	0	0	0	72	0	0	
Vernal Pool Complex	11,284 ¹⁸	15 ¹⁸	0	0	0 ^{17,18,19}	0	0	0	0	0	0	0	0	0	0	67	0	0	
Other Natural Seasonal Wetland	276	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Nontidal Freshwater Perennial Emergent Wetland	1,385	1	1	0	5	25	1	0	0	0	0	0	0	0	0	127	0	6	
Nontidal Perennial Aquatic	5,489	2	55	0	7	24	12	28	16	0	0	0	0	0	0	299	0	34	
Managed Wetland	70,698	7	0	0	28	24	44	0	0	0	0	0	0	0	0	13,778	0	72	
Inland Dune Scrub	19	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Cultivated Lands	481,909	1,448	3,140	199	1,196	629	363	2,087	1,194	960	3,593	2,000	1,950	0	0	55,372	199	2,753	
Total	772,364	1,885	3,465	200	3,533	1,238	757	2,212	1,285	971	3,991	2,000	1,950	50	35	73,170	200	5,575	

N/A = Not available.

¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Nonfederal Actions.

² Existing habitat and habitat loss are estimated using natural community models created from detailed vegetation mapping, See Chapter 2, Existing Ecological Conditions, Section 2.3, for a complete description of mapping methods. Effects on natural communities will be tracked during implementation through on-the-ground surveys performed by qualified biologists.

³ See Table 5.J.1, *Quantitative Effects Analysis Methods and Assumptions*, in Appendix 5.J, *Effects on Natural Communities, Wildlife, and Plants*, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, *Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat*, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.

⁴ Permanent and temporary effects assessed under CM1 Water Facilities and Operation are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Covered Activities and Associated Federal Actions, Section 4.1.3.1, Tunnel/Pipeline Facility Construction and Operations, for a complete description of all activities assessed under CM1.

⁵ This represents the maximum area potentially necessary for storing reusable tunnel material. This material will likely be moved to other sites for use in levee build-up and restoration, and the affected area will likely be restored. While this effect is categorized as

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Natural Community	Total Existing Modeled Habitat in the Plan Area ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline/Conveyance Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent ^{4,20}	Permanent Reusable Tunnel Material ⁵	Temporary (Borrow and Spoil) ^{4,6,20}	Temporary ⁴	Permanent ⁷	Temporary ⁷	Permanent ¹¹	Temporary ¹¹	Permanent	Permanent ¹²	Permanent ¹³	Permanent ¹³	Permanent	Permanent ¹³	Permanent ¹³	Permanent ¹⁶	Temporary (Borrow and Spoil) ¹⁶
<p>permanent, because there is no assurance that the material will eventually be moved, the effect will likely be temporary. Furthermore, the amount of storage area needed for reusable tunnel material is flexible (based on height of storage piles and other factors) and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.</p> <p>⁶ Borrow/Spoil Area: Borrow: a location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil. While these impacts are considered “temporary”, because affected lands will be restored when conveyance facility construction is complete, for the purposes of determining net effects, impacts are considered “permanent”.</p> <p>⁷ Permanent and temporary effects assessed under CM2 Yolo Bypass Fisheries Enhancement include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.</p> <p>⁸ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>⁹ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>¹⁰ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>¹¹ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>¹² Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>¹³ Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>¹⁴ Tidal mudflat features were not mapped in the BDCP vegetation layer.</p> <p>¹⁵ Effects on tidal wetland communities are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below MLLW in Suisun and MLLW + 1 ft. in the rest of the Delta. See Table 5.J-1 for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>¹⁶ Totals may not sum due to rounding.</p> <p>¹⁷ Loss reduced to zero. Although the temporary powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools be avoided during temporary powerline installation.</p> <p>¹⁸ Of the 11,284 acres of vernal pool complex natural community, 2,576 acres are considered “degraded”. Of the original (some impacts subsequently reduced, see footnotes 17 and 19) 15 acres of permanent loss (CM1), 0 acres of temporary loss (CM1), and 28 acres of permanent loss (CM4), 7 acres, 2 acres, and 370 acres of loss are to degraded vernal pool complex, respectively.</p> <p>¹⁹ Total permanent loss reduced from 201 acres (CM4) to 28 acres. This reduction is based on a 10-acre cap for total loss of wetted acres, assuming 15% density of vernal pools in the area affected. Acreage of vernal pool complex loss may be higher if actual vernal pool density is lower. The maximum acreage loss is based on loss of wetted acres and not total vernal pool complex acreage.</p> <p>²⁰ Current proposed transmission line alignment extends outside the Plan Area, although final alignment is unknown. Acreage loss associated with transmission line construction outside the Plan Area is included in this column. Plan Area will be adjusted if needed for final plan when transmission line alignment is further designed.</p>																		

1
2

1 Table 5.J-9. Near Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity

Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}														Plan Area Total ⁷	
		CM4 Tidal Natural Communities Restoration														Permanent (Acres) ^{7,8,9}	Conversion (Acres)
		Suisun Marsh							Delta								
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3			
Mammals																	
Riparian brush rabbit																	
<i>Riparian habitat</i>	2,909	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Grassland habitat</i>	3,103	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	6,011	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Riparian woodrat																	
<i>Habitat</i>	2,166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	2,166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Salt marsh harvest mouse																	
<i>Tidal brackish emergent wetland primary</i>	3,641	0	0	64	0	0	0	0	0	0	0	0	0	0	0	0	64
<i>Tidal brackish emergent wetland secondary</i>	2,718	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland secondary</i>	749	1	7	0	0	0	0	0	0	0	0	0	0	0	0	0 ¹⁵	8
<i>Managed wetland—wetland primary, low long-term conservation value</i>	21,891	30	0	534	577	770	3	0	0	0	0	0	0	0	0	1,349	564
<i>Managed wetland—wetland secondary, low long-term conservation value</i>	2,800	11	0	229	50	23	0	0	0	0	0	0	0	0	0	74	241
<i>Managed wetland—upland, low long-term conservation value</i>	3,787	8	0	64	70	24	0	0	0	0	0	0	0	0	94	71	
Total	35,588	49	7	892	697	817	3	0	0	0	0	0	0	0	0	1,517	948
San Joaquin kit fox																	
<i>Breeding, foraging, and dispersal habitat</i>	5,327	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	5,327	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Suisun shrew																	
<i>Primary habitat</i>	3,128	0	0	58	0	0	0	0	0	0	0	0	0	0	0	58	0
<i>Secondary habitat</i>	4,387	10	5	0	22	9	0	0	0	0	0	0	0	0	0	31 ¹⁵	15
Total	7,515	10	5	58	22	9	0	0	0	0	0	0	0	0	0	89	15
Birds																	
California black rail																	
<i>Primary habitat</i>	7,467	0	0	69	0	0	0	0	0	0	0	0	0	1	0	2	69
<i>Secondary habitat</i>	17,915	49	0	0	402	532	2	0	0	0	1	0	0	0	0	936 ¹⁵	50
Total	25,382	49	0	69	402	532	2	0	0	0	1	0	1	0	938	120	
California clapper rail¹³																	
<i>Primary habitat</i>	296	0	0	26	0	0	0	0	0	0	0	0	0	0	0	0	26
<i>Secondary habitat</i>	6,420	2	5	0	0	0	0	0	0	0	0	0	0	0	0	0 ¹⁵	7
Total	6,716	2	5	26	0	0	0	0	0	0	0	0	0	0	0	0	33

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM4 Tidal Natural Communities Restoration													Plan Area Total ⁷	
		Suisun Marsh						Delta						Permanent (Acres) ^{7,8,9}	Conversion (Acres)	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2			Subtidal 3
California least tern																
<i>Nesting and Migratory Habitat</i>	86,263	0	0	1	0	0	0	0	1	1	11	12	4	0	30	0
Total	86,263	0	0	1	0	0	0	0	1	1	11	12	4	0	30	0
Greater sandhill crane																
<i>Roosting and foraging - Permanent</i>	7,340	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Roosting and foraging - Temporary</i>	16,522	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Foraging</i>	162,164	0	0	0	0	0	0	0	2	0	709	852	363	26	1,951	0
Total	186,025	0	0	0	0	0	0	0	2	0	709	852	363	26	1,951	0
Least Bell's vireo																
<i>Migratory and breeding</i>	14,528	1	0	3	1	1	0	0	0	0	282	6	4	2	299	0
Total	14,528	1	0	3	1	1	0	0	0	0	282	6	4	2	299	0
Suisun song sparrow																
<i>Primary habitat</i>	3,722	0	0	54	0	0	0	0	0	0	0	0	0	0	0	54
<i>Secondary habitat</i>	23,986	53	5	0	432	605	2	0	0	0	0	0	0	0	1,040 ¹⁵	58
Total	27,707	53	5	54	432	605	2	0	0	0	0	0	0	0	1,040	112
Swainson's hawk																
<i>Foraging habitat</i>	470,324	27	29	192	59	43	0	0	773	152	3,742	882	352	27	6,278	0
<i>Nesting habitat</i>	9,796	0	0	4	1	0	0	0	0	0	164	2	1	1	173	0
Total	480,120	27	29	196	60	43	0	0	773	152	3,906	884	354	27	6,451	0
Tricolored blackbird																
<i>Breeding habitat-ag foraging</i>	100,198	0	0	0	0	0	0	0	106	0	79	0	0	0	106	79
<i>Breeding habitat-foraging</i>	58,181	19	2	286	153	99	0	0	1	0	1	0	0	0	272	288
<i>Breeding habitat-nesting</i>	1,741	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Nonbreeding habitat-foraging ag</i>	194,251	0	0	0	0	0	0	0	332	8	2,065	488	312	19	1,159	2,065
<i>Nonbreeding habitat-roosting</i>	28,066	36	1	0	230	171	1	0	116	0	0	4	2	1	562	0
<i>Nonbreeding habitat-foraging</i>	34,308	0	0	0	0	0	0	0	41	3	355	37	16	6	104	355
Total	416,745	54	2	286	383	270	2	0	595	11	2,499	529	331	26	2,203	2,787
Western burrowing owl																
<i>High-value habitat</i>	149,783	17	11	122	98	52	0	0	623	5	2,037	143	157	33	3,297	0
<i>Low-value habitat</i>	251,767	1	17	23	3	1	0	0	141	148	1,505	397	64	2	2,300	0
Total	401,550	17	28	145	100	53	0	0	764	152	3,541	540	220	35	5,597	0
Western yellow-billed cuckoo																
<i>Breeding habitat</i>	1,970	0	0	0	0	0	0	0	0	0	1	1	1	0	3	0
<i>Migratory habitat</i>	10,425	0	0	0	0	0	0	0	0	0	216	2	1	1	221	0
Total	12,395	0	0	0	0	0	0	0	0	0	217	3	2	1	224	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM4 Tidal Natural Communities Restoration													Plan Area Total ⁷	
		Suisun Marsh						Delta						Permanent (Acres) ^{7,8,9}	Conversion (Acres)	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2			Subtidal 3
White-tailed kite																
<i>Breeding habitat</i>	14,069	0	0	4	1	0	0	0	0	0	218	3	2	1	230	0
<i>Foraging habitat</i>	500,365	0	32	986	733	931	3	0	0	152	3,578	870	353	27	7,667	0
Total	514,434	0	32	991	733	931	3	0	0	153	3,796	873	355	28	7,896	0
Yellow-breasted chat																
<i>Primary nesting and migratory habitat</i>	8,178	0	0	0	0	0	0	0	0	0	83	1	2	1	87	0
<i>Secondary nesting and migratory habitat</i>	5,528	0	0	0	0	0	0	0	0	0	199	5	2	1	206	0
<i>Suisun Marsh/Upper Yolo Bypass nest and migratory habitat</i>	841	1	0	3	1	1	0	0	0	0	0	0	0	0	5	0
Total	14,547	1	0	3	1	1	0	0	0	0	282	6	4	2	299	0
Reptiles																
Giant garter snake																
<i>Aquatic - tidal</i>	12,097	0	0	0	0	0	0	0	2	0	0	0	0	0	2	0
<i>Aquatic - nontidal¹⁹</i>	19,027	0	0	0	0	0	0	0	5	2	89	9	3	0	109	0
<i>Upland-high</i>	21,581	0	0	0	0	0	0	0	43	1	239	32	32	17	364	0
<i>Upland-moderate</i>	25,407	0	0	0	0	0	0	0	70	43	552	26	7	1	700	0
<i>Upland-low</i>	5,683	0	0	0	0	0	0	0	11	0	116	1	1	1	129	0
Total	83,796	0	0	0	0	0	0	0	131	48	996	69	43	20	1,305	0
<i>Aquatic breeding, foraging, and movement (miles)</i>	2,784	0	0	0	0	0	0	0	3	1	30	5	3	1	44	0
Western pond turtle																
<i>Aquatic habitat¹⁰</i>	81,588	45	0	0	0	0	0	0	0	0	0	0	0	0	45	0
<i>Upland nesting and overwintering habitat</i>	16,043	2	1	48	41	16	0	0	2	0	24	2	0	0	136	0
<i>Upland nesting and overwintering habitat-NHD</i>	12,615	5	5	12	7	2	0	0	9	1	88	9	5	2	144	0
Total	110,245	52	6	60	48	18	0	0	10	1	111	11	6	2	326	0
<i>Aquatic habitat linear (miles) - NHD</i>	1,418	0	0	1	1	1	0	0	1	0	14	3	2	0	24	0
Amphibians																
California red-legged frog																
<i>Aquatic habitat</i>	159	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland cover and dispersal habitat</i>	7,766	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	7,925	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Aquatic habitat (miles)</i>	30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
California tiger salamander																
<i>Aquatic breeding habitat</i>	7,845	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Terrestrial cover and aestivation</i>	28,173	0	0	0	0	0	0	0	182	0	21	0	0	0	203	0
Total	36,018	0	0	0	0	0	0	0	182	0	21	0	0	0	203	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM4 Tidal Natural Communities Restoration													Plan Area Total ⁷	
		Suisun Marsh						Delta						Permanent (Acres) ^{7,8,9}	Conversion (Acres)	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2			Subtidal 3
Invertebrates																
Valley elderberry longhorn beetle																
<i>Riparian vegetation</i>	17,464	0	0	0	0	0	0	0	0	0	286	6	4	2	298	0
<i>Nonriparian channels and grasslands</i>	16,585	1	0	9	4	0	0	0	7	0	59	13	6	1	100	0
Total	34,048	1	0	9	4	0	0	0	7	1	345	19	10	2	398	0
California linderiella																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Conservancy fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Longhorn fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Midvalley fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Vernal pool fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Vernal pool tadpole shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28	0

1
2

1 Table 5.J-9. Near Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity (cont'd)

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			
Mammals																		
Riparian brush rabbit																		
<i>Riparian habitat</i>	2,909	3	0	0	1	0	0	0	0	0	0	0	0	0	0	3	0	1
<i>Grassland habitat</i>	3,103	124	0	0	54	0	0	0	0	0	0	0	0	0	0	124	0	54
Total	6,011	127	0	0	54	0	0	0	0	0	0	0	0	0	0	127	0	54
Riparian woodrat																		
<i>Habitat</i>	2,166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	2,166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Salt marsh harvest mouse																		
<i>Tidal brackish emergent wetland primary</i>	3,641	0	0	0	0	0	0	0	0	0	0	0	0	0	0	64	0	0
<i>Tidal brackish emergent wetland secondary</i>	2,718	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland secondary</i>	749	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Managed wetland—wetland primary, low long-term conservation value</i>	21,891	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1,913	0	0
<i>Managed wetland—wetland secondary, low long-term conservation value</i>	2,800	0	0	0	0	0	0	0	0	0	0	0	0	0	0	315	0	0
<i>Managed wetland—upland, low long-term conservation value</i>	3,787	0	0	0	0	0	0	0	0	0	0	0	0	0	0	165	0	0
Total	35,588	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2,465	0	0
San Joaquin kit fox																		
<i>Breeding, foraging, and dispersal habitat</i>	5,327	155	52	0	103	0	0	0	0	0	0	0	0	3	0	210	0	103
Total	5,327	155	52	0	103	0	0	0	0	0	0	0	0	3	0	210	0	103
Suisun shrew																		
<i>Primary habitat</i>	3,128	0	0	0	0	0	0	0	0	0	0	0	0	0	0	58	0	0
<i>Secondary habitat</i>	4,387	0	0	0	0	0	0	0	0	0	0	0	0	0	0	47 ¹⁵	0	0
Total	7,515	0	0	0	0	0	0	0	0	0	0	0	0	0	0	105	0	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			
Birds																		
California black rail																		
Primary habitat	7,467	0	0	0	18	5	0	0	0	0	0	0	0	0	0	76	0	18
Secondary habitat	17,915	0	0	0	0	0	0	0	0	0	0	0	0	0	0	986 ¹⁵	0	0
Total	25,382	0	0	0	18	5	0	0	0	0	0	0	0	0	0	1,062	0	18
California clapper rail¹³																		
Primary habitat	296	0	0	0	0	0	0	0	0	0	0	0	0	0	0	26	0	0
Secondary habitat	6,420	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7 ¹⁵	0	0
Total	6,716	0	0	0	0	0	0	0	0	0	0	0	0	0	0	33	0	0
California least tern																		
Nesting and Migratory Habitat	86,263	178	0	0	2,101	8	11	0	0	0	0	0	0	0	0	216	0	2,112
Total	86,263	178	0	0	2,101	8	11	0	0	0	0	0	0	0	0	216	0	2,112
Greater sandhill crane																		
Roosting and foraging - Permanent	7,340	0	0	0	8	0	0	0	0	0	0	0	0	0	0	0	0	8
Roosting and foraging - Temporary	16,522	0 ¹⁴	0	0	16	0	0	0	0	0	0	0	0	0	0	0	0	16
Foraging	162,164	352	2,347	183	778	0	0	0	0	0	0	257	567	1	0	5,474	183	778
Total	186,025	352	2,347	183	802	0	0	0	0	0	0	257	567	1	0	5,474	183	802
Least Bell's vireo																		
Migratory and breeding	14,528	11	18	1	22	83	88	0	0	0	0	0	0	0	0	411	1	110
Total	14,528	11	18	1	22	83	88	0	0	0	0	0	0	0	0	411	1	110
Suisun song sparrow																		
Primary habitat	3,722	0	0	0	0	0	0	0	0	0	0	0	0	0	0	54	0	0
Secondary habitat	23,986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1,097 ¹⁵	0	0
Total	27,707	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1,151	0	0
Swainson's hawk																		
Foraging habitat	470,324	1,100	3,235	183	1,113	996	504	0	0	13	0	1,054	527	13	35	13,251	183	1,617
Nesting habitat	9,796	8	10	0	18	79	54	0	0	0	0	0	0	0	0	270	0	72
Total	480,120	1,108	3,245	183	1,131	1,075	558	0	0	13	0	1,054	527	13	35	13,521	183	1,689
Tricolored blackbird																		
Breeding habitat-ag foraging	100,198	634	795	81	148	477	84	0	0	2	0	867	126	0	0	3,086	81	232
Breeding habitat-foraging	58,181	161	52	0	114	105	155	0	0	4	0	0	0	13	35	930	0	268

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			
<i>Breeding habitat-nesting</i>	1,741	4	0	1	2	13	75	0	0	0	0	0	0	0	0	17	1	77
<i>Nonbreeding habitat-foraging ag</i>	194,251	203	2,124	0	575	0	54	0	0	7	0	120	397	0	0	6,074	0	628
<i>Nonbreeding habitat-roosting</i>	28,066	7	12	0	20	8	0	0	0	0	0	0	0	0	0	590	0	20
<i>Nonbreeding habitat-foraging</i>	34,308	48	197	0	47	0	0	0	0	0	0	0	0	0	0	704	0	47
Total	416,745	1,057	3,180	82	905	603	367	0	0	13	0	987	523	13	35	11,401	82	1,273
Western burrowing owl																		
<i>High-value habitat</i>	149,783	340	541	0	351	882	245	0	0	4	0	206	63	13	35	5,381	0	596
<i>Low-value habitat</i>	251,767	689	2,324	101	588	98	144	0	0	9	0	749	371	0	0	6,540	101	732
Total	401,550	1,030	2,864	102	939	979	389	0	0	13	0	955	434	13	35	11,921	102	1,328
Western yellow-billed cuckoo																		
<i>Breeding habitat</i>	1,970	3	6	0	1	26	5	0	0	0	0	0	0	0	0	38	0	5
<i>Migratory habitat</i>	10,425	4	10	0	18	57	83	0	0	0	0	0	0	0	0	292	0	101
Total	12,395	7	16	0	19	83	88	0	0	0	0	0	0	0	0	330	0	106
White-tailed kite																		
<i>Breeding habitat</i>	14,069	10	16	0	23	82	88	0	0	0	0	0	0	0	0	338	0	110
<i>Foraging habitat</i>	500,365	1,100	3,239	183	1,112	1,008	516	0	0	13	0	0	0	13	35	13,075	183	1,629
Total	514,434	1,111	3,255	183	1,135	1,090	604	0	0	13	0	0	0	13	35	13,413	183	1,739
Yellow-breasted chat																		
<i>Primary nesting and migratory habitat</i>	8,178	7	10	0	6	9	58	0	0	0	0	0	0	0	0	113	0	64
<i>Secondary nesting and migratory habitat</i>	5,528	3	8	1	16	3	0	0	0	0	0	0	0	0	0	220	1	16
<i>Suisun Marsh/Upper Yolo Bypass nest and migratory habitat</i>	841	0	0	0	0	71	29	0	0	0	0	0	0	0	0	76	0	29
Total	14,547	10	18	1	22	83	88	0	0	0	0	0	0	0	0	410	1	110
Reptiles																		
Giant garter snake																		
<i>Aquatic - tidal</i>	12,097	16	1	0	55	9	2	0	0	0	0	0	0	0	0	28	0	57
<i>Aquatic - nontidal¹⁹</i>	19,027	10	56	0	13	59	13	0	0	0	0	0	0	0	0	235	0	26
<i>Upland-high</i>	21,581	66	106	0	48	178	158	0	0	0	0	0	0	0	0	715	0	206
<i>Upland-moderate</i>	25,407	167	54	0	135	60	61	0	0	0	0	0	0	0	35	1,017	0	196

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			
<i>Upland-low</i>	5,683	14	4	0	5	1	0	0	0	0	0	0	0	0	0	148	0	5
Total	83,796	274	222	0	257	306	234	0	0	0	0	0	0	0	35	2,142	0	491
<i>Aquatic breeding, foraging, and movement (miles)</i>	2,784	7	6	0	6	5	9	0	0	0	0	0	0	0	0	61	0	15
Western pond turtle																		
<i>Aquatic habitat¹⁰</i>	81,588	180	57	0	2,098	37	23	0	0	0	0	0	0	0	0	320	0	2,120
<i>Upland nesting and overwintering habitat</i>	16,043	105	97	0	34	109	70	0	0	4	0	0	0	0	0	451	0	104
<i>Upland nesting and overwintering habitat-NHD²⁰</i>	12,615	30	47	0	34	21	49	0	0	0	0	0	0	0	0	242	0	83
Total	110,245	315	201	0	2,166	167	141	0	0	4	0	0	0	0	0	1,012	0	2,307
<i>Aquatic habitat linear (miles) – NHD²⁰</i>	1,418	3	6	0	3	1	3	0	0	0	0	0	0	0	0	34	0	6
Amphibians																		
California red-legged frog																		
<i>Aquatic habitat</i>	159	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
<i>Upland cover and dispersal habitat</i>	7,766	6	0	0	39	0	0	0	0	0	0	0	0	8	0	14	0	39
Total	7,925	7	0	0	39	0	0	0	0	0	0	0	0	8	0	15	0	39
<i>Aquatic habitat (miles)</i>	30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
California tiger salamander																		
<i>Aquatic breeding habitat</i>	7,845	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Terrestrial cover and aestivation</i>	28,173	6	0	0	32	42	0	0	0	0	0	0	0	12	35	298	0	32
Total	36,018	6	0	0	32	42	0	0	0	0	0	0	0	12	35	298	0	32
Invertebrates																		
Valley elderberry longhorn beetle																		
<i>Riparian vegetation</i>	17,464	16	18	1	29	83	76	0	0	0	0	0	0	0	0	415	1	105
<i>Nonriparian channels and grasslands</i>	16,585	126	101	0	62	41	94	0	0	0	0	0	0	0	0	369	0	156
Total	34,048	142	119	1	90	125	170	0	0	0	0	0	0	0	0	784	1	261
California linderiella																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	35	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	43	0	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)
Conservancy fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	35	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	43	0	0
Longhorn fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	35	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	43	0	0
Midvalley fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	35	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	43	0	0
Vernal pool fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	35	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	43	0	0
Vernal pool tadpole shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	35	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	43	0	0

¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Non-federal Actions.

² Existing habitat and habitat loss are estimated using habitat models created from detailed vegetation mapping. See Appendix 2.A, *Covered Species Accounts*, for a complete description of species-specific mapping methods. Effects on species' habitat will be tracked during implementation through on-the-ground surveys performed by qualified biologists.

³ See Table 5.J.1, *Quantitative Effects Analysis Methods and Assumptions*, in Appendix 5.J, *Effects on Natural Communities, Wildlife, and Plants*, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, *Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat*, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.

⁴ Permanent and temporary effects assessed under CM1 are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Section 4.2.1.1, North Delta Diversions Construction and Operations, for a complete description of all activities assessed under CM1.

⁵ Borrow/Spoil Area Borrow: location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil.

⁶ Permanent and temporary effects assessed under CM2 include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.

⁷ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, *Quantitative Effects Analysis Methods*

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent (Acres) ⁴	Permanent - Reusable Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Permanent (Acres)
<p><i>and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>⁸ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>⁹ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>¹⁰ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>¹¹ Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>¹² Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>¹³ Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁴ AMM30 (Appendix 3.C) requires a reroute of the transmission line so it does not affect a roost site. This will reduce impacts on roosting and foraging habitat by 29 acres.</p> <p>¹⁵ Although the tidal restoration model results in some decreases in acreage of natural community loss between near term and late long-term due to tidal damping and sea level rise, for permitting purposes the maximum acreage of loss is shown for late long-term.</p> <p>¹⁶ Because decimal places are not shown in this table, in some cases, a row total may be larger by one or two acres than the result obtained by manually summing numbers across columns.</p> <p>¹⁷ Reusable tunnel material is flexible and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.</p> <p>¹⁸ Loss reduced to zero. Although the temporary transmission powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools complex be avoided during transmission powerline installation.</p> <p>¹⁹ Rice loss from CM8 and CM10 are not included in this analysis as rice conversion in Conservation Zone 2 will be avoided. This table will be updated for all other species in the next version.</p> <p>²⁰ For western pond turtle NHD model types, a 35% habitat suitability correction factor was applied to existing modeled habitat and covered activity loss acreage as it was determined that, in the Plan Area, approximately 35% of all channels and ditches mapped in the NHD layer are likely suitable for western pond turtle. See Appendix 2.A, <i>Covered Species Accounts</i>, Section 2.A.29, for more details.</p> <p>NHD = National Hydrologic Database; SWP = State Water Project; CVP = Central Valley Project.</p>																		

1
2

1 Table 5.J-10. Early Long-Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh						Delta						Plan Area Total ⁷		
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
Mammals																
Riparian brush rabbit																
<i>Riparian habitat</i>	2,909	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Grassland habitat</i>	3,103	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	6,011	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Riparian woodrat																
<i>Habitat</i>	2,166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	2,166	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Salt marsh harvest mouse																
<i>Tidal brackish emergent wetland primary</i>	3,641	0	0	66	0	0	0	0	0	0	0	0	0	0	0	66
<i>Tidal brackish emergent wetland secondary</i>	2,718	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland secondary</i>	749	1	7	0	0	0	0	0	0	0	0	0	0	0	0 ¹⁵	8
<i>Managed wetland—wetland primary, low long-term conservation value</i>	21,891	15	36	670	825	1,067	6	0	0	0	0	0	0	0	1,898	721
<i>Managed wetland—wetland secondary, low long-term conservation value</i>	2,800	5	16	226	87	52	1	0	0	0	0	0	0	0	140	248
<i>Managed wetland—upland, low long-term conservation value</i>	3,787	5	9	144	174	61	1	0	0	0	0	0	0	0	236	158
Total	35,588	26	69	1,106	1,086	1,181	8	0	0	0	0	0	0	0	2,275	1,201
San Joaquin kit fox																
<i>Breeding, foraging, and dispersal habitat</i>	5,327	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	5,327	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Suisun shrew																
<i>Primary habitat</i>	3,128	0	0	59	0	0	0	0	0	0	0	0	0	0	59	0
<i>Secondary habitat</i>	4,387	7	17	0	58	22	1	0	0	0	0	0	0	0	81	24
Total	7,515	7	17	59	58	22	1	0	0	0	0	0	0	0	140	24
Birds																
California black rail																
<i>Primary habitat</i>	7,467	0	0	71	0	0	0	0	0	0	0	0	1	0	2	71
<i>Secondary habitat</i>	17,915	29	63	0	607	757	4	0	0	0	1	0	0	0	1,367 ¹⁵	93
Total	25,382	29	63	71	607	757	4	0	0	0	1	0	1	0	1,369	164
California clapper rail¹³																
<i>Primary habitat</i>	296	0	0	26	0	0	0	0	0	0	0	0	0	0	0	26
<i>Secondary habitat</i>	6,420	3	6	0	0	0	0	0	0	0	0	0	0	0	0 ¹⁵	8
Total	6,716	3	6	26	0	0	0	0	0	0	0	0	0	0	0	35
California least tern																
<i>Nesting and Migratory Habitat</i>	86,263	0	0	1	0	0	0	0	1	1	13	13	4	0	33	0
Total	86,263	0	0	1	0	0	0	0	1	1	13	13	4	0	33	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh							Delta						Plan Area Total ⁷	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
Greater sandhill crane																
<i>Roosting and foraging - Permanent</i>	7,340	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Roosting and foraging - Temporary</i>	16,522	0	0	0	0	0	0	0	0	0	41	0	0	0	41	0
<i>Foraging</i>	162,164	0	0	0	0	0	0	0	11	0	1,407	827	381	29	2,655	0
Total	186,025	0	0	0	0	0	0	0	11	0	1,448	827	381	29	2,696	0
Least Bell's vireo																
<i>Migratory and breeding</i>	14,528	1	1	6	1	1	0	0	0	0	381	6	4	2	403	0
Total	14,528	1	1	6	1	1	0	0	0	0	381	6	4	2	403	0
Suisun song sparrow																
<i>Primary habitat</i>	3,722	0	0	55	0	0	0	0	0	0	0	0	0	0	0	55
<i>Secondary habitat</i>	23,986	32	70	0	623	851	5	0	0	0	0	0	0	0	1,479 ¹⁵	102
Total	27,707	32	70	55	624	851	5	0	0	0	0	0	0	0	1,479	157
Swainson's hawk																
<i>Foraging habitat</i>	470,324	24	45	324	216	97	0	0	1,073	192	7,437	2,007	604	37	12,057	0
<i>Nesting habitat</i>	9,796	0	0	4	2	0	0	0	0	0	217	2	2	1	228	0
Total	480,120	24	46	328	218	97	0	0	1,073	193	7,654	2,009	605	38	12,285	0
Tricolored blackbird																
<i>Breeding habitat-ag foraging</i>	100,198	0	0	0	0	0	0	0	554	143	2,341	421	7	1	1,126	2,341
<i>Breeding habitat-foraging</i>	58,181	11	20	294	281	165	1	0	7	0	98	2	0	0	468	413
<i>Breeding habitat-nesting</i>	1,741	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1
<i>Nonbreeding habitat-foraging ag</i>	194,251	0	0	0	0	0	0	0	223	27	2,870	986	548	29	1,813	2,870
<i>Nonbreeding habitat-roosting</i>	28,066	23	43	0	315	282	3	0	91	0	0	4	3	1	765	0
<i>Nonbreeding habitat-foraging</i>	34,308	0	0	0	0	0	0	0	42	3	547	37	17	6	105	547
Total	416,745	34	63	294	596	448	4	0	917	174	5,857	1,451	574	37	4,277	6,172
Western burrowing owl																
<i>High-value habitat</i>	149,783	15	24	212	219	105	1	0	886	4	4,039	146	157	34	5,840	0
<i>Low-value habitat</i>	251,767	0	15	25	6	2	0	0	192	157	2,994	1,323	298	13	5,025	0
Total	401,550	15	38	237	225	107	1	0	1,077	160	7,033	1,469	455	46	10,865	0
Western yellow-billed cuckoo																
<i>Breeding habitat</i>	1,970	0	0	0	0	0	0	0	0	0	52	0	1	0	53	0
<i>Migratory habitat</i>	10,425	0	0	0	0	0	0	0	0	0	248	2	1	1	254	0
Total	12,395	0	0	0	0	0	0	0	0	0	300	3	2	2	307	0
White-tailed kite																
<i>Breeding habitat</i>	14,069	0	0	4	2	0	0	0	0	0	284	3	2	2	298	0
<i>Foraging habitat</i>	500,365	0	79	1,054	1,124	1,360	23	0	0	193	7,288	1,999	605	38	13,764	0
Total	514,434	0	79	1,059	1,127	1,360	23	0	0	193	7,572	2,001	607	40	14,061	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh							Delta						Plan Area Total ⁷	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
Yellow-breasted chat																
<i>Primary nesting and migratory habitat</i>	8,178	0	0	0	0	0	0	0	0	0	105	1	2	1	109	0
<i>Secondary nesting and migratory habitat</i>	5,528	0	0	0	0	0	0	0	0	0	276	4	2	1	283	0
<i>Suisun Marsh/Upper Yolo Bypass nest and migratory habitat</i>	841	1	1	6	1	1	0	0	0	0	0	0	0	0	11	0
Total	14,547	1	1	6	1	1	0	0	0	0	381	6	4	2	403	0
Reptiles																
Giant garter snake																
<i>Aquatic - tidal</i>	12,097	0	0	0	0	0	0	0	2	0	0	0	0	0	2	0
<i>Aquatic - nontidal¹⁹</i>	19,027	0	0	0	0	0	0	0	8	0	134	12	3	0	158	0
<i>Upland-high</i>	21,581	0	0	0	0	0	0	0	41	0	420	33	32	17	544	0
<i>Upland-moderate</i>	25,407	0	0	0	0	0	0	0	66	1	722	69	12	1	871	0
<i>Upland-low</i>	5,683	0	0	0	0	0	0	0	6	0	122	1	1	1	131	0
Total	83,796	0	0	0	0	0	0	0	122	2	1,398	115	48	20	1,705	0
<i>Aquatic breeding, foraging, and movement (miles)</i>	2,784	0	0	0	0	0	0	0	4	1	49	11	7	2	74	0
Western pond turtle																
<i>Aquatic habitat¹⁰</i>	81,588	45	0	0	0	0	0	0	0	0	0	0	0	0	45	0
<i>Upland nesting and overwintering habitat</i>	16,043	2	4	108	98	35	1	0	2	0	44	2	0	0	295	0
<i>Upland nesting and overwintering habitat-NHD²⁰</i>	12,615	4	7	11	15	7	0	0	7	1	164	10	6	2	235	0
Total	110,245	52	11	120	113	41	1	0	9	1	208	12	6	2	576	0
<i>Aquatic habitat linear (miles) - NHD²⁰</i>	1,418	0	0	1	1	1	0	0	2	1	26	8	4	1	46	0
Amphibians																
California red-legged frog																
<i>Aquatic habitat</i>	159	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland cover and dispersal habitat</i>	7,766	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	7,925	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Aquatic habitat (miles)</i>	30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
California tiger salamander																
<i>Aquatic breeding habitat</i>	7,845	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Terrestrial cover and aestivation</i>	28,173	0	0	0	0	0	0	0	140	0	135	0	0	0	275	0
Total	36,018	0	0	0	0	0	0	0	140	0	135	0	0	0	275	0
Invertebrates																
Valley elderberry longhorn beetle																
<i>Riparian vegetation</i>	17,464	0	0	0	0	0	0	0	0	0	389	7	5	2	403	0
<i>Nonriparian channels and grasslands</i>	16,585	1	1	14	9	2	0	0	10	0	107	15	6	1	164	0
Total	34,048	1	1	14	9	2	0	0	10	0	497	22	10	2	568	0
California linderiella																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh						Delta						Plan Area Total ⁷		
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
<i>Degraded vernal pool complex</i>	2,713	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Conservancy fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Longhorn fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Midvalley fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Vernal pool fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Vernal pool tadpole shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38	0

1
2

1 **Table 5.J-10. Early Long-Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity (cont'd)**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²			
Mammals																		
Riparian brush rabbit																		
<i>Riparian habitat</i>	2,909	3	0	0	1	0	0	6	6	0	0	0	0	0	0	9	0	7
<i>Grassland habitat</i>	3,103	124	0	0	54	0	0	3	6	0	0	0	0	0	127	0	60	
Total	6,011	127	0	0	54	0	0	8	12	0	0	0	0	0	136	0	67	
Riparian woodrat																		
<i>Habitat</i>	2,166	0	0	0	0	0	0	6	6	0	0	0	0	0	6	0	6	
Total	2,166	0	0	0	0	0	0	6	6	0	0	0	0	0	6	0	6	
Salt marsh harvest mouse																		
<i>Tidal brackish emergent wetland primary</i>	3,641	0	0	0	0	0	0	0	0	0	0	0	0	0	66	0	0	
<i>Tidal brackish emergent wetland secondary</i>	2,718	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Upland secondary</i>	749	0	0	0	0	0	0	0	0	0	0	0	0	0	8	0	0	
<i>Managed wetland—wetland primary, low long-term conservation value</i>	21,891	0	0	0	0	0	0	0	0	0	0	0	0	0	2,619	0	0	
<i>Managed wetland—wetland secondary, low long-term conservation value</i>	2,800	0	0	0	0	0	0	0	0	0	0	0	0	0	387	0	0	
<i>Managed wetland—upland, low long-term conservation value</i>	3,787	0	0	0	0	0	0	0	0	0	0	0	0	0	395	0	0	
Total	35,588	0	0	0	0	0	0	0	0	0	0	0	0	0	3,476	0	0	
San Joaquin kit fox																		
<i>Breeding, foraging, and dispersal habitat</i>	5,327	155	52	0	103	0	0	0	0	0	0	0	0	4	211	0	103	
Total	5,327	155	52	0	103	0	0	0	0	0	0	0	0	4	211	0	103	
Suisun shrew																		
<i>Primary habitat</i>	3,128	0	0	0	0	0	0	0	0	0	0	0	0	0	59	0	0	
<i>Secondary habitat</i>	4,387	0	0	0	0	0	0	0	0	0	0	0	0	0	105	0	0	

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶
Total	7,515	0	0	0	0	0	0	0	0	0	0	0	0	0	0	164	0	0
Birds																		
California black rail																		
Primary habitat	7,467	0	0	0	18	5	0	0	0	0	0	0	0	0	0	77	0	18
Secondary habitat	17,915	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1,460 ¹⁵	0	0
Total	25,382	0	0	0	18	5	0	0	0	0	0	0	0	0	0	1,538	0	18
California clapper rail¹³																		
Primary habitat	296	0	0	0	0	0	0	0	0	0	0	0	0	0	0	26	0	0
Secondary habitat	6,420	0	0	0	0	0	0	0	0	0	0	0	0	0	0	9 ¹⁵	0	0
Total	6,716	0	0	0	0	0	0	0	0	0	0	0	0	0	0	35	0	0
California least tern																		
Nesting and Migratory Habitat	86,263	178	0	0	2,101	8	11	0	0	0	0	0	0	0	0	219	0	2,112
Total	86,263	178	0	0	2,101	8	11	0	0	0	0	0	0	0	0	219	0	2,112
Greater sandhill crane																		
Roosting and foraging - Permanent	7,340	0	0	0	8	0	0	0	0	0	0	0	0	0	0	0	0	8
Roosting and foraging - Temporary	16,522	0 ¹⁴	0	0	16	0	0	0	0	0	0	0	0	0	0	41	0	16
Foraging	162,164	352	2,347	183	778	0	0	0	0	0	333	750	3	0	6,439	183	778	
Total	186,025	352	2,347	183	802	0	0	0	0	0	333	750	3	0	6,481	183	802	
Least Bell's vireo																		
Migratory and breeding	14,528	11	18	1	22	83	88	6	4	0	0	0	0	0	0	521	1	114
Total	14,528	11	18	1	22	83	88	6	4	0	0	0	0	0	0	521	1	114
Suisun song sparrow																		
Primary habitat	3,722	0	0	0	0	0	0	0	0	0	0	0	0	0	55	0	0	
Secondary habitat	23,986	0	0	0	0	0	0	0	0	0	0	0	0	0	1,581 ¹⁵	0	0	
Total	27,707	0	0	0	0	0	0	0	0	0	0	0	0	0	1,637	0	0	
Swainson's hawk																		
Foraging habitat	470,324	1,100	3,235	183	1,113	996	504	197	122	726	0	1,368	746	20	35	20,479	183	1,739
Nesting habitat	9,796	8	10	0	18	79	54	6	6	0	0	0	0	0	330	0	78	
Total	480,120	1,108	3,245	183	1,131	1,075	558	202	128	726	0	1,368	746	20	35	20,809	183	1,817

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)	
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)				Permanent (Acres) ¹²
Tricolored blackbird																			
<i>Breeding habitat-ag foraging</i>	100,198	634	795	81	148	477	84	98	57	705	0	1,125	237	0	0	7,537	81	289	
<i>Breeding habitat-foraging</i>	58,181	161	52	0	114	105	155	11	11	7	0	0	0	20	35	1,272	0	279	
<i>Breeding habitat-nesting</i>	1,741	4	0	1	2	13	75	0	0	0	0	0	0	0	0	18	1	77	
<i>Nonbreeding habitat-foraging ag</i>	194,251	203	2,124	0	575	0	54	56	32	14	0	155	525	0	0	7,760	0	660	
<i>Nonbreeding habitat-roosting</i>	28,066	7	12	0	20	8	0	0	0	0	0	0	0	0	0	793	0	20	
<i>Nonbreeding habitat-foraging</i>	34,308	48	197	0	47	0	0	0	0	0	0	0	0	0	0	897	0	47	
Total	416,745	1,057	3,180	82	905	603	367	165	100	726	0	1,281	761	20	35	18,277	82	1,373	
Western burrowing owl																			
<i>High-value habitat</i>	149,783	340	541	0	351	882	245	13	13	7	0	268	86	20	35	8,030	0	609	
<i>Low-value habitat</i>	251,767	689	2,324	101	588	98	144	146	83	719	0	972	509	0	0	10,482	101	815	
Total	401,550	1,030	2,864	102	939	979	389	159	96	726	0	1,240	594	20	35	18,512	102	1,424	
Western yellow-billed cuckoo																			
<i>Breeding habitat</i>	1,970	3	6	0	1	26	5	0	0	0	0	0	0	0	0	89	0	5	
<i>Migratory habitat</i>	10,425	4	10	0	18	57	83	0	0	0	0	0	0	0	0	325	0	102	
Total	12,395	7	16	0	19	83	88	0	0	0	0	0	0	0	0	413	0	107	
White-tailed kite																			
<i>Breeding habitat</i>	14,069	10	16	0	23	82	88	6	6	0	0	0	0	0	0	411	0	116	
<i>Foraging habitat</i>	500,365	1,100	3,239	183	1,112	1,008	516	197	123	726	0	0	0	20	35	20,089	183	1,752	
Total	514,434	1,111	3,255	183	1,135	1,090	604	203	130	726	0	0	0	20	35	20,501	183	1,869	
Yellow-breasted chat																			
<i>Primary nesting and migratory habitat</i>	8,178	7	10	0	6	9	58	6	4	0	0	0	0	0	0	141	0	68	
<i>Secondary nesting and migratory habitat</i>	5,528	3	8	1	16	3	0	0	0	0	0	0	0	0	0	297	1	16	
<i>Suisun Marsh/Upper Yolo Bypass nest and migratory habitat</i>	841	0	0	0	0	71	29	0	0	0	0	0	0	0	0	82	0	29	
Total	14,547	10	18	1	22	83	88	6	4	0	0	0	0	0	0	520	1	114	
Reptiles																			

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)	
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)				Permanent (Acres) ¹²
Giant garter snake																			
<i>Aquatic - tidal</i>	12,097	16	1	0	55	9	2	0	0	0	0	0	0	0	0	27	0	58	
<i>Aquatic - nontidal¹⁹</i>	19,027	10	56	0	13	59	13	20	10	0	0	0	0	0	0	303	0	36	
<i>Upland-high</i>	21,581	66	106	0	48	178	158	0	0	0	0	0	0	0	0	894	0	206	
<i>Upland-moderate</i>	25,407	167	54	0	135	60	61	6	8	0	0	0	0	0	35	1,193	0	204	
<i>Upland-low</i>	5,683	14	4	0	5	1	0	4	3	0	0	0	0	0	0	154	0	8	
Total	83,796	274	222	0	257	306	234	30	22	0	0	0	0	0	35	2,572	0	513	
<i>Aquatic breeding, foraging, and movement (miles)</i>	2,784	7	6	0	6	5	9	0	0	0	0	0	0	0	0	91	0	15	
Western pond turtle																			
<i>Aquatic habitat¹⁰</i>	81,588	180	57	0	2,098	37	23	19	9	0	0	0	0	0	0	339	0	2,129	
<i>Upland nesting and overwintering habitat</i>	16,043	105	97	0	34	109	70	3	8	6	0	0	0	0	0	615	0	112	
<i>Upland nesting and overwintering habitat-NHD²⁰</i>	12,615	30	47	0	34	21	49	0	0	0	0	0	0	0	0	333	0	83	
Total	110,245	315	201	0	2,166	167	141	22	17	6	0	0	0	0	0	1,287	0	2,324	
<i>Aquatic habitat linear (miles) – NHD²⁰</i>	1,418	3	6	0	3	1	3	0	0	0	0	0	0	0	0	56	0	6	
Amphibians																			
California red-legged frog																			
<i>Aquatic habitat</i>	159	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	
<i>Upland cover and dispersal habitat</i>	7,766	6	0	0	39	0	0	0	0	0	0	0	0	11	0	17	0	39	
Total	7,925	7	0	0	39	0	0	0	0	0	0	0	0	11	0	18	0	39	
<i>Aquatic habitat (miles)</i>	30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
California tiger salamander																			
<i>Aquatic breeding habitat</i>	7,845	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
<i>Terrestrial cover and aestivation</i>	28,173	6	0	0	32	42	0	0	0	0	0	0	0	17	35	375	0	32	
Total	36,018	6	0	0	32	42	0	0	0	0	0	0	0	17	35	375	0	32	

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			
Invertebrates																		
Valley elderberry longhorn beetle																		
<i>Riparian vegetation</i>	17,464	16	18	1	29	83	76	6	6	0	0	0	0	0	0	526	1	111
<i>Nonriparian channels and grasslands</i>	16,585	126	101	0	62	41	94	3	6	0	0	0	0	0	0	436	0	162
Total	34,048	142	119	1	90	125	170	9	13	0	0	0	0	0	0	962	1	273
California linderiella																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	45	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Conservancy fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	45	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Longhorn fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	45	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Midvalley fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	45	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Vernal pool fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	45	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Vernal pool tadpole shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	45	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)	
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²				
<p>¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Non-federal Actions.</p> <p>² Existing habitat and habitat loss are estimated using habitat models created from detailed vegetation mapping. See Appendix 2.A, <i>Covered Species Accounts</i>, for a complete description of species-specific mapping methods. Effects on species' habitat will be tracked during implementation through on-the-ground surveys performed by qualified biologists.</p> <p>³ See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, <i>Effects on Natural Communities, Wildlife, and Plants</i>, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, <i>Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat</i>, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.</p> <p>⁴ Permanent and temporary effects assessed under CM1 are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Section 4.2.1.1, North Delta Diversions Construction and Operations, for a complete description of all activities assessed under CM1.</p> <p>⁵ Borrow/Spoil Area Borrow: location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil.</p> <p>⁶ Permanent and temporary effects assessed under CM2 include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.</p> <p>⁷ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>⁸ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>⁹ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>¹⁰ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>¹¹ Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>¹² Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>¹³ Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁴ AMM30 (Appendix 3.C) requires a reroute of the transmission line so it does not affect a roost site. This will reduce impacts on roosting and foraging habitat by 29 acres.</p> <p>¹⁵ Although the tidal restoration model results in some decreases in acreage of natural community loss between near term and late long-term due to tidal damping and sea level rise, for permitting purposes the maximum acreage of loss is shown for late long-term.</p> <p>¹⁶ Because decimal places are not shown in this table, in some cases, a row total may be larger by one or two acres than the result obtained by manually summing numbers across columns.</p> <p>¹⁷ Reusable tunnel material is flexible and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.</p> <p>¹⁸ Loss reduced to zero. Although the temporary transmission powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools complex be avoided during transmission powerline installation.</p> <p>¹⁹ Rice loss from CM8 and CM10 are not included in this analysis as rice conversion in Conservation Zone 2 will be avoided. This table will be updated for all other species in the next version.</p> <p>²⁰ For western pond turtle NHD model types, a 35% habitat suitability correction factor was applied to existing modeled habitat and covered activity loss acreage as it was determined that, in the Plan Area, approximately 35% of all channels and ditches mapped in the NHD layer are likely suitable for western pond turtle. See Appendix 2.A, <i>Covered Species Accounts</i>, Section 2.A.29, for more details.</p> <p>NHD = National Hydrologic Database; SWP = State Water Project; CVP = Central Valley Project.</p>																			

1
2

1 **Table 5.J-11. Late Long-Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh							Delta						Plan Area Total ⁷	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
Mammals																
Riparian brush rabbit																
<i>Riparian habitat</i>	2,909	0	0	0	0	0	0	0	5	0	7	3	3	1	19	0
<i>Grassland habitat</i>	3,103	0	0	0	0	0	0	0	3	0	11	1	2	0	18	0
Total	6,011	0	0	0	0	0	0	0	8	0	19	4	4	1	37	0
Riparian woodrat																
<i>Habitat</i>	2,166	0	0	0	0	0	0	0	4	0	5	1	0	0	10	0
Total	2,166	0	0	0	0	0	0	0	4	0	5	1	0	0	10	0
Salt marsh harvest mouse																
<i>Tidal brackish emergent wetland primary</i>	3,641	0	0	67	0	0	0	0	0	0	0	0	0	0	0	67
<i>Tidal brackish emergent wetland secondary</i>	2,718	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland secondary</i>	749	1	7	0	0	0	0	0	0	0	0	0	0	0	1 ¹⁵	8
<i>Managed wetland—wetland primary, low long-term conservation value</i>	21,891	5	7	991	807	3,353	160	0	0	0	0	0	0	0	4,320	1,003
<i>Managed wetland—wetland secondary, low long-term conservation value</i>	2,800	2	3	336	135	317	14	0	0	0	0	0	0	0	467	340
<i>Managed wetland—upland, low long-term conservation value</i>	3,787	6	9	158	164	419	5	0	0	0	0	0	0	0	588	174
Total	35,588	13	26	1,552	1,107	4,090	179	0	0	0	0	0	0	0	5,376	1,592
San Joaquin kit fox																
<i>Breeding, foraging, and dispersal habitat</i>	5,327	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	5,327	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Suisun shrew																
<i>Primary habitat</i>	3,128	0	0	60	0	0	0	0	0	0	0	0	0	0	60	0
<i>Secondary habitat</i>	4,387	7	17	0	97	208	12	0	0	0	0	0	0	0	318 ¹⁵	24
Total	7,515	7	17	60	97	208	12	0	0	0	0	0	0	0	377	24
Birds																
California black rail																
<i>Primary habitat</i>	7,467	0	0	71	0	0	0	0	5	0	0	1	1	1	3	76
<i>Secondary habitat</i>	17,915	29	52	0	587	2,240	118	0	0	0	12	5	0	0	2,951 ¹⁵	93
Total	25,382	29	52	71	587	2,240	118	0	5	0	12	6	2	1	2,954	168
California clapper rail¹³																
<i>Primary habitat</i>	296	0	0	27	0	0	0	0	0	0	0	0	0	0	0	27
<i>Secondary habitat</i>	6,420	5	3	0	0	0	0	0	0	0	0	0	0	0	0 ¹⁵	8

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh							Delta						Plan Area Total ⁷	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
Total	6,716	5	3	27	0	0	0	0	0	0	0	0	0	0	0	35
California least tern																
<i>Nesting and Migratory Habitat</i>	86,263	0	0	1	0	0	0	0	1	1	14	8	9	0	36	0
Total	86,263	0	0	1	0	0	0	0	1	1	14	8	9	0	36	0
Greater sandhill crane																
<i>Roosting and foraging - Permanent</i>	7,340	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Roosting and foraging - Temporary</i>	16,522	0	0	0	0	0	0	0	0	0	41	0	0	0	41	0
<i>Foraging</i>	162,164	0	0	0	0	0	0	0	1	0	1,467	514	614	117	2,713	0
Total	186,025	0	0	0	0	0	0	0	1	0	1,508	514	614	117	2,754	0
Least Bell's vireo																
<i>Migratory and breeding</i>	14,528	2	1	8	1	2	0	0	0	0	477	23	25	6	545	0
Total	14,528	2	1	8	1	2	0	0	0	0	477	23	25	6	545	0
Suisun song sparrow																
<i>Primary habitat</i>	3,722	0	0	55	0	0	0	0	0	0	0	0	0	0	0	55
<i>Secondary habitat</i>	23,986	53	70	0	657	2,712	140	0	0	0	0	0	0	0	3,510 ¹⁵	123
Total	27,707	53	70	55	657	2,712	140	0	0	0	0	0	0	0	3,510	178
Swainson's hawk																
<i>Foraging habitat</i>	470,324	62	63	411	349	666	11	0	1,319	3	17,988	6,280	7,393	2,814	37,359	0
<i>Nesting habitat</i>	9,796	0	0	2	4	2	0	0	0	0	258	12	15	2	295	0
Total	480,120	62	63	413	353	668	11	0	1,319	3	18,246	6,292	7,408	2,816	37,654	0
Tricolored blackbird																
<i>Breeding habitat-ag foraging</i>	100,198	2	0	0	0	0	0	0	338	0	3,635	1,335	1,093	47	2,814	3,635
<i>Breeding habitat-foraging</i>	58,181	10	11	382	299	692	18	0	38	0	254	28	16	2	1,102	647
<i>Breeding habitat-nesting</i>	1,741	0	0	0	0	0	0	0	6	0	34	10	4	0	21	34
<i>Nonbreeding habitat-foraging ag</i>	194,251	0	0	0	0	0	0	0	530	1	8,716	2,991	4,115	851	8,489	8,716
<i>Nonbreeding habitat-roosting</i>	28,066	5	7	0	404	1,119	29	0	41	0	0	13	10	5	1,633	0
<i>Nonbreeding habitat-foraging</i>	34,308	0	0	8	44	465	7	0	38	3	651	33	49	33	672	659
Total	416,745	17	19	391	746	2,276	54	0	992	3	13,291	4,410	5,287	939	14,732	13,692
Western burrowing owl																
<i>High-value habitat</i>	149,783	39	40	324	216	795	81	0	620	3	6,253	783	617	158	9,929	0
<i>Low-value habitat</i>	251,767	0	4	44	21	14	0	0	478	0	9,281	3,919	3,751	2,226	19,739	0
Total	401,550	39	44	368	236	809	81	0	1,098	3	15,534	4,702	4,368	2,384	29,668	0
Western yellow-billed cuckoo																
<i>Breeding habitat</i>	1,970	0	0	0	0	0	0	0	0	0	86	13	10	1	110	0
<i>Migratory habitat</i>	10,425	0	0	0	0	0	0	0	0	0	290	6	9	4	310	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh							Delta						Plan Area Total ⁷	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
Total	12,395	0	0	0	0	0	0	0	0	0	376	19	20	5	420	0
White-tailed kite																
<i>Breeding habitat</i>	14,069	0	0	2	4	2	0	0	0	0	339	15	17	3	383	0
<i>Foraging habitat</i>	500,365	0	71	1,372	1,133	4,528	425	0	0	3	17,811	6,227	7,240	2,815	41,625	0
Total	514,434	0	71	1,374	1,137	4,530	425	0	0	3	18,151	6,242	7,257	2,818	42,008	0
Yellow-breasted chat																
<i>Primary nesting and migratory habitat</i>	8,178	0	0	0	0	0	0	0	0	0	149	14	16	3	182	0
<i>Secondary nesting and migratory habitat</i>	5,528	0	0	0	0	0	0	0	0	0	328	9	9	3	349	0
<i>Suisun Marsh/Upper Yolo Bypass nest and migratory habitat</i>	841	2	1	8	1	2	0	0	0	0	0	0	0	0	14	0
Total	14,547	2	1	8	1	2	0	0	0	0	478	23	25	6	545	0
Reptiles																
Giant garter snake																
<i>Aquatic - tidal</i>	12,097	0	0	0	0	0	0	0	1	0	0	0	0	0	2	0
<i>Aquatic - nontidal¹⁹</i>	19,027	0	0	0	0	0	0	0	37	0	237	38	60	21	393	0
<i>Upland-high</i>	21,581	0	0	0	0	0	0	0	33	0	477	17	26	39	594	0
<i>Upland-moderate</i>	25,407	0	0	0	0	0	0	0	60	0	1,019	128	140	28	1,375	0
<i>Upland-low</i>	5,683	0	0	0	0	0	0	0	7	0	137	4	3	2	154	0
Total	83,796	0	0	0	0	0	0	0	139	1	1,870	188	230	90	2,518	0
<i>Aquatic breeding, foraging, and movement (miles)</i>	2,784	0	0	0	0	0	0	0	8	0	73	18	23	16	138	0
Western pond turtle																
<i>Aquatic habitat¹⁰</i>	81,588	45	0	0	0	0	0	0	0	0	0	0	0	0	45	0
<i>Upland nesting and overwintering habitat</i>	16,043	3	5	86	95	139	2	0	13	0	113	6	5	4	473	0
<i>Upland nesting and overwintering habitat-NHD²⁰</i>	12,615	9	9	48	11	64	1	0	12	1	203	16	15	11	399	0
Total	110,245	57	14	134	107	203	3	0	24	1	316	22	20	15	917	0
<i>Aquatic habitat linear (miles) - NHD²⁰</i>	1,418	0	0	2	2	7	1	0	4	0	43	15	20	12	106	0
Amphibians																
California red-legged frog																
<i>Aquatic habitat</i>	159	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland cover and dispersal habitat</i>	7,766	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	7,925	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Aquatic habitat (miles)</i>	30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
California tiger salamander																
<i>Aquatic breeding habitat</i>	7,845	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Terrestrial cover and aestivation</i>	28,173	0	0	0	0	0	0	0	101	0	404	5	6	0	517	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
CM4 Tidal Natural Communities Restoration																
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Suisun Marsh							Delta						Plan Area Total ⁷	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Permanent (Acres) ^{7,8,9}	Conversion (Acres)
Total	36,018	0	0	0	0	0	0	0	101	0	404	5	6	0	517	0
Invertebrates																
Valley elderberry longhorn beetle																
<i>Riparian vegetation</i>	17,464	0	0	0	0	0	0	0	0	0	490	25	28	8	552	0
<i>Nonriparian channels and grasslands</i>	16,585	1	2	11	10	18	0	0	18	0	149	16	22	13	260	0
Total	34,048	1	2	11	10	18	0	0	18	0	640	42	50	21	813	0
California linderiella																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Conservancy fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Longhorn fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Midvalley fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Vernal pool fairy shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Vernal pool tadpole shrimp																
<i>Vernal pool complex</i>	8,759	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Degraded vernal pool complex</i>	2,713	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52	0

1
2

1 Table 5.J-11. Late Long-Term Wildlife Modeled Habitat Loss and Conversion by Covered Activity (cont'd)

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			
Mammals																		
Riparian brush rabbit																		
<i>Riparian habitat</i>	2,909	3	0	0	1	0	0	43	35	0	0	0	0	0	0	65	0	35
<i>Grassland habitat</i>	3,103	124	0	0	54	0	0	26	20	0	0	0	0	0	0	168	0	74
Total	6,011	127	0	0	54	0	0	69	54	0	0	0	0	0	0	232	0	109
Riparian woodrat																		
<i>Habitat</i>	2,166	0	0	0	0	0	0	41	33	0	0	0	0	0	0	51	0	33
Total	2,166	0	0	0	0	0	0	41	33	0	0	0	0	0	0	51	0	33
Salt marsh harvest mouse																		
<i>Tidal brackish emergent wetland primary</i>	3,641	0	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
<i>Tidal brackish emergent wetland secondary</i>	2,718	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Upland secondary</i>	749	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Managed wetland—wetland primary, low long-term conservation value</i>	21,891	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5,323	0	0
<i>Managed wetland—wetland secondary, low long-term conservation value</i>	2,800	0	0	0	0	0	0	0	0	0	0	0	0	0	0	807	0	0
<i>Managed wetland—upland, low long-term conservation value</i>	3,787	0	0	0	0	0	0	0	0	0	0	0	0	0	0	762	0	0
Total	35,588	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6,968	0	0
San Joaquin kit fox																		
<i>Breeding, foraging, and dispersal habitat</i>	5,327	155	52	0	103	0	0	0	0	0	0	0	0	8	0	214	0	103
Total	5,327	155	52	0	103	0	0	0	0	0	0	0	0	8	0	214	0	103
Suisun shrew																		
<i>Primary habitat</i>	3,128	0	0	0	0	0	0	0	0	0	0	0	0	0	0	60	0	0

Maximum Allowable Habitat Loss by Covered Activity^{1,2,3}

Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶
<i>Secondary habitat</i>	4,387	0	0	0	0	0	0	0	0	0	0	0	0	0	0	342 ¹⁵	0	0
Total	7,515	0	0	0	0	0	0	0	0	0	0	0	0	0	0	401	0	0
Birds																		
California black rail																		
<i>Primary habitat</i>	7,467	0	0	0	18	5	0	0	0	0	0	0	0	0	0	83	0	18
<i>Secondary habitat</i>	17,915	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3,043 ¹⁵	0	0
Total	25,382	0	0	0	18	5	0	0	0	0	0	0	0	0	0	3,127	0	18
California clapper rail¹³																		
<i>Primary habitat</i>	296	0	0	0	0	0	0	0	0	0	0	0	0	0	0	27	0	0
<i>Secondary habitat</i>	6,420	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8 ¹⁵	0	0
Total	6,716	0	0	0	0	0	0	0	0	0	0	0	0	0	0	35	0	0
California least tern																		
<i>Nesting and Migratory Habitat</i>	86,263	178	0	0	2,101	8	11	2	5	0	0	0	0	0	0	224	0	2,116
Total	86,263	178	0	0	2,101	8	11	2	5	0	0	0	0	0	0	224	0	2,116
Greater sandhill crane																		
<i>Roosting and foraging - Permanent</i>	7,340	0	0	0	8	0	0	0	0	0	0	0	0	0	0	0	0	8
<i>Roosting and foraging - Temporary</i>	16,522	0 ¹⁴	0	0	16	0	0	0	0	0	0	0	0	0	0	41	0	16
<i>Foraging</i>	162,164	352	2,347	183	778	0	0	0	0	0	0	300	1,350	4	0	7,065	183	778
Total	186,025	352	2,347	183	802	0	0	0	0	0	0	300	1,350	4	0	7,107	183	802
Least Bell's vireo																		
<i>Migratory and breeding</i>	14,528	11	18	1	22	83	88	28	21	0	0	0	0	0	0	685	1	131
Total	14,528	11	18	1	22	83	88	28	21	0	0	0	0	0	0	685	1	131
Suisun song sparrow																		
<i>Primary habitat</i>	3,722	0	0	0	0	0	0	0	0	0	0	0	0	0	0	55	0	0
<i>Secondary habitat</i>	23,986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3633 ¹⁵	0	0
Total	27,707	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3,688	0	0
Swainson's hawk																		
<i>Foraging habitat</i>	470,324	1,100	3,235	183	1,113	996	504	1,820	1,036	971	3,991	1,849	1,440	50	35	52,845	183	2,653
<i>Nesting habitat</i>	9,796	8	10	0	18	79	54	38	31	0	0	0	0	0	0	430	0	104

Maximum Allowable Habitat Loss by Covered Activity^{1,2,3}

Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶
Total	480,120	1,108	3,245	183	1,131	1,075	558	1,857	1,067	971	3,991	1,849	1,440	50	35	53,275	183	2,756
Tricolored blackbird																		
<i>Breeding habitat-ag foraging</i>	100,198	634	795	81	148	477	84	503	275	7	0	1,521	568	0	0	10,954	81	507
<i>Breeding habitat-foraging</i>	58,181	161	52	0	114	105	155	47	30	11	0	0	0	44	35	2,204	0	298
<i>Breeding habitat-nesting</i>	1,741	4	0	1	2	13	75	4	2	0	0	0	0	0	0	77	1	79
<i>Nonbreeding habitat-foraging ag</i>	194,251	203	2,124	0	575	0	54	652	367	953	3,991	210	945	0	0	26,282	0	995
<i>Nonbreeding habitat-roosting</i>	28,066	7	12	0	20	8	0	1	1	0	0	0	0	0	0	1,662	0	22
<i>Nonbreeding habitat-foraging</i>	34,308	48	197	0	47	0	0	3	3	0	0	0	0	7	0	1,586	0	50
Total	416,745	1,057	3,180	82	905	603	367	1,211	678	971	3,991	1,731	1,513	50	35	42,766	82	1,950
Western burrowing owl																		
<i>High-value habitat</i>	149,783	340	541	0	351	882	245	142	83	11	0	362	159	50	35	12,450	0	679
<i>Low-value habitat</i>	251,767	689	2,324	101	588	98	144	1,452	827	960	3,991	1,314	952	0	0	31,519	101	1,558
Total	401,550	1,030	2,864	102	939	979	389	1,594	910	971	3,991	1,675	1,111	50	35	43,969	102	2,237
Western yellow-billed cuckoo																		
<i>Breeding habitat</i>	1,970	3	6	0	1	26	5	6	5	0	0	0	0	0	0	150	0	11
<i>Migratory habitat</i>	10,425	4	10	0	18	57	83	16	11	0	0	0	0	0	0	397	0	112
Total	12,395	7	16	0	19	83	88	21	17	0	0	0	0	0	0	547	0	123
White-tailed kite																		
<i>Breeding habitat</i>	14,069	10	16	0	23	82	88	42	33	0	0	0	0	0	0	533	0	144
<i>Foraging habitat</i>	500,365	1,100	3,239	183	1,112	1,008	516	1,706	968	971	3,991	1,849	1,440	50	35	57,015	183	2,597
Total	514,434	1,111	3,255	183	1,135	1,090	604	1,748	1,001	971	3,991	1,849	1,440	50	35	57,548	183	2,740
Yellow-breasted chat																		
<i>Primary nesting and migratory habitat</i>	8,178	7	10	0	6	9	58	23	15	0	0	0	0	0	0	232	0	79
<i>Secondary nesting and migratory habitat</i>	5,528	3	8	1	16	3	0	5	6	0	0	0	0	0	0	367	1	22
<i>Suisun Marsh/Upper Yolo Bypass nest and migratory habitat</i>	841	0	0	0	0	71	29	0	0	0	0	0	0	0	0	85	0	29
Total	14,547	10	18	1	22	83	88	28	21	0	0	0	0	0	0	684	1	131

Maximum Allowable Habitat Loss by Covered Activity^{1,2,3}

Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			
Reptiles																		
Giant garter snake																		
<i>Aquatic - tidal</i>	12,097	16	1	0	55	9	2	2	3	0	0	0	0	0	0	28	0	60
<i>Aquatic - nontidal¹⁹</i>	19,027	10	56	0	13	59	13	34	21	0	0	0	0	0	0	553	0	47
<i>Upland-high</i>	21,581	66	106	0	48	178	158	0	0	0	0	0	0	0	0	944	0	206
<i>Upland-moderate</i>	25,407	167	54	0	135	60	61	27	24	0	0	0	0	0	35	1,718	0	220
<i>Upland-low</i>	5,683	14	4	0	5	1	0	20	18	0	0	0	0	0	0	193	0	23
Total	83,796	274	222	0	257	306	234	82	65	0	0	0	0	0	35	3,437	0	556
<i>Aquatic breeding, foraging, and movement (miles)</i>	2,784	7	6	0	6	5	9	2	1	0	0	0	0	0	0	156	0	16
Western pond turtle																		
<i>Aquatic habitat¹⁰</i>	81,588	180	57	0	2,098	37	23	32	21	0	0	0	0	0	0	351	0	2,141
<i>Upland nesting and overwintering habitat</i>	16,043	105	97	0	34	109	70	12	15	10	0	0	0	0	0	805	0	119
<i>Upland nesting and overwintering habitat-NHD²⁰</i>	12,615	30	47	0	34	21	49	4	2	0	0	0	0	0	0	501	0	85
Total	110,245	315	201	0	2,166	167	141	48	39	10	0	0	0	0	0	1,657	0	2,346
<i>Aquatic habitat linear (miles) - NHD²⁰</i>	1,418	3	6	0	3	1	3	2	1	0	0	0	0	0	0	118	0	7
Amphibians																		
California red-legged frog																		
<i>Aquatic habitat</i>	159	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
<i>Upland cover and dispersal habitat</i>	7,766	6	0	0	39	0	0	0	0	0	0	0	24	0	30	0	39	
Total	7,925	7	0	0	39	0	0	0	0	0	0	0	24	0	31	0	39	
<i>Aquatic habitat (miles)</i>	30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
California tiger salamander																		
<i>Aquatic breeding habitat</i>	7,845	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Terrestrial cover and aestivation</i>	28,173	6	0	0	32	42	0	0	0	0	0	0	40	35	639	0	32	
Total	36,018	6	0	0	32	42	0	0	0	0	0	0	40	35	639	0	32	
Invertebrates																		

Maximum Allowable Habitat Loss by Covered Activity^{1,2,3}

Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶
Valley elderberry longhorn beetle																		
<i>Riparian vegetation</i>	17,464	16	18	1	29	83	76	43	35	0	0	0	0	0	0	712	1	140
<i>Nonriparian channels and grasslands</i>	16,585	126	101	0	62	41	94	9	14	0	0	0	0	0	0	538	0	170
Total	34,048	142	119	1	90	125	170	52	49	0	0	0	0	0	0	1,250	1	310
California linderiella																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	59	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
Conservancy fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	59	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
Longhorn fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	59	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
Midvalley fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	59	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
Vernal pool fairy shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	59	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
Vernal pool tadpole shrimp																		
<i>Vernal pool complex</i>	8,759	8	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	8	0	0
<i>Degraded vernal pool complex</i>	2,713	7	0	0	0 ¹⁸	0	0	0	0	0	0	0	0	0	0	59	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0

Maximum Allowable Habitat Loss by Covered Activity^{1,2,3}

Covered Wildlife Species	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)			

¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Non-federal Actions.

² Existing habitat and habitat loss are estimated using habitat models created from detailed vegetation mapping. See Appendix 2.A, *Covered Species Accounts*, for a complete description of species-specific mapping methods. Effects on species' habitat will be tracked during implementation through on-the-ground surveys performed by qualified biologists.

³ See Table 5.J.1, *Quantitative Effects Analysis Methods and Assumptions*, in Appendix 5.J, *Effects on Natural Communities, Wildlife, and Plants*, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, *Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat*, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.

⁴ Permanent and temporary effects assessed under CM1 are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Section 4.2.1.1, North Delta Diversions Construction and Operations, for a complete description of all activities assessed under CM1.

⁵ Borrow/Spoil Area Borrow: location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil.

⁶ Permanent and temporary effects assessed under CM2 include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.

⁷ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, *Quantitative Effects Analysis Methods and Assumptions*, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.

⁸ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, *BDCP Tidal Habitat Evolution Assessment*) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.

⁹ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.

¹⁰ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.

¹¹ Based on restoration design assumptions described in Appendix 5.E, *Habitat Restoration*, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.

¹² Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.

¹³ Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.

¹⁴ AMM30 (Appendix 3.C) will require a reroute of the transmission line so it does not affect a roost site. This will reduce impacts on roosting and foraging habitat by 29 acres.

¹⁵ Although the tidal restoration model results in some decreases in acreage of natural community loss between near term and late long-term due to tidal damping and sea level rise, for permitting purposes the maximum acreage of loss is shown for late long-term.

¹⁶ Because decimal places are not shown in this table, in some cases, a row total may be larger by one or two acres than the result obtained by manually summing numbers across columns.

¹⁷ Reusable tunnel material is flexible and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.

¹⁸ Loss reduced to zero. Although the temporary transmission powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools complex be avoided during transmission powerline installation.

¹⁹ Rice loss from CM8 and CM10 are not included in this analysis as rice conversion in Conservation Zone 2 will be avoided. This table will be updated for all other species in the next version.

²⁰ For western pond turtle NHD model types, a 35% habitat suitability correction factor was applied to existing modeled habitat and covered activity loss acreage as it was determined that, in the Plan Area, approximately 35% of all channels and ditches mapped in the NHD layer are likely suitable for western pond turtle. See Appendix 2.A, *Covered Species Accounts*, Section 2.A.29, for more details.

NHD = National Hydrologic Database; SWP = State Water Project; CVP = Central Valley Project.

1
2

1 **Table 5.J-12. Near Term Plant Modeled Habitat Loss and Conversion by Covered Activity**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																	
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM4 Tidal Natural Communities Restoration													Plan Area Total ⁷		
		Suisun Marsh						Delta						Permanent (Acres) ^{7,8,9}	Conversion (Acres)		
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2			Subtidal 3	
Plants																	
Brittlescale total ¹³	451	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heartscale total	6,451	0	0	1	0	0	0	0	0	0	0	0	0	0	0	2	0
San Joaquin spearscale total	14,477	0	0	1	0	0	0	0	164	0	31	1	0	0	196	0	0
Carquinez goldenbush total ¹³	1,346	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Delta button celery total	3,361	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Delta mudwort total	6,081	0	0	0	0	0	0	0	0	0	0	3	1	2	5	0	0
Mason's lilaepsis total	6,081	0	0	0	0	0	0	0	0	0	0	3	1	2	5	0	0
Delta tule pea total ¹⁴	5,853	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Suisun Marsh aster total ¹⁴	5,853	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Side-flowering skullcap total ¹⁵	2,497	0	0	0	0	0	0	0	1	0	0	1	1	1	3	0	0
Slough thistle total	1,834	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Soft bird's-beak total	1,228	0	0	72	0	0	0	0	0	0	0	0	0	0	72	0	0
Suisun thistle total	1,281	0	0	72	0	0	0	0	0	0	0	0	0	0	72	0	0
Vernal Pool Plants																	
Alkali milk-vetch																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	24	0	3	0	0	0	27	0	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28²⁰	0	0
Legenere																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	24	0	3	0	0	0	27	0	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28²⁰	0	0
Heckard's peppergrass																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	24	0	3	0	0	0	27	0	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28²⁰	0	0
Boggs Lake hedge-hyssop																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	24	0	3	0	0	0	27	0	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28²⁰	0	0
Dwarf downingia																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	24	0	3	0	0	0	27	0	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	1	0	0	0	0	24	0	3	0	0	0	28²⁰	0	0

2

1 **Table 5.J-12. Near Term Plant Modeled Habitat Loss and Conversion by Covered Activity (cont'd)**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction				
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
Plants																			
Brittlescale total ¹³	451	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Heartscale total	6,451	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	
San Joaquin spearscale total	14,477	23	30	0	29	56	0	0	0	0	0	0	0	0	0	304	0	30	
Carquinez goldenbush total ¹³	1,346	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Delta button celery total	3,361	34	39	0	23	0	0	0	0	0	0	0	0	0	0	72	0	23	
Delta mudwort total	6,081	12	3	0	15	3	2	0	0	0	0	0	0	0	0	23	0	17	
Mason's lilaepsis total	6,081	12	3	0	15	3	2	0	0	0	0	0	0	0	0	23	0	17	
Delta tule pea total ¹⁴	5,853	2	0	0	1	0	0	0	0	0	0	0	0	0	0	3	0	1	
Suisun Marsh aster total ¹⁴	5,853	2	0	0	1	0	0	0	0	0	0	0	0	0	0	3	0	1	
Side-flowering skullcap total ¹⁵	2,497	3	0	0	5	0	0	0	0	0	0	0	0	0	0	7	0	5	
Slough thistle total	1,834	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soft bird's-beak total	1,228	0	0	0	0	0	0	0	0	0	0	0	0	0	0	72	0	0	
Suisun thistle total	1,281	0	0	0	0	0	0	0	0	0	0	0	0	0	0	72	0	0	
Vernal Pool Plants																			
Alkali milk-vetch																0	0	0	
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	34	0	0	
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	15	0	0	0¹⁹	0	0	0	0	0	0	0	0	0	0	43	0	0	
Legenere																			
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	34	0	0	

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction				
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	15	0	0	0 ¹⁹	0	0	0	0	0	0	0	0	0	0	43	0	0	
Heckard's peppergrass																0	0	0	
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	34	0	0	
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	15	0	0	0 ¹⁹	0	0	0	0	0	0	0	0	0	0	43	0	0	
Boggs Lake hedge-hyssop																0	0	0	
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	34	0	0	
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	15	0	0	0 ¹⁹	0	0	0	0	0	0	0	0	0	0	43	0	0	
Dwarf downingia																0	0	0	
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	34	0	0	
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	15	0	0	0 ¹⁹	0	0	0	0	0	0	0	0	0	0	43	0	0	

¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Non-federal Actions.

² Existing habitat and habitat loss are estimated using habitat models created from detailed vegetation mapping. See Appendix 2.A, *Covered Species Accounts*, for a complete description of species-specific mapping methods. Effects on species' habitat will be tracked during implementation through on-the-ground surveys performed by qualified biologists.

³ See Table 5.J.1, *Quantitative Effects Analysis Methods and Assumptions*, in Appendix 5.J, *Effects on Natural Communities, Wildlife, and Plants*, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, *Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat*, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.

⁴ Permanent and temporary effects assessed under CM1 are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Covered Activities and Associated Federal Actions, Section 4.2.1.1 North Delta Diversions Construction and Operations for a complete description of all activities assessed under CM1.

⁵ Borrow/Spoil Area: Borrow: location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil.

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																			
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss			
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction				
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
<p>⁶ Permanent and temporary effects assessed under CM2 include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.</p> <p>⁷ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>⁸ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>⁹ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>¹⁰ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>¹¹ Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>¹² Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>¹³ Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁴ Based on the hypothetical tidal restoration footprint, an estimated 2 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁵ Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁶ Reusable tunnel material is flexible and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.</p> <p>¹⁷ Because decimal places are not shown in this table, in some cases, a row total may be larger by one or two acres than the result obtained by manually summing numbers across columns.</p> <p>¹⁸ Although the tidal restoration model results in some decreases in acreage of natural community loss between near term and late long-term due to tidal damping and sea level rise, for permitting purposes the maximum acreage of loss is shown for late long-term.</p> <p>¹⁹ Loss reduced to zero. Although the temporary transmission powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools complex be avoided during transmission powerline installation.</p> <p>²⁰ Total permanent loss reduced from 201 acres (CM4) to 28 acres. This reduction is based on a 10-acre cap for total loss of wetted acres, assuming 15% density of vernal pools in the area affected. Acreage of vernal pool complex loss may be higher if actual vernal pool density is lower. The maximum acreage loss is based on loss of wetted acres and not total vernal pool complex acreage.</p> <p>NHD = National Hydrologic Database; SWP = State Water Project; CVP = Central Valley Project.</p>																			

1 **Table 5.J-13. Early Long-Term Plant Modeled Habitat Loss and Conversion by Covered Activity**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM4 Tidal Natural Communities Restoration													Plan Area Total ⁷	
		Suisun Marsh						Delta						Permanent (Acres) ^{7,8,9}	Conversion (Acres)	
		High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2			Subtidal 3
Plants																
Brittlescale total ¹³	451	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heartscale total	6,451	0	0	1	0	0	0	0	4	0	83	1	0	0	90	
San Joaquin spearscale total	14,477	0	0	0	0	0	0	0	123	0	255	1	0	0	380	
Carquinez goldenbush total ¹³	1,346	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Delta button celery total	3,361	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Delta mudwort total	6,081	0	0	0	0	0	0	0	0	0	0	3	1	2	6	
Mason's lilaepsis total	6,081	0	0	0	0	0	0	0	0	0	0	3	1	2	6	
Delta tule pea total ¹⁴	5,853	0	0	0	4	4	0	0	0	0	0	19	15	7	50 ²¹	
Suisun Marsh aster total ¹⁴	5,853	0	0	0	4	4	0	0	0	0	0	19	15	7	50 ²¹	
Side-flowering skullcap total ¹⁵	2,497	0	0	0	0	0	0	0	1	0	0	1	1	1	4	
Slough thistle total	1,834	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soft bird's-beak total	1,228	0	0	73	0	0	0	0	0	0	0	0	0	0	73	
Suisun thistle total	1,281	0	0	73	0	0	0	0	0	0	0	0	0	0	73	
Vernal Pool Plants																
Alkali milk-vetch																
Vernal pool complex	8,709	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	19	0	18	0	0	0	37	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38²⁰	0
Legenere																
Vernal pool complex	8,709	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	19	0	18	0	0	0	37	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38²⁰	0
Heckard's peppergrass																
Vernal pool complex	8,709	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	19	0	18	0	0	0	37	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38²⁰	0
Boggs Lake hedge-hyssop																
Vernal pool complex	8,709	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	19	0	18	0	0	0	37	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38²⁰	0
Dwarf downingia																
Vernal pool complex	8,709	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	19	0	18	0	0	0	37	0
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	0	0	0	0	0	0	0	19	0	18	0	0	0	38²⁰	0

2

1 **Table 5.J-13. Early Long-Term Plant Modeled Habitat Loss and Conversion by Covered Activity (cont'd)**

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶
Plants																		
Brittlescale total ¹³	451	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Heartscale total	6,451	0	0	0	0	0	0	0	0	0	0	0	0	0	0	90	0	0
San Joaquin spearscale total	14,477	23	30	0	29	56	0	0	1	0	0	0	0	0	0	488	0	30
Carquinez goldenbush total ¹³	1,346	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Delta button celery total	3,361	34	39	0	23	0	0	0	3	0	0	0	0	0	0	72	0	25
Delta mudwort total	6,081	12	3	0	15	3	2	0	0	0	0	0	0	0	0	24	0	17
Mason's lilaepsis total	6,081	12	3	0	15	3	2	0	0	0	0	0	0	0	0	24	0	17
Delta tule pea total ¹⁴	5,853	2	0	0	1	0	0	0	0	0	0	0	0	0	0	52	0	1
Suisun Marsh aster total ¹⁴	5,853	2	0	0	1	0	0	0	0	0	0	0	0	0	0	52	0	1
Side-flowering skullcap total ¹⁵	2,497	3	0	0	5	0	0	0	0	0	0	0	0	0	0	7	0	5
Slough thistle total	1,834	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	3
Soft bird's-beak total	1,228	0	0	0	0	0	0	0	0	0	0	0	0	0	0	73	0	0
Suisun thistle total	1,281	0	0	0	0	0	0	0	0	0	0	0	0	0	0	73	0	0
Vernal Pool Plants																		
Alkali milk-vetch																0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	44	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)	Temporary (Acres)
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Legenere																0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	44	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Heckard's peppergrass																0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	44	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Boggs Lake hedge-hyssop																0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	44	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0
Dwarf downingia																0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	44	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	53	0	0

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent (Acres) ⁴	Permanent - Reusable Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)
<p>¹ The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Non-federal Actions.</p> <p>² Existing habitat and habitat loss are estimated using habitat models created from detailed vegetation mapping. See Appendix 2.A, <i>Covered Species Accounts</i>, for a complete description of species-specific mapping methods. Effects on species' habitat will be tracked during implementation through on-the-ground surveys performed by qualified biologists.</p> <p>³ See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, <i>Effects on Natural Communities, Wildlife, and Plants</i>, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, <i>Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat</i>, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.</p> <p>⁴ Permanent and temporary effects assessed under CM1 are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Covered Activities and Associated Federal Actions, Section 4.2.1.1 North Delta Diversions Construction and Operations for a complete description of all activities assessed under CM1.</p> <p>⁵ Borrow/Spoil Area: Borrow: location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil.</p> <p>⁶ Permanent and temporary effects assessed under CM2 include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.</p> <p>⁷ Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>⁸ Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>⁹ Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>¹⁰ Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>¹¹ Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>¹² Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>¹³ Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁴ Based on the hypothetical tidal restoration footprint, an estimated 2 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁵ Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>¹⁶ Reusable tunnel material is flexible and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.</p> <p>¹⁷ Because decimal places are not shown in this table, in some cases, a row total may be larger by one or two acres than the result obtained by manually summing numbers across columns.</p> <p>¹⁸ Although the tidal restoration model results in some decreases in acreage of natural community loss between near term and late long-term due to tidal damping and sea level rise, for permitting purposes the maximum acreage of loss is shown for late long-term.</p> <p>¹⁹ Loss reduced to zero. Although the temporary transmission powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM3 requires that wetted acres of alkali seasonal wetlands and vernal pools complex be avoided during transmission powerline installation.</p> <p>²⁰ Total permanent loss reduced from 201 acres (CM4) to 28 acres. This reduction is based on a 10-acre cap for total loss of wetted acres, assuming 15% density of vernal pools in the area affected. Acreage of vernal pool complex loss may be higher if actual vernal pool density is lower. The maximum acreage loss is based on loss of wetted acres and not total vernal pool complex acreage.</p> <p>NHD = National Hydrologic Database; SWP = State Water Project; CVP = Central Valley Project.</p>																		

1 **Table 5.J-14. Late Long-Term Plant Modeled Habitat Loss and Conversion by Covered Activity**

Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}														Plan Area Total ⁷	
		CM4 Tidal Natural Communities Restoration									Delta					Permanent (Acres) ^{7,8,9}	Conversion (Acres)
		Suisun Marsh			Delta			Delta			Delta						
High Tidal Brackish Marsh	Mid Tidal Brackish Marsh	Low Tidal Brackish Marsh	Intertidal Mudflat	Subtidal 1	Subtidal 2	Subtidal 3	Ecotone	Intertidal Mudflat	Tidal Freshwater Marsh	Subtidal 1	Subtidal 2	Subtidal 3	Subtidal 1	Subtidal 2	Subtidal 3		
Plants																	
Brittlescale total ¹³	451	0	0	0	0	0	0	0	0	0	20	0	0	0	20 ²¹	0	
Heartscale total	6,451	1	0	1	0	0	0	0	38	0	253	7	6	0	306	0	
San Joaquin spearscale total	14,477	0	0	1	0	0	0	0	83	0	525	4	8	0	622	0	
Carquinez goldenbush total ¹³	1,346	0	0	0	0	0	0	0	0	0	50	0	0	0	50 ²¹	0	
Delta button celery total	3,361	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Delta mudwort total	6,081	0	0	0	0	0	0	0	0	0	0	2	2	2	6	0	
Mason's lilaeopsis total	6,081	0	0	0	0	0	0	0	0	0	0	2	2	2	6	0	
Delta tule pea total ¹⁴	5,853	0	0	0	4	4	0	0	0	0	0	19	15	7	50 ²¹	0	
Suisun Marsh aster total ¹⁴	5,853	0	0	0	4	4	0	0	0	0	0	19	15	7	50 ²¹	0	
Side-flowering skullcap total ¹⁵	2,497	0	0	0	0	0	0	0	1	0	0	1	1	1	4	0	
Slough thistle total	1,834	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Soft bird's-beak total	1,228	0	0	73	0	0	0	0	0	0	0	0	0	0	73	0	
Suisun thistle total	1,281	0	0	73	0	0	0	0	0	0	0	0	0	0	73	0	
Vernal Pool Plants																	
Alkali milk-vetch																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	9	0	41	1	0	0	51	0	
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52²⁰	0	
Legenere																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	9	0	41	1	0	0	51	0	
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52²⁰	0	
Heckard's peppergrass																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	9	0	41	1	0	0	51	0	
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52²⁰	0	
Boggs Lake hedge-hyssop																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	9	0	41	1	0	0	51	0	
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52²⁰	0	
Dwarf downingia																	
Vernal pool complex	8,709	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	
Degraded vernal pool complex	2,576	0	0	0	0	0	0	0	9	0	41	1	0	0	51	0	
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	0	0	1	0	0	0	0	9	0	41	1	0	0	52²⁰	0	

2

1 Table 5.J-14. Late Long-Term Plant Modeled Habitat Loss and Conversion by Covered Activity (cont'd)

Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)
Plants																		
Brittlescale total ¹³	451	0	0	0	0	0	0	0	0	0	0	0	0	0	20	0	0	
Heartscale total	6,451	0	0	0	0	0	0	0	0	0	0	0	0	0	306	0	0	
San Joaquin spearscale total	14,477	23	30	0	29	56	0	1	1	0	0	0	0	0	731	0	30	
Carquinez goldenbush total ¹³	1,346	0	0	0	0	0	0	0	0	0	0	0	0	0	50	0	0	
Delta button celery total	3,361	34	39	0	23	0	0	7	8	0	0	0	0	0	79	0	31	
Delta mudwort total	6,081	12	3	0	15	3	2	1	2	0	0	0	0	0	25	0	19	
Mason's lilaepsis total	6,081	12	3	0	15	3	2	1	2	0	0	0	0	0	25	0	19	
Delta tule pea total ¹⁴	5,853	2	0	0	1	0	0	0	0	0	0	0	0	0	52	0	2	
Suisun Marsh aster total ¹⁴	5,853	2	0	0	1	0	0	0	0	0	0	0	0	0	52	0	2	
Side-flowering skullcap total ¹⁵	2,497	3	0	0	5	0	0	1	1	0	0	0	0	0	8	0	6	
Slough thistle total	1,834	0	0	0	0	0	0	50 ²¹	6	0	0	0	0	0	50	0	6	
Soft bird's-beak total	1,228	0	0	0	0	0	0	0	0	0	0	0	0	0	73	0	0	
Suisun thistle total	1,281	0	0	0	0	0	0	0	0	0	0	0	0	0	73	0	0	
Vernal Pool Plants																		
Alkali milk-vetch																		
Vernal pool complex	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	
Degraded vernal pool complex	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	58	0	0	
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0	
Legenere																		
Vernal pool complex	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0	
Degraded vernal pool complex	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	58	0	0	
Alkali seasonal wetland	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0	


Maximum Allowable Habitat Loss by Covered Activity ^{1,2,3}																		
Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)
Heckard's peppergrass														0		0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	58	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
Boggs Lake hedge-hyssop														0		0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	58	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0
Dwarf downingia														0		0	0	0
<i>Vernal pool complex</i>	8,709	8	0	0	0	0	0	0	0	0	0	0	0	0	0	9	0	0
<i>Degraded vernal pool complex</i>	2,576	7	0	0	0	0	0	0	0	0	0	0	0	0	0	58	0	0
<i>Alkali seasonal wetland</i>	188	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	11,472	15	0	0	0	0	0	0	0	0	0	0	0	0	0	67	0	0

Maximum Allowable Habitat Loss by Covered Activity^{1,2,3}

Resource	Total Existing Modeled Habitat in the Plan Area (Acres) ²	CM1 Water Facilities and Operation				CM2 Yolo Bypass Fisheries Enhancement		CM5 Seasonally Inundated Floodplain Restoration		CM7 Riparian Natural Community Restoration		CM8 Grassland Restoration	CM10 Nontidal Marsh Natural Community Restoration	CM11 Natural Community Enhancement and Management	CM18 Conservation Hatcheries	Maximum Allowable Habitat Loss		
		Tunnel/Pipeline Facilities Construction				Fremont Weir and Yolo Bypass Improvements		Levee Construction		Riparian Restoration as Part of Tidal Natural Communities Restoration	Riparian Restoration as Part of Seasonal Floodplain Restoration		Construction and Inundation	Construction of Recreational-Related Facilities	Construction			
		Permanent (Acres) ⁴	Permanent - Reusable Tunnel Material (Acres) ¹⁷	Temporary (Borrow and Spoil) (Acres) ^{4,5}	Temporary (Acres) ⁴	Permanent (Acres) ⁶	Temporary (Acres) ⁶	Permanent (Acres) ¹⁰	Temporary (Acres) ¹⁰	Permanent (Acres)	Permanent (Acres) ¹¹		Permanent (Acres) ¹²	Permanent (Acres) ¹²	Permanent (Acres)	Permanent (Acres) ¹²	Permanent (Acres) ¹⁶	Temporary (Borrow and Spoil) (Acres)
<p>1 The following covered activities and associated federal actions (listed here by the header/category as described in Chapter 4, Covered Activities and Associated Federal Actions) are assumed not to have footprint impacts on natural communities or species habitat: Operations and Maintenance of Existing SWP Facilities; Power Generation Water Use - Mirant Delta, LLC activities; Activities to Reduce Contaminants; Activities to Reduce Predators and Other Sources of Direct Mortality; Monitoring and Research Programs; Emergency Actions; CVP Operations and Maintenance; and Joint Federal and Non-federal Actions.</p> <p>2 Existing habitat and habitat loss are estimated using habitat models created from detailed vegetation mapping, See Appendix 2.A, <i>Covered Species Accounts</i>, for a complete description of species-specific mapping methods. Effects on species' habitat will be tracked during implementation through on-the-ground surveys performed by qualified biologists.</p> <p>3 See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, <i>Effects on Natural Communities, Wildlife, and Plants</i>, for a description methods and assumptions relevant to estimating natural community loss by covered activity type and Table 5.J.3, <i>Key Assumptions Related to Tidal Restoration Effects on Covered Species Habitat</i>, for a list of assumptions used to determine permanent loss or conversion as a result of inundation caused by tidal restoration.</p> <p>4 Permanent and temporary effects assessed under CM1 are associated with construction of the following conveyance-related facilities: forebay, intake facilities, permanent access roads, shaft locations, and transmission lines. See Chapter 4, Covered Activities and Associated Federal Actions, Section 4.2.1.1 North Delta Diversions Construction and Operations for a complete description of all activities assessed under CM1.</p> <p>5 Borrow/Spoil Area: Borrow: location from where construction material, such as sand or clay, will be taken. Spoil: area where construction by-products, such as removed earth, will be placed and stored. Borrow/spoil: an area that will originally be used for borrow and then later be used for spoil.</p> <p>6 Permanent and temporary effects assessed under CM2 include activities associated with Fremont Weir improvements, Putah Creek realignment activities, Lisbon weir and fish crossing improvements, and Sacramento Weir improvements.</p> <p>7 Inundation is tidal flooding of existing wetland habitat as a result of tidal restoration actions. Inundation can cause permanent loss of habitat from either the removal of habitat or the conversion of one habitat type to another. See Table 5.J.1, <i>Quantitative Effects Analysis Methods and Assumptions</i>, in Appendix 5.J, for a description of relevant assumptions. All construction is assumed to occur within the inundation footprint.</p> <p>8 Permanent loss calculations are based on hypothetical tidal restoration designs and include those areas modeled by ESAPWA (Appendix 3.B, <i>BDCP Tidal Habitat Evolution Assessment</i>) to be below extreme high water elevation. See Table 5.J.1 in Appendix 5.J, for methods and assumptions used to apply the hypothetical footprint to determine effects.</p> <p>9 Tidal restoration is expected to include riparian restoration where elevations are favorable. Permanent loss from riparian restoration was determined by non-GIS methods. See Table 5.J.1, in Appendix 5.J, for a complete list of methods and assumptions.</p> <p>10 Calculation of effects based on hypothetical floodplain restoration designs. See Table 5.J.1 in Appendix 5.J, for details.</p> <p>11 Based on restoration design assumptions described in Appendix 5.E, <i>Habitat Restoration</i>, and effects analysis assumptions detailed in Table 5.J.1 in Appendix 5.J.</p> <p>12 Permanent loss was determined based on non-GIS methods described in Table 5.J.1 in Appendix 5.J.</p> <p>13 Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>14 Based on the hypothetical tidal restoration footprint, an estimated 2 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>15 Based on the hypothetical tidal restoration footprint, an estimated 4 acres of habitat will be lost or converted. However, to provide flexibility in implementation of tidal restoration projects, the take limit is set higher than the amount of loss estimated under the hypothetical footprint.</p> <p>16 Reusable tunnel material is flexible and the footprint used in the effects analysis is based on a worst case scenario: the actual area to be affected by reusable tunnel material storage will likely be less than the estimated acreage.</p> <p>17 Because decimal places are not shown in this table, in some cases, a row total may be larger by one or two acres than the result obtained by manually summing numbers across columns.</p> <p>18 Although the tidal restoration model results in some decreases in acreage of natural community loss between near term and late long-term due to tidal damping and sea level rise, for permitting purposes the maximum acreage of loss is shown for late long-term.</p> <p>19 Loss reduced to zero. Although the temporary transmission powerline footprint overlaps with 2 acres of alkali seasonal wetland complex and 16 acres of vernal pool complex in Conservation Zone 8, AMM30 requires that wetted acres of alkali seasonal wetlands and vernal pools complex be avoided during transmission powerline installation.</p> <p>20 Total permanent loss reduced from 372 acres (CM4) to 52 acres. This reduction is based on a 10-acre cap for total loss of wetted acres, assuming 15% density of vernal pools in the area affected. Acreage of vernal pool complex loss may be higher if actual vernal pool density is lower. The maximum acreage loss is based on loss of wetted acres and not total vernal pool complex acreage.</p> <p>21 To allow for flexibility in implementation and to address uncertainty related to the hypothetical restoration footprints, maximum loss from CM4 has been increased from 4 to 20 acres for brittlescale, 4 to 50 acres for Carquinez goldenbush, and from 1 to 50 acres for delta tule pea and Suisun marsh aster. Maximum loss from CM5 has been increased from 5 to 50 acres for slough thistle.</p> <p>NHD = National Hydrologic Database; SWP = State Water Project; CVP = Central Valley Project.</p>																		


1 **5.J.1 Reference**

- 2 Patterson, Laura. Staff Environmental Scientist, California Department of Water Resources. 2012b.
3 September 11, 2012—Email to Rebecca Sloan detailing methods and results of an analysis to
4 determine percent western pond turtle habitat in the National Hydrologic Dataset (NHD) in the
5 Plan Area.




K:\Projects\1\DWRI\00000_00_BDCP\mapdocs\Public\Draft\DWIR_Screencheck\Appendix_5\J\Fig_05J_01_Difference_Proposed_Existing_Yolo_Scenario1_20130904.mxd Date: 9/16/2013 Time: 3:37:52 PM 29391

Figure 5.J-1
Difference between Proposed and Existing
Yolo Bypass Flows: Scenario 1




K:\Projects\1\DWRI\00000_00_BDCP\mapdocs\Public\Draft\DWIR_Screencheck\Appendix_5\J\Fig_05J_02_Difference_Proposed_Existing_Yolo_Scenario2_20130904.mxd Date: 9/16/2013 Time: 3:57:40 PM 29391

Figure 5.J-2
Difference between Proposed and Existing
Yolo Bypass Flows: Scenario 2



K:\Projects\1\DWRI\00000_00_BDCP\mapdocs\Public\Draft\DWIR_Screencheck\Appendix_5.J\Fig_05J_03_Difference_Proposed_Existing_Yolo_Scenario3_20130904.mxd Date: 9/16/2013 Time: 3:35:02 PM 29391

Figure 5.J-3
 Difference between Proposed and Existing
 Yolo Bypass Flows: Scenario 3



K:\Projects\1\DWRI\00000_00_BDCP\mapdocs\Public\Draft\DWIR_Screencheck\Appendix_5\J\Fig_05J_04_Difference_Proposed_Existing_Yolo_Scenario4_20130904.mxd Date: 9/16/2013 Time: 3:48:26 PM 29391

Figure 5.J-4
 Difference between Proposed and Existing
 Yolo Bypass Flows: Scenario 4






Figure 5.J-5
Difference between Proposed and Existing
Yolo Bypass Flows: Scenario 5



K:\Projects\1\DWRI\00000_00_BDCP\mapdocs\Public\Draft\DWIR_Screencheck\Appendix_5\J\Fig_05J_06_Difference_Proposed_Existing_Yolo_Scenario6_20130904.mxd Date: 9/16/2013 Time: 3:53:27 PM 29391

Figure 5.J-6
 Difference between Proposed and Existing
 Yolo Bypass Flows: Scenario 6



K:\Projects\1\DWRI\00000_00_BDCP\mapdocs\Public\Draft\DWIR_Screencheck\Appendix_5\J\Fig_05J_07_Difference_Proposed_Existing_Yolo_Scenario7_20130904.mxd Date: 9/16/2013 Time: 3:53:38 PM 29391

Figure 5.J-7
 Difference between Proposed and Existing
 Yolo Bypass Flows: Scenario 7

1
2
3

Attachment 5J.A
**Construction-Related Nitrogen Deposition on
BDCP Natural Communities**

Date:	October 1, 2013
To:	Laura King Moon, Project Manager, BDCP California Department of Water Resources
From:	Paola Bernazzani, M.S., Senior Conservation Biologist, ICF International
Subject:	Construction-Related Nitrogen Deposition on BDCP Natural Communities

1

2 The primary purpose of this memorandum is to provide context and describe the potential effects of
3 nitrogen deposition from Bay Delta Conservation Plan (BDCP) covered activities with respect to
4 natural communities and associated plants and invertebrates in the Plan Area.

5 Introduction

6 BDCP construction activities will require the use of cars, trucks, and machinery that release small
7 amounts of atmospheric nitrogen through the combustion and emissions process associated with
8 motorized vehicles. Emissions will be largely limited to the construction phase of development,
9 which is anticipated to last approximately 9 years. Following combustion, reactive nitrogen is blown
10 downwind and deposited on the landscape, where it acts as a fertilizer (Bay Area Open Space
11 Council 2011) (see Exhibit 1 for details on the nitrogen deposition process). This depositional
12 nitrogen can affect biogeochemical processes and species composition in terrestrial ecosystems,
13 which are largely nitrogen limited (Pardo et al. 2011; Bay Area Open Space Council 2011). Nitrogen
14 can be directly absorbed by plant leaves or taken up by roots through the process of dry deposition,
15 the most common form of deposition in the Central Valley. Increased nitrogen favors nonnative
16 annual grasses and other weeds that crowd out native plants, change fire regimes, and displace rare
17 species adapted to low-nitrogen conditions.

18 Aquatic natural communities are not addressed in this memo because nitrogen deposition to Delta
19 waters from airborne sources is insignificant compared to other sources of nitrogen; in particular,
20 the ammonium from wastewater discharges and agricultural runoff. High concentrations of
21 ammonium are a concern in the Delta because the ammonium inhibits uptake of nitrate by
22 phytoplankton, contributing to declines in the production of phytoplankton at the base of the Delta's
23 pelagic food web (Wilkerson et al. 2006; Dugdale et al. 2007; Glibert et al. 2011) (Chapter 3, Section
24 3.5, *Important Regional Actions*, and Appendix 5.F, *Biological Stressors on Covered Fish*).

25 In California, there are several terrestrial natural communities known to be susceptible to the
26 biological effects of nitrogen deposition, including coastal sage scrub, desert scrub, and serpentine
27 grassland (Weiss 2006). Although the Plan Area does not contain any "susceptible" natural

1 communities, the following natural communities in the Plan Area may be sensitive to nitrogen
2 deposition (Figure 1) (Weiss 2006).

- 3 • Grasslands
- 4 • Vernal pools (includes vernal pools and alkali seasonal wetlands)
- 5 • Salt marsh (tidal brackish emergent wetland)
- 6 • Freshwater marsh (tidal freshwater emergent wetland)


7 The effects of nitrogen deposition on nonserpentine annual grasslands are similar to those on
8 serpentine grasslands, with increased nonnative, invasive plants displacing native grasses and
9 herbs. In addition, vernal pools and alkali seasonal wetlands appear to be particularly vulnerable to
10 overgrowth and invasion by nonnative, annual plants (Marty 2005). Weeds such as yellow
11 starthistle react positively to increased nitrogen availability because they have high growth
12 potential and can rapidly respond to increased nutrient levels. In general, salt and freshwater marsh
13 communities are nitrogen limited, and adding nitrogen could shift plant composition by affecting
14 plant productivity.

15 The covered activities that may deposit nitrogen, the potential effects on vulnerable communities in
16 the Plan Area, and the context for understanding the effects of nitrogen deposition from covered
17 activities are discussed below.

18 **Covered Activities**

19 Construction activities use trucks and other mechanized equipment that release atmospheric
20 nitrogen via fossil-fuel combustion. These activities will include the construction of the conveyance
21 facilities and restoration sites and associated operations and maintenance. The water conveyance
22 facilities will be constructed with three intakes located at the northern end of the Sacramento–San
23 Joaquin River Delta (Delta) with a transmission line to deliver power to the project. Construction is
24 scheduled to be completed in phases between 2016 and 2024. Construction-related nitrogen
25 emissions will originate primarily from construction equipment and employee vehicle exhaust and
26 concrete batching from onsite plants. The highest levels of nitrogen emissions are expected at utility
27 and construction sites along the tunnel conveyance alignment. These emissions will be temporary
28 and will cease when construction activities are completed.

1



2

3

Figure 1. Potential BDCP Sources of Nitrogen and Natural Communities that May Be Sensitive to Effects of Nitrogen Deposition

1 Restoration activities under *CM2 Yolo Bypass Fisheries Enhancement*, *CM3 Natural Communities*
 2 *Protection and Restoration*, *CM4 Tidal Natural Communities Restoration*, *CM5 Seasonally Inundated*
 3 *Floodplain Restoration*, *CM6 Channel Margin Enhancement*, *CM7 Riparian Natural Community*
 4 *Restoration*, *CM8 Grassland Natural Community Restoration*, *CM9 Vernal Pool and Alkali Seasonal*
 5 *Wetland Complex Restoration*, and *CM10 Nontidal Marsh Restoration* require the use of construction
 6 equipment and would also generate small amounts of atmospheric nitrogen. Emissions would result
 7 from temporary earth-moving activities that require the use of heavy equipment and from ongoing
 8 restoration or monitoring activities that result in additional traffic on roads and highways in and
 9 around Suisun Marsh and the Yolo Bypass. The amount of emissions from these activities will be
 10 negligible, and emissions resulting from restoration activities were not modeled as part of the air
 11 quality analysis.

12 In addition, operations and maintenance activities could result in nitrogen emissions originating
 13 from vehicle and maintenance equipment exhaust and electrical generation. In general, future
 14 emissions are anticipated to decrease because of continuing improvements in vehicle and
 15 equipment engine technology. Operations and maintenance activities and construction at
 16 restoration sites would contribute a negligible amount of nitrogen relative to construction of
 17 conveyance facilities and other ongoing sources of nitrogen in the Central Valley (see *Baseline*
 18 *Conditions* section). For discussion purposes, this memo focuses on emissions from conveyance
 19 facilities construction.


20 **Baseline Conditions**

21 In 2002, modeling of nitrogen deposition in California estimated deposition rates of up to
 22 15 kilograms nitrogen (kg-N)/hectare (ha)/year (yr) (81.7 pounds nitrogen (lbs-N)/acre/yr) from
 23 urban and agricultural sources. The Central Valley is recognized as a hotspot of nitrogen deposition,
 24 with deposition values in the Plan Area ranging from 2.1 to 10 kg-N/ha/yr (11.4 to 54.5
 25 lbs/acre/yr). The southern portion of the Sacramento Valley received 6 to 8 kg-N/ha/yr (32.7 to
 26 43.6 lbs/acre/yr). Areas around Modesto (near but outside the Plan Area) received up to 14 kg-
 27 N/ha/yr (76.3 lbs/acre/yr), and in the Bay Area the maximum deposition was 9 kg-N/ha/yr
 28 (76.3 lbs/acre/yr) (Weiss 2006).

29 Nitrogen deposition above 5 kg-N/ha/yr (27.2 lbs/acre/yr) is known to result in exotic grass
 30 invasion on serpentine soils, and similar effects are expected for other annual grassland ecosystems
 31 and vernal pools in California (Weiss 2006; Fenn et al. 2010). Based on these analyses, current
 32 sources of nitrogen have already exceeded these thresholds in and around the Plan Area.

33 Deposition studies have not been done with respect to future emissions for the BDCP or for the Plan
 34 Area *per se*. However, the *California Almanac of Emissions and Air Quality* (2009) provides current
 35 and future values for the average annual emissions of the three air quality districts (Table 1) that
 36 overlap with the proposed conveyance facilities under Alternative (Figure 2).

1



2

3

Figure 2. Air Quality Districts in Plan Area

1 **Table 1. Previous and Estimated Future Nitrogen Oxide Emissions**

Year	Nitrogen Oxide Annual Average Emissions (tons/day) By Air Quality District		
	Bay Area	Sacramento	San Joaquin
2005	488	292	595
2010	414	249	524
2015	335	198	398
2020	284	161	316

Source: California Environmental Protection Agency 2009.

2

3 As mentioned above, there are four land cover types identified by Weiss (2006) as potentially
 4 vulnerable in the Plan Area: vernal pools, grasslands, saltwater marsh (tidal brackish emergent
 5 wetland) and freshwater marsh (tidal freshwater emergent wetland). The Plan Area supports an
 6 estimated 8,547 acres of vernal pools, 78,624 acres of grasslands, 8,501 acres of tidal brackish
 7 wetlands, and 8,953 acres of freshwater emergent wetlands. The location of these natural
 8 communities is mapped in relation to the proposed conveyance facilities, powerlines, and primary
 9 access routes for construction vehicles (Figure 1). The potential effects of deposition on these
 10 natural communities from construction of conveyance facilities is discussed below.

11 **Potential Effects**

12 **Natural Communities and Species in the Plan Area**

13 Generally, invasive nonnative plants may compete with native plants for water, nutrients, light, and
 14 germination sites, and invasive plants are considered a threat to most of the covered plant species.
 15 Perennial pepperweed (*Lepidium latifolium*) is a specific threat to soft bird's-beak (*Chloropyron*
 16 *molle* subsp. *molle*) and Suisun thistle (*Cirsium hydrophilum* var. *hydrophilum*). Perennial ryegrass
 17 (*Festuca perennis*, formerly *Lolium perenne*) invades seasonally moist grasslands, seasonal wetlands,
 18 and vernal pools, and is a threat to legumere (*Legenere limosa*), alkali milk-vetch (*Astragalus tener*
 19 var. *tener*), and other covered plants of these habitats. The invasive aquatic plant water hyacinth
 20 (*Eichhornia crassipes*), which forms large mats of floating vegetation, is a specific threat to Mason's
 21 lilaepsis (Zebell and Fiedler 1996). The potential effects of nitrogen deposition are described by
 22 natural community type below.

23 **Vernal Pools**

24 Small, annual plants in vernal pools and alkali seasonal wetlands are susceptible to overgrowth by
 25 invasive grasses, which can shorten hydroperiods and place associated species at risk (Marty 2005).
 26 Annual grass invasions in vernal pools have been documented in the Sacramento Valley (Gerhardt
 27 and Collinge 2003) and may be a major threat to ungrazed vernal pools (Marty 2005). Given the
 28 responses of annual grasses to additional nitrogen, the intensity of annual grass invasions in vernal

1 pools and alkali seasonal wetlands may increase as the result of increased nitrogen deposition in the
 2 Plan Area. Covered plants and invertebrates in vernal pool habitats include the conservancy fairy
 3 shrimp (*Branchinecta conservatio*), vernal pool tadpole shrimp (*Lepidurus packardii*), longhorn fairy
 4 shrimp (*Branchinecta longiantenna*), vernal pool fairy shrimp (*Branchinecta lynchi*), California
 5 linderiella (*Linderiella occidentalis*), midvalley fairy shrimp (*Branchinecta mesovallensis*), alkali milk-
 6 vetch (*Astragalus tener* var. *tener*), Boggs Lake hedge-hyssop (*Gratiola heterosepala*), delta button
 7 celery (*Eryngium racemosum*), dwarf downingia (*Downingia pusilla*), Heckard's peppergrass
 8 (*Lepidium latipes* var. *heckardii*), brittlescale (*Atriplex depressa*), heartscale (*Atriplex cordulata*), San
 9 Joaquin spearscale (*Atriplex joaquiniana*), and legenere (*Legenere limosa*).

10 Annual Grasslands

11 Although California grasslands are dominated by invasive annual grasses, they do support
 12 wildflower and native bunchgrass grassland concentrations. Increased levels of nitrogen deposition
 13 can stimulate annual grass growth, thus further adversely affecting these native concentrations,
 14 particularly in areas where the soils are nutrient poor, such as on rocky outcrops or steep slopes
 15 (Weiss 2006).

16 Tidal Brackish Emergent Wetlands

17 Productivity in salt marshes (tidal brackish emergent wetland) is limited by nitrogen (Morris 1991).
 18 While salt marshes are major sites for denitrification, additional inputs of nitrogen may exceed the
 19 capacity of salt marshes to remove nitrogen from the system and could subsequently alter species
 20 assemblages associated with this natural community type. Many salt marshes are already subject to
 21 elevated nitrogen due to sewage effluent and agricultural runoff, and, while the direct effects of
 22 atmospheric nitrogen deposition on California salt marshes have not been assessed, additional
 23 inputs of nitrogen are likely to exacerbate issues associated with invasive plants. Covered plants in
 24 tidal brackish habitats include the Suisun thistle (*Cirsium hydrophilum* var. *hydrophilum*), soft bird's
 25 beak (*Cordylanthus mollis* ssp. *mollis*), Suisun Marsh aster (*Symphyotrichum lentum*), Delta tule pea
 26 (*Lathyrus jepsonii* var. *jepsonii*), Mason's lilaeopsis (*Lilaeopsis masonii*), and delta mudwort
 27 (*Limosella subulata*).

28 Tidal Freshwater Emergent Wetlands

29 Productivity in freshwater marshes (tidal freshwater emergent wetlands) may be limited by
 30 nitrogen (Morris 1991). Because of anoxic conditions and an abundance of organic matter,
 31 freshwater marshes, like salt marshes described above, remove nitrogen from the system and may
 32 be altered by an abundance of nitrogen. Covered plants in freshwater emergent wetland habitats
 33 include the Suisun Marsh aster (*Symphyotrichum lentum*), Delta tule pea (*Lathyrus jepsonii* var.
 34 *jepsonii*), Mason's lilaeopsis (*Lilaeopsis masonii*), and delta mudwort (*Limosella subulata*).

35 Analysis and Discussion

36 Maximum daily emissions associated with covered activities are presented in Chapter 22 of the
 37 Draft BDCP environmental impact report/environmental impact statement (EIR/EIS) (California

1 Department of Water Resources et al. 2012) for the three air quality districts that overlap with the
 2 conveyance facilities in Alternative 4. Results (originally in pounds/day) were converted to tons/day
 3 and are summarized in Table 2.

4 **Table 2. Projected Maximum Daily Nitrogen Oxide Emissions from Construction of Conveyance**
 5 **Facilities**

Year	Maximum Daily Nitrogen Oxide Emissions (tons/day) By Air Quality District		
	Bay Area	Sacramento	San Joaquin
Source: Draft BDCP EIR/EIS Chapter 22, Table 22-86, for Alternative 4 (converted from pounds/day to tons/day) (California Department of Water Resources et al. 2012)			

6
 7 The projected total nitrogen emissions (Table 1) are compared to projected nitrogen emissions from
 8 covered activities (Table 2) to quantify the relative contribution of covered activities to estimated
 9 regional emissions of nitrogen.

10 Table 3 lists the BDCP contribution by percentage for the three air quality districts that overlap with
 11 the proposed conveyance facilities. Results are given for the years 2015/2016 and 2020¹. In all
 12 cases, the BDCP contribution of nitrogen deposition to the estimated annual average total is less
 13 than 0.2%, with concentrations in most basins less than 0.08%.

14 **Table 3. Projected Annual Average Nitrogen Emissions from the BDCP as a Percentage**
 15 **of Total Projected Emissions, by Air Quality District**

Year	Percentage BDCP Emissions		
	Bay Area	Sacramento	San Joaquin
2015/2016	0.002	0.074	0.024
2020	0.164	0.122	0.041
Sources: California Department of Water Resources et al. 2012 (Chapter 22, Table 22-86, for Alternative 4). California Environmental Protection Agency, Air Resources Board 2009.			

16
 17 In addition, there is a considerable distance between covered activities that will temporarily emit
 18 nitrogen and most covered natural communities potentially sensitive to nitrogen (Figure 1 and
 19 Table 4). Furthermore, the direction of the prevailing winds is west to east (Western Regional
 20 Climate Center 2012). With most grasslands, vernal pools, and marshes in the Plan Area lying west
 21 of the conveyance facilities, most emissions will be transported away from areas of potential
 22 concern. The exception is the Stone Lakes Wildlife Refuge complex, located east of the conveyance
 23 facilities and discussed further below.

¹ The only year that specifically overlaps both the BDCP emissions analysis and the *California Almanac of Emissions and Air Quality* value is 2020. The years 2015 (California Environmental Protection Agency, Air Resources Board. 2009) and 2016 (BDCP Draft EIR/EIS [California Department of Water Resources et al. 2012]) were also compared, since the BDCP projections begin year 2016 (not 2015).

Table 4. Communities that May be Sensitive to Nitrogen Deposition within 5 Kilometers of Proposed Conveyance Facilities

Natural Community Type	Acres	Percent of Total
Vernal pools (includes alkali seasonal wetlands)	1,770	14
Annual grassland	16,716	21
Tidal brackish emergent wetlands	0	0
Tidal freshwater emergent wetlands	1,684	19

With respect to potential effects on natural communities, the following observations are made.

- In the Plan Area, the grassland natural community is often found adjacent to wetland and riparian natural communities. As indicated in Figure 1 and Table 4, most of this community in the Plan Area is over 5 kilometers from emissions locations and west of the proposed facilities. The Byron Hills Area is directly adjacent to proposed construction at the southern end of the facilities. Significant grassland areas that include vernal pools and alkali seasonal wetlands are located here. Temporary nitrogen deposition resulting from construction could affect grasslands in the Byron Hills. However, prevailing winds in this area will likely blow most deposition away from grasslands in that area. Also, the Byron Hills area is a target for acquisition and management, including weed management through grazing, which will likely offset any effects of increased deposition.
- The vernal pool complex, including alkali seasonal wetlands, and associated grasslands are rare in the Plan Area and generally found only in a few locations along the margins of the Plan Area, including the Stone Lakes National Wildlife Refuge, adjacent to and east of proposed construction of conveyance facilities. The North Stone Lake unit of the Stone Lake Wildlife Refuge contains one of the only remaining undeveloped grassland units in the eastern Delta region (U.S. Fish and Wildlife Service 2007), as well as large complexes of vernal pools. Based on proximity to the facilities and its location downwind of construction, this area could be affected by the temporary increases in nitrogen deposition associated with conveyance construction. However, weed control and targeted grazing in the refuge are anticipated to control invasive plants, which might proliferate in an ungrazed system. Grazing throughout the refuge is conducted from November through June to reduce competition between vernal pool plants and nonnative species such as annual ryegrass and yellow starthistle, in accordance with the *Stone Lake Comprehensive Conservation Plan* (U.S. Fish and Wildlife Service 2007).
- Remnant patches of tidal brackish and freshwater emergent wetland natural community are found in the western portion of the Delta. Small patches of tidal brackish marshes are found on islands west of Sherman Island and in Suisun Marsh. Freshwater emergent wetlands are found near the confluence of the Sacramento and San Joaquin Rivers, along Lindsey Slough and the Yolo Bypass, along the mainstem and several channels of the San Joaquin, Old, and Middle Rivers, Lost Slough, and the area where the Cosumnes and Mokelumne Rivers join the Delta. Most of these areas are from 5 to 20 kilometers west of the proposed conveyance facilities and powerlines and are unlikely to experience significant negative effects from temporary, construction-related nitrogen deposition.

1 Conclusions

2 Nitrogen emissions from covered activities will not negatively affect natural communities and
3 covered species in the Plan Area for the following reasons.

- 4 • The covered activities will make a negligible contribution to projected emissions in the region
5 (less than 0.2%).
- 6 • The construction activities will be temporary (less than 9 years).
- 7 • There is a substantial distance between the nitrogen sources and potentially sensitive
8 communities.
- 9 • Nitrogen emissions will be transported away from most sensitive communities in the Plan Area
10 because of prevailing wind conditions.
- 11 • In the grassland and vernal pool natural community portion of the Stone Lakes Wildlife Refuge,
12 where negative effects are most likely to occur, a weed control and grazing plan are already in
13 place.

14 Moreover, planned management of the BDCP reserve system (*CM11 Natural Communities*
15 *Enhancement and Management*), which includes invasive vegetation control measures, is expected to
16 minimize the potential adverse effects of nitrogen deposition on protected grasslands, vernal pools,
17 and marshes in the Plan Area.

18 References Cited

19 Bay Area Open Space Council. 2011. *The Conservation Lands Network: San Francisco Bay Area Upland*
20 *Habitat Goals Project Report*. Berkeley, CA

21 California Department of Water Resources, Bureau of Reclamation, U.S. Fish and Wildlife Service,
22 and National Marine Fisheries Service. 2012. *Environmental Impact Report/Environmental*
23 *Impact Statement for the Bay Delta Conservation Plan*. Prepared by ICF International.
24 Sacramento, CA. February.

25 California Environmental Protection Agency, Air Resources Board. 2009. *California Almanac of*
26 *Emissions and Air Quality*. Air Resources Board. Last revised: January 14, 2010. Available:
27 <<http://www.arb.ca.gov/aqd/almanac/almanac09/almanac09.htm>>.

28 Dugdale, R. C., F. P. Wilkerson, V. E. Hogue, and A. Marchi. 2007. The Role of Ammonium and Nitrate
29 In Spring Bloom Development in San Francisco Bay. *Estuarine, Coastal, and Shelf Science* 73:17-
30 29.

31 Fenn, M. E., E. B. Allen, S. B. Weiss, S. Jovan, L. H. Geiser, G. S. Tonnesen, R. F. Johnson, L. E. Rao, B. S.
32 Gimeno, F. Yuan, T. Meixner, and A. Bytnerowicz. 2010. Nitrogen Critical Loads and Management
33 Alternatives for N-Impacted Ecosystems in California. *Journal of Environmental Management*
34 91:2402-2423.


- 1 Fenn, M. E., J. S. Baron, E. B. Allen, H. M. Rueth, K. R. Nydick, L. Geiser, W. D. Bowman, J. O. Sickman, T.
2 Meixner, D. W. Johnson, and P. Neitlich. 2003. Ecological Effects of Nitrogen Deposition in the
3 Western United States. *Bioscience* 53:404–420.
- 4 Fenn M. E., M. A. Poth,, J. D. Aber, J. S. Baron, B. T. Bormann, D. W. Johnson, A. D. Lemly, S. G. McNulty,
5 D. F. Ryan, and R. Stottlemeyer. 1998. Nitrogen Excess in North American Ecosystem:
6 Predisposing Factors, Ecosystem Responses, and Management Strategies. *Ecological Applications*
7 8:706–733.
- 8 Galloway, J. N., J. D. Aber, J. W. Erisman, S. P. Seitzinger, R. W. Howarth, E. B. Cowling, and B. J. Cosby.
9 2003. The Nitrogen Cascade. *BioScience* 53:341–356.
- 10 Gerhardt, F. and S. K. Collinge. 2003. Exotic Plant Invasions of Vernal Pools in the Central Valley of
11 California, USA. *Journal of Biogeography* 30(7): 1043–1052.
- 12 Glibert, P. M., D. Fullerton, J. M. Burkholder, J. C. Cornwell, and T. M. Kana. 2011. Ecological
13 Stoichiometry, Biogeochemical Cycling, Invasive Species and Aquatic Food Webs: San Francisco
14 Estuary and Comparative Systems. *Reviews in Fisheries Science* 19:358–417.
- 15 Marty, J. 2005. Effects of Cattle Grazing on Diversity in Ephemeral Wetlands. *Conservation Biology*
16 19:1626–1632.
- 17 Matson, P. K., A. Lohse, et al. 2002. The Globalization of Nitrogen Deposition: Consequences for
18 Terrestrial Ecosystems. *AMBIO: A Journal of the Human Environment* 31(2): 113–119.
- 19 Morris, J. T. 1991. Effects of Nitrogen Loading on Wetland Ecosystems with Particular Reference to
20 Atmospheric Deposition. *Annual Review of Ecology and Systematics* 22: 257–279.
- 21 Pardo, L. H., M. J. Robin-Abbott, and C. T. Driscoll (eds.). 2011. *Assessment of Nitrogen Deposition*
22 *Effects and Empirical Critical Loads of Nitrogen for Ecoregions of the United States*. U.S.
23 Department of Agriculture, Forest Service, Northern Research Station, General Technical Report
24 NRS-80.
- 25 U.S. Fish and Wildlife Service. 2007. *Stone Lakes National Wildlife Refuge Comprehensive*
26 *Conservation Plan*. January. USFWS California Nevada Operations Refuge Planning Office,
27 Sacramento, CA, and Stone Lakes National Wildlife Refuge, Elk Grove, CA. Available:
28 <<http://www.fws.gov/stonelakes/SL%20CCP%20Final%20main%20body.pdf>>. Accessed:
29 November 2012.
- 30 Weiss, S. B. 2006. *Impacts of Nitrogen Deposition on California Ecosystems and Biodiversity*. California
31 Energy Commission, PIER Energy-Related Environmental Research. CEC-500-2005-165.
- 32 Western Regional Climate Center. 2012: *Prevailing Wind Direction*. Available:
33 <<http://www.wrcc.dri.edu/htmlfiles/westwinddir.html>>. Accessed: November 2012.
- 34 Wilkerson, F. P., R. C. Dugdale, V. E. Hogue, and A. Marchi. 2006. Phytoplankton Blooms and Nitrogen
35 Productivity in San Francisco Estuary. *Estuaries and Coasts* 29: 401–416.
- 36 Zebell, R. and P. Fiedler. 1996. *Restoration and Recovery of Mason's lilaeopsis: Phase II*. Final Report
37 to the California Department of Fish and Game, Plant Conservation Program.
- 38

Exhibit 1

N-Deposition Pathway

Most of the nitrogen in the atmosphere is in the form of the inert nitrogen gas, dinitrogen (N₂). The primary emissions of nitrogen from anthropogenic sources are nitrogen oxide (NO) and nitrogen dioxide (NO₂), collectively referred to as nitrogen oxides (NO_x). Chemical processes in the atmosphere convert NO and NO₂ to other forms of nitrogen. Of particular interest is the formation of nitric acid (HNO₃), which reacts with ammonia in the atmosphere to form ammonium nitrate, which is biologically available nitrogen deposited to the ground (Matson et al. 2002). In a process known as nitrification, soil microbes oxidize the ammonium (NH₄⁺) in ammonium nitrate to produce nitrite (NO₂⁻), which is metabolized further to produce nitrate (NO₃⁻), the form of nitrogen assimilated by plants. Denitrifying bacteria convert nitrate back to nitrogen gas, providing a pathway for nitrogen cycling back to the atmosphere.

Studies have shown a range of ecosystem responses to elevated inputs of nitrogen from atmospheric deposition, with particular responses depending on multiple, interacting factors such as climate, land use, the ecosystem's current nitrogen status, and the extent and level of nitrogen additions (Fenn et al. 1998; Matson et al. 2002; Fenn et al. 2003; Pardo et al. 2011). For example, nitrogen-poor ecosystems like grasslands tend to accumulate additional nitrogen, while wetlands have a high capacity for removing nitrogen through denitrification (Galloway et al. 2003).



Note: Biological processes are labeled in bold italics, and the lighter arrows show deposition pathways.

Source: Figure 1 from Weiss 2006.

Figure A. Simplified Nitrogen Cycle

1 Exhibit 1 References Cited

- 2 Fenn, M. E., J. S. Baron, E. B. Allen, H. M. Rueth, K. R. Nydick, L. Geiser, W. D. Bowman, J. O. Sickman, T.
3 Meixner, D. W. Johnson, and P. Neitlich. 2003. Ecological Effects of Nitrogen Deposition in the
4 Western United States. *Bioscience* 53:404–420.
- 5 Fenn M. E., M. A. Poth,, J. D. Aber, J. S. Baron, B. T. Bormann, D. W. Johnson, A. D. Lemly, S. G. McNulty,
6 D. F. Ryan, and R. Stottlemyer. 1998. Nitrogen Excess in North American Ecosystem:
7 Predisposing Factors, Ecosystem Responses, and Management Strategies. *Ecological Applications*
8 8:706–733.
- 9 Galloway, J. N., J. D. Aber, J. W. Erisman, S. P. Seitzinger, R. W. Howarth, E. B. Cowling, and B. J. Cosby.
10 2003. The Nitrogen Cascade. *BioScience* 53:341–356.
- 11 Matson, P. K., A. Lohse, et al. 2002. The Globalization of Nitrogen Deposition: Consequences for
12 Terrestrial Ecosystems. *AMBIO: A Journal of the Human Environment* 31(2): 113–119.
- 13 Pardo, L. H., M. J. Robin-Abbott, and C. T. Driscoll (eds.). 2011. *Assessment of Nitrogen Deposition*
14 *Effects and Empirical Critical Loads of Nitrogen for Ecoregions of the United States*. U.S.
15 Department of Agriculture, Forest Service, Northern Research Station, General Technical Report
16 NRS-80.
- 17 Weiss, S. B. 2006. *Impacts of Nitrogen Deposition on California Ecosystems and Biodiversity*. California
18 Energy Commission, PIER Energy-Related Environmental Research. CEC-500-2005-165.
19 Appendix B is available from: <www.energy.ca.gov/.../CEC-500.../CEC-500-2005-165-APB.XLS>.

1
2
3

Attachment 5J.B
**Natural Community Restoration and Protection
Contributing to Covered Species Conservation**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

Attachment 5J.B

**Natural Community Restoration and Protection
Contributing to Covered Species Conservation**

Contents

	Page
Attachment 5J.B Natural Community Restoration and Protection Contributing to Covered Species Conservation.....	5J.B-1

Tables

	Page
5J.B-1 Natural Community Restoration Contributing to Covered Species Conservation— Wildlife.....	5J.B-3
5J.B-2 Natural Community Protection Contributing to Covered Species Conservation— Wildlife.....	5J.B-13
5J.B-3 Natural Community Restoration Contributing to Covered Species Conservation— Plants.....	5J.B-22
5J.B-4 Natural Community Protection Contributing to Covered Species Conservation— Plants.....	5J.B-25

Attachment 5J.B

Natural Community Restoration and Protection Contributing to Covered Species Conservation

The biological goals and objectives have been designed to provide species conservation through the protection and restoration of the natural communities on which those species depend. Planning at the natural community scale benefits species by protecting ecosystem processes that maintain species' habitat and by preserving connectivity between species' habitat types (e.g., between foraging and breeding habitat). However, species habitat is often comprises portions of one or more natural communities. This approach makes calculating total, net benefits for any one species difficult when there are no species-specific habitat conservation requirements. There are two main complications with estimating net benefits: 1) how to determine what portion of any given natural community conservation commitment will benefit any one species, and 2) to what species can you apply any given natural community conservation commitment. As an example, Swainson's hawk forages in many different habitat types (e.g., cultivated lands, managed wetlands, grasslands). There are specific Swainson's hawk conservation commitments for cultivated lands and grasslands but not for managed wetlands. Swainson's hawk will most assuredly benefit from managed wetland protection, but quantifying the benefit is difficult. The approach used to estimate species-specific benefits from a natural community objective is described in the paragraphs below. Table 5J.B-1 and Table 5J.B-2 provide the restoration and protection benefit estimates for wildlife, respectively, and Table 5J.B-3 and Table 5J.B-4 present these same results for plants. The species-specific benefit conclusion for each natural community model type is presented in bold text in one of the last three columns of each table.

The approach to estimating net benefits included the five basic steps described below.

1. Of the total modeled habitat acreage, quantify the contribution of each natural community. This is done in GIS where the habitat model is intersected with the natural community layer. These results are presented in each table below in the *Acreage of Modeled Habitat Comprising the Natural Community* column.
2. Determine what proportion (or percentage) of the natural community is included in the species modeled habitat. This is done by dividing the amount of the natural community that overlaps with the species model, described in step 1 above, by the total amount of the natural community in the Plan Area (presented in the *Total Acres of Natural Community in the Plan Area* column of each table below). The result is presented in the column titled *Percentage of Modeled Habitat Comprising the Natural Community*.
3. Identify whether a natural community restoration or protection objective will contribute to conservation of the species in question. For example, there is a managed wetland objective to protect 8,100 acres of managed wetlands in Suisun Marsh. The greater sandhill crane habitat model includes managed wetlands; however, the crane's range does not extend to Suisun Marsh, so this natural community objective would not be applied to the greater sandhill crane.
4. Calculate the estimated contribution of natural community protection or restoration to species conservation. This is done by multiplying the acreage of natural community restoration and

1 protection by the *Percentage of Modeled Habitat Comprising the Natural Community*. The result
2 is presented in the fifth column.

3 5. Identify the natural community or species-specific objectives that were created specifically to
4 benefit the species in question. These acreages are placed in one of the last two columns in each
5 table, depending on whether the objective is specific to a natural community or species.

6 Whenever a natural community or species-specific objective is identified as benefiting a species, the
7 entirety of that acreage commitment is counted as a benefit for the species. The estimated
8 contribution of natural community restoration or protection is only used in the absence of a specific
9 natural community or species objective. The acreage number chosen to estimate total benefits is
10 presented in bold text.

11 The natural community benefit estimates for each model type are totaled and carried forward to the
12 *BDCP Conservation* columns of the wildlife and plant net effects tables in Chapter 5, *Effects Analysis*,
13 Table 5.6-7 and Table 5.6-8, respectively. Estimated benefits are also discussed in the *Beneficial*
14 *Effects* section of each covered wildlife and plant species (see Section 5.6, *Effects on Covered Wildlife*
15 *and Plant Species*, for more details).

1 Table 5J.B-1. Natural Community Restoration Contributing to Covered Species Conservation—Wildlife

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Mammals							
Riparian brush rabbit							
<i>Riparian Habitat</i>							
Valley/Foothill Riparian	2,909	17,966	16.2	5,000	809		800
<i>Grassland Habitat</i>							
Grassland	3,094	78,047	4.0	2,000	79		
Riparian woodrat							
<i>Habitat</i>							
Valley/Foothill Riparian	2,166	17,966	12.1	5,000	603		300
Salt marsh harvest mouse							
<i>Managed Wetland—Upland Low, Long-Term Conservation Value</i>							
Managed Wetland	3,787	70,798	5.3	0	0		
<i>Managed Wetland—Wetland Primary Low, Long-Term Conservation Value</i>							
Managed Wetland	21,891	70,798	30.9	0	0		
<i>Managed Wetland—Wetland Secondary Low, Long-Term Conservation Value</i>							
Managed Wetland	2,800	70,798	4.0	0	0		
<i>Tidal Brackish Emergent Wetland Primary</i>							
Tidal Brackish Emergent Wetland	3,342	8,501	39.3	1,500	590	1,500	
<i>Tidal Brackish Emergent Wetland Secondary</i>							
Tidal Brackish Emergent Wetland	2,718	8,501	32.0	4,500	1,439	4,500	
<i>Upland Secondary</i>							
Grassland	491	78,047	0.6	2,000	13		
Tidal Brackish Emergent Wetland	189	8,501	2.2	1,500	33		
San Joaquin kit fox							
<i>Breeding, Foraging, and Dispersal Habitat</i>							
Grassland	5,098	78,047	6.5	2,000	131		
Vernal Pool Complex	229	12,132	1.9	67	1		
Suisun shrew							
<i>Primary Habitat</i>							

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Grassland	12	78,047	0.0	2,000	0		
Tidal Brackish Emergent Wetland	3,001	8,501	35.3	1,500	530	1,500	
<i>Secondary Habitat</i>							
Grassland	219	78,047	0.3	2,000	6		
Managed Wetland	1,825	70,798	2.6	0	0		
Tidal Brackish Emergent Wetland	2,181	8,501	25.7	4,500	1,155	4,500	
Birds							
California black rail							
<i>Primary Habitat</i>							
Nontidal Freshwater Perennial Emergent Wetland	715	1,509	47.4	800 ^a	379		
Tidal Brackish Emergent Wetland	3,760	8,501	44.2	1,500	664	1,500	
Tidal Freshwater Emergent Wetland	1,458	8,856	16.5	24,000	3,951		1,700
<i>Secondary Habitat</i>							
Managed Wetland	12,957	70,798	18.3	0	0		
Nontidal Freshwater Perennial Emergent Wetland	66	1,509	4.4	800 ^a	35		
Tidal Brackish Emergent Wetland	2,022	8,501	23.8	4,500	1,070	4,500	
Tidal Freshwater Emergent Wetland	2,797	8,856	31.6	24,000	7,580		
California clapper rail							
<i>Primary Habitat</i>							
Tidal Brackish Emergent Wetland	248	8,501	2.9	1,500	44	1,500	
<i>Secondary Habitat</i>							
Tidal Brackish Emergent Wetland	5,324	8,501	62.6	4,500	2,818	4,500	
Tidal Freshwater Emergent Wetland	753	8,856	8.5	0 ^b	0		
Greater sandhill crane							
<i>Roosting - Permanent</i>							
Cultivated land	5,237	487,106	1.1	75	1		75
Grassland	628	78,047	0.8	0 ^c	0		
Managed Wetland	1,097	70,798	1.5	500	8		500

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Nontidal Freshwater Perennial Emergent Wetland	43	1,509	2.9	0 ^c	0		
Nontidal Perennial Aquatic	121	5,567	2.2	0 ^c			
<i>Roosting - Temporary</i>							
Cultivated land	14,573	487,106	3.0	0	0		
Grassland	341	78,047	0.4	0 ^c	0		
Managed Wetland	1,008	70,798	1.4	0	0		
Nontidal Freshwater Perennial Emergent Wetland	73	1,509	8.3	0 ^d	0		
Nontidal Perennial Aquatic	191	5,567	8.3	0 ^d	0		
<i>Foraging</i>							
Cultivated land	135,413	487,106	27.8	0	0		
Alkali Seasonal Wetland Complex	22	3,723	0.6	0 ^c	0		
Grassland	21,032	78,047	26.9	0 ^c	0		
Managed Wetland	3,713	70,798	5.2	0 ^e	0		
Nontidal Perennial Aquatic	0	5,567	0.0	400	0		
Other Natural Seasonal Wetland	184	842	21.9	0	0		
Vernal Pool Complex	1,799	12,132	14.8	0 ^c	0		
Least Bell's vireo							
<i>Nesting and Migratory Habitat</i>							
Valley/Foothill Riparian	14,206	17,966	79.1	5,000	3,954		1,000 ^f
Suisun song sparrow							
<i>Primary Habitat</i>							
Tidal Brackish Emergent Wetland	3,221	8,501	37.9	1,500	568	1,500	
Tidal Freshwater Emergent Wetland	339	8,856	3.8	0 ^g	0		
<i>Secondary Habitat</i>							
Managed Wetland	18,125	70,798	25.6	0 ^h	0		
Tidal Brackish Emergent Wetland	2,990	8,501	35.2	4,500	1,583	4,500	
Tidal Freshwater Emergent Wetland	2,455	8,856	27.7	0 ^g	0		
Swainson's hawk							
<i>Foraging Habitat</i>							

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Cultivated land	361,365	487,106	74.2	0	0		
Alkali Seasonal Wetland Complex	3,261	3,723	87.6	0 ⁱ	0		
Grassland	71,343	78,047	91.4	2,000	1,828	2,000	
Managed Wetland	22,304	70,798	31.5	0	0		
Other Natural Seasonal Wetland	259	842	30.8	0	0		
Vernal Pool Complex	11,246	12,132	92.7	0 ⁱ	0		
<i>Nesting Habitat</i>							
Valley/Foothill Riparian	9,388	17,966	52.3	5,000	2,613		
Tricolored blackbird							
<i>Breeding Habitat—Ag Foraging</i>							
Cultivated land	100,198	487,106	20.6	0	0		
<i>Breeding Habitat—Foraging</i>							
Alkali Seasonal Wetland Complex	3,463	3,723	93.0	72	67		
Grassland	38,819	78,047	49.7	2,000	995		
Managed Wetland	6,991	70,798	9.9	0 ⁱ	0		
Other Natural Seasonal Wetland	188	842	22.3	0	0		
Tidal Brackish Emergent Wetland	773	8,501	9.1	6,000	546		
Vernal Pool Complex	7,940	12,132	65.4	67	44		
<i>Breeding Habitat—Nesting</i>							
Managed Wetland	57	70,798	0.1	500	0		
Nontidal Freshwater Perennial Emergent Wetland	279	1,509	18.5	800 ^a	148		
Valley/Foothill Riparian	1,405	17,966	7.8	5,000	391		
<i>Nonbreeding Habitat—Foraging Ag</i>							
Cultivated land	194,251	487,106	39.9	0	0		
<i>Nonbreeding Habitat—Roosting</i>							
Managed Wetland	9,889	70,798	14.0	500	70		
Nontidal Freshwater Perennial Emergent Wetland	935	1,509	61.9	800 ^a	496		
Tidal Brackish Emergent Wetland	4,880	8,501	57.4	6,000	3,444		
Tidal Freshwater Emergent Wetland	8,413	8,856	95.0	24,000	22,800		
Valley/Foothill Riparian	3,805	17,966	21.2	5,000	1,059		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
<i>Nonbreeding Habitat—Foraging</i>							
Alkali Seasonal Wetland Complex	122	3,723	3.3	72	2		
Grassland	32,213	78,047	41.3	2,000	825		
Managed Wetland	1,588	70,798	2.2	500	11		
Tidal Brackish Emergent Wetland	145	8,501	1.7	6,000	102		
Vernal Pool Complex	228	12,132	1.9	67	1		
Western burrowing owl							
<i>High-Value Habitat</i>							
Cultivated land	68,761	487,106	14.1	0	0		
Alkali Seasonal Wetland Complex	3,081	3,723	82.8	72	60		
Grassland	59,437	78,047	76.2	2,000	1,523		
Managed Wetland	7,365	70,798	10.4	0	0		
Vernal Pool Complex	10,706	12,132	88.2	67	59		
Other Natural Seasonal Wetland	0	842	0.0	0	0		
<i>Low-Value Habitat</i>							
Cultivated land	235,559	487,106	48.4	0	0		
Alkali Seasonal Wetland Complex	122	3,723	3.3	72	2		
Grassland	28	78,047	0.0	2,000	1		
Managed Wetland	14,567	70,798	20.6	0	0		
Other Natural Seasonal Wetland	242	842	28.7	0	0		
Western Yellow-billed Cuckoo							
<i>Breeding Habitat</i>							
Valley/Foothill Riparian	1,970	17,966	11.0	5,000	548	500	
<i>Migratory Habitat</i>							
Valley/Foothill Riparian	10,409	17,966	57.9	5,000	2,897		
White-tailed kite							
<i>Breeding/Roosting</i>							
Valley/Foothill Riparian	13,655	17,966	76.0	5,000	3,800		
<i>Foraging</i>							
Cultivated land	357,626	487,106	73.4	0	0		
Alkali Seasonal Wetland Complex	3,450	3,723	92.7	72	67		
Grassland	74,961	78,047	96.0	2,000	1,921		
Managed Wetland	50,808	70,798	71.8	0	0		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Other Natural Seasonal Wetland	261	842	31.0	0	0		
Vernal Pool Complex	11,282	12,132	93.0	67	62		
Yellow-breasted chat							
<i>Primary Nesting and Migratory Habitat</i>							
Valley/Foothill Riparian	8,178	17,966	45.5	5,000	2,276	1,000	
<i>Secondary Nesting and Migratory Habitat</i>							
Valley/Foothill Riparian	5,528	17,966	30.8	5,000	1,538		
<i>Suisun Marsh/Upper Yolo Bypass Nest and Migratory Habitat</i>							
Valley/Foothill Riparian	520	17,966	2.9	5,000	145		
Reptiles							
Giant Garter Snake							
<i>Aquatic—Tidal</i>							
Tidal Perennial Aquatic	6,430	86,263	7.5	0	0		
Tidal Freshwater Emergent Wetland	5,667	8,856	64.0	24,000	15,357		1,250 ^k
<i>Aquatic—Nontidal</i>							
Cultivated land	12,337	487,106	2.5	0	0		
Nontidal Freshwater Perennial Emergent Wetland	1,359	1,509	90.0	800	720		733 ^l
Nontidal Perennial Aquatic	5,331	5,567	95.8	400	383		1,467 ^l
<i>Upland—High</i>							
Cultivated land	5,071	487,106	1.0	0	0		
Alkali Seasonal Wetland Complex	644	3,723	17.3	72	12		
Grassland	14,490	78,047	18.6	2,000	371		700 ^m
Managed Wetland	923	70,798	1.3	500 ⁿ	7		
Vernal Pool Complex	454	12,132	3.7	67	3		
<i>Upland—Moderate</i>							
Cultivated land	3,406	487,106	0.7	0	0		
Alkali Seasonal Wetland Complex	230	3,723	6.2	72	4		
Grassland	8,375	78,047	10.7	2,000	215		
Managed Wetland	5,113	70,798	7.2	500	36		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Vernal Pool Complex	609	12,132	5.0	67	3		
<i>Upland—Low</i>							
Managed Wetland	31	70,798	0.0	500	0		
Vernal Pool Complex	1	12,132	0.0	67	0		
Western pond turtle							
<i>Aquatic Habitat</i>							
Cultivated land	15	487,106	0.0	0	0		
Grassland	0	78,047	0.0	2,000	0		
Managed Wetland	10,820	70,798	15.3	500	76		
Nontidal Freshwater Perennial Emergent Wetland	864	1,509	57.3	800	458	1,200	
Nontidal Perennial Aquatic	5,489	5,567	98.6	400	394		
Tidal Brackish Emergent Wetland	5,768	8,501	67.9	6,000	4,071		
Tidal Freshwater Emergent Wetland	8,855	8,856	100.0	24,000	23,997		
Tidal Perennial Aquatic	49,759	86,263	57.7	0	0		
<i>Upland Nesting and Overwintering Habitat</i>							
Cultivated land	150	487,106	0.0	0	0		
Grassland	13,983	78,047	17.9	2,000	358		
Managed Wetland	1,159	70,798	1.6	500	8		
Tidal Brackish Emergent Wetland	160	8,501	1.9	6,000	113		
Tidal Perennial Aquatic	1	86,263	0.0	0	0		
Valley/Foothill Riparian	2	17,966	0.0	6,000	1		
<i>Upland Nesting and Overwintering Habitat—NHD</i>							
Cultivated land	114	487,106	0.0	0	0°		
Grassland	31,186	78,047	40.0	2,000	799°		
Managed Wetland	2,923	70,798	4.1	500	21°		
Tidal Brackish Emergent Wetland	141	8,501	1.7	6,000	100°		
Valley/Foothill Riparian	74	17,966	0.4	5,000	21°		
Amphibians							
California red-legged frog							
<i>Aquatic Habitat</i>							
Managed Wetland	23	70,798	0.0	0°	0		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Nontidal Freshwater Perennial Emergent Wetland	34	1,509	2.3	0 ^p	0		
Nontidal Perennial Aquatic	84	5,567	1.5	0 ^p			
Tidal Freshwater Emergent Wetland	6	8,856	0.1	24,000	16		
<i>Upland Cover and Dispersal Habitat</i>							
Grassland	6,729	78,047	8.6	2,000	172		
Valley/Foothill Riparian	636	17,966	3.5	5,000	177		
Vernal Pool Complex	402	12,132	3.3	67	2		
California tiger salamander							
<i>Aquatic Breeding Habitat</i>							
Vernal Pool Complex	7,845	12,132	64.7	67	43		
<i>Terrestrial Cover and Aestivation</i>							
Alkali Seasonal Wetland Complex	2,352	3,723	63.2	72	45		
Grassland	23,342	78,047	29.9	2,000	598		
Invertebrates							
Valley elderberry longhorn beetle							
<i>Riparian Vegetation</i>							
Valley/Foothill Riparian	17,451	17,966	97.1	5,000	4,857		
<i>Non-Riparian Channels and Grasslands</i>							
Grassland	15,943	78,047	20.4	0 ^r	0		
California linderiella							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
Vernal Pool Complex	8,571	12,132	70.6	67	47		
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Conservancy fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
Vernal Pool Complex	8,571	12,132	70.6	67	47		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Longhorn fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
Vernal Pool Complex	8,571	12,132	70.6	67	47		
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Midvalley fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
Vernal Pool Complex	8,571	12,132	70.6	67	47		
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Vernal pool fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
Vernal Pool Complex	8,571	12,132	70.6	67	47		
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Vernal pool tadpole shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
Vernal Pool Complex	8,571	12,132	70.6	67	47		
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		

BGOs = Biological Goals and Objectives

- a The 1,200-acre nontidal emergent wetland restoration under GGS1.1 assumes 2/3 nontidal perennial aquatic and 1/3 nontidal emergent wetland.
- b Freshwater emergent wetland restoration under Objective TFEW1.1 is not likely to overlap with locations that benefit the California clapper rail.
- c Grassland, vernal pool, and alkali seasonal wetland protection likely to occur outside the range of greater sandhill crane.
- d The 1,200-acres of nontidal emergent wetland restoration under GGS1.1 do not overlap with the range of the greater sandhill crane.
- e All 500 acres of managed wetland restoration under Objective GSHC1.4 will benefit roosting habitat.

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
f	1,000 acres of early- to mid-successional riparian maintained under Objective VFRN2.2 assumed to benefit least Bell's vireo.						
g	24,000 acres of tidal freshwater emergent wetland restored under Objective TFEW1.1 are outside the range of the Suisun song sparrow.						
h	500 acres of restored managed wetland under Objective GSHC1.4 are outside the range of the Suisun song sparrow.						
i	Assuming no benefit from vernal pool or alkali season wetland natural community restoration because Objective VPNC 1.2 and ASWNC1.2 commit to no net loss of habitat						
j	500 acres of restored managed wetland under Objective GSHC1.4 are not likely to benefit tricolored blackbird.						
k	Objectives GGS1.4 and GGS2.3 provide for the conservation of 4,240 acres of rice or equivalent; assume 1,250 acres of muted tidal restoration (part of 65,00-acre commitment), 1,000 acres as nontidal restoration, 1,000 acres of rice protection, and 1,000 acres as upland protection.						
l	Objectives GGS1.4 and GGS2.3 provide for the conservation of 4,240 acres of rice or equivalent; assume 1,250 acres of muted tidal restoration (part of 65,00-acre commitment), 1,000 acres of nontidal wetland restoration, 1,000 acres of rice protection, and 1,000 acres as upland protection. In addition, 1,200 acres of nontidal marsh will be restored under Objective GGS1.1. This is a total of 2,200 acres of nontidal restoration, 1/3 of which is assumed to be nontidal emergent wetland and 2/3 of which is assumed to be nontidal perennial aquatic.						
m	Of the 400 acres of grassland created or protected under Objectives GGS1.2 and 2.3, assume 200 acres protected and 200 acres restored. Additionally, for the 1,000 acres of grassland protected or created as "rice or equivalent" under Objectives GGS1.4 and GGS3.1 assume 500 acres are protected and 500 are restored.						
n	A portion of the managed wetlands restored for greater sandhill crane under Objective GSHC1.4 could potentially support GGS.						
o	35% of total benefit calculated here will be carried forward to Table 5.6-7 <i>Net Effects, Wildlife</i> , see Appendix 2A.29 <i>Western Pond Turtle Species Account</i> for details.						
p	500 acres of restored managed wetland under Objective GSHC1.4 are outside the range of the California red-legged frog.						
q	The 1,200-acres of nontidal emergent wetland restoration under GGS1.1 do not overlap with the range of the California red-legged frog.						
r	Grassland restoration under Objective GNC1.1 will not contribute habitat for valley elderberry longhorn beetle.						

1
2

1 Table 5J.B-2. Natural Community Protection Contributing to Covered Species Conservation—Wildlife

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
Mammals							
Riparian brush rabbit							
<i>Riparian Habitat</i>							
Valley/Foothill Riparian	2,909	17,966	16.2	750	121		200
<i>Grassland Habitat</i>							
Grassland	3,094	78,047	4.0	8,000	317		
Riparian woodrat							
<i>Habitat</i>							
Valley/Foothill Riparian	2,166	17,966	12.1	750	90		
Salt marsh harvest mouse							
<i>Managed Wetland—Upland Low, Long-Term Conservation Value</i>							
Managed Wetland	3,787	70,798	5.3	0 ^a	0		
<i>Managed Wetland—Wetland Primary Low, Long-Term Conservation Value</i>							
Managed Wetland	21,891	70,798	30.9	1,500	464	1,500	
<i>Managed Wetland—Wetland Secondary Low, Long-Term Conservation Value</i>							
Managed Wetland	2,800	70,798	4.0	0 ^a	0		
<i>Tidal Brackish Emergent Wetland Primary</i>							
Tidal Brackish Emergent Wetland	3,342	8,501	39.3		0		
<i>Tidal Brackish Emergent Wetland Secondary</i>							
Tidal Brackish Emergent Wetland	2,718	8,501	32.0	0	0		
<i>Upland Secondary</i>							
Grassland	491	78,047	0.6	8,000	50		
Tidal Brackish Emergent Wetland	189	8,501	2.2	0	0		
San Joaquin kit fox							
<i>Breeding, Foraging, and Dispersal Habitat</i>							
Grassland	5,098	78,047	6.5	8,000	523	1,000	
Vernal Pool Complex	229	12,132	1.9	600	11		
Suisun shrew							
<i>Primary Habitat</i>							
Grassland	12	78,047	0.0	8,000	1		
Tidal Brackish Emergent Wetland	3,001	8,501	35.3	0	0		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
<i>Secondary Habitat</i>							
Grassland	219	78,047	0.3	8,000	22		
Managed Wetland	1,825	70,798	2.6	8,100	209		
Tidal Brackish Emergent Wetland	2,181	8,501	25.7	0	0		
Birds							
California black rail							
<i>Primary Habitat</i>							
Nontidal Freshwater Perennial Emergent Wetland	715	1,509	47.4	0	0		
Tidal Brackish Emergent Wetland	3,760	8,501	44.2	0	0		
Tidal Freshwater Emergent Wetland	1,458	8,856	16.5	0	0		
<i>Secondary Habitat</i>							
Managed Wetland	12,957	70,798	18.3	1,500	275		
Nontidal Freshwater Perennial Emergent Wetland	66	1,509	4.4	0	0		
Tidal Brackish Emergent Wetland	2,022	8,501	23.8	0	0		
Tidal Freshwater Emergent Wetland	2,797	8,856	31.6	0	0		
California clapper rail							
<i>Primary Habitat</i>							
Tidal Brackish Emergent Wetland	248	8,501	2.9	0	0		
<i>Secondary Habitat</i>							
Tidal Brackish Emergent Wetland	5,324	8,501	62.6	0	0		
Tidal Freshwater Emergent Wetland	753	8,856	8.5	0	0		
Greater sandhill crane							
<i>Roosting - Permanent</i>							
Cultivated land	5,237	487,106	1.1	0 ^b	0		
Grassland	628	78,047	0.8	0 ^c	0		
Managed Wetland	1,097	70,798	1.5	0	0		
Nontidal Freshwater Perennial Emergent Wetland	43	1,509	2.9	0 ^d	0		
Nontidal Perennial Aquatic	121	5,567	2.2	0 ^d	0		
<i>Roosting - Temporary</i>							
Cultivated land	14,573	487,106	3.0	0 ^b	0		
Grassland	341	78,047	0.4	0 ^c	0		
Managed Wetland	1,008	70,798	1.4	0	0		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
Nontidal Freshwater Perennial Emergent Wetland	73	1,509	4.9	0 ^d	0		
Nontidal Perennial Aquatic	191	5,567	3.4	0 ^d	0		
<i>Foraging</i>							
Cultivated land	135,413	487,106	27.8	48,600	13,511		7,300
Alkali Seasonal Wetland Complex	22	3,723	0.6	0 ^c	0		
Grassland	21,032	78,047	26.9	0 ^c	0		
Managed Wetland	3,713	70,798	5.2	0	0		
Nontidal Perennial Aquatic	0	5,567	0.0	0	0		
Other Natural Seasonal Wetland	184	842	21.9	0	0		
Vernal Pool Complex	1,799	12,132	14.8	0 ^c	0		
Least Bell's vireo							
<i>Nesting and Migratory Habitat</i>							
Valley/Foothill Riparian	14,206	17,966	79.1	750	593		
Suisun song sparrow							
<i>Primary Habitat</i>							
Tidal Brackish Emergent Wetland	3,221	8,501	37.9	0	0		
Tidal Freshwater Emergent Wetland	339	8,856	3.8	0	0		
<i>Secondary Habitat</i>							
Managed Wetland	18,125	70,798	25.6	1,500 ^e	384		
Tidal Brackish Emergent Wetland	2,990	8,501	35.2	0	0		
Tidal Freshwater Emergent Wetland	2,455	8,856	27.7	0	0		
Swainson's hawk							
<i>Foraging Habitat</i>							
Cultivated land	361,365	487,106	74.2	48,600	36,054		43,325
Alkali Seasonal Wetland Complex	3,261	3,723	87.6	150	131		150
Grassland	71,343	78,047	91.4	8,000	7,313		8,000
Managed Wetland	22,304	70,798	31.5	8,100	2,552		
Other Natural Seasonal Wetland	259	842	30.8	0	0		
Vernal Pool Complex	11,246	12,132	92.7	600	556		600
<i>Nesting Habitat</i>							
Valley/Foothill Riparian	9,388	17,966	52.3	750	392		
Tricolored blackbird							
<i>Breeding Habitat—Ag Foraging</i>							
Cultivated land	100,198	487,106	20.6	48,600	9,997		11,050

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
<i>Breeding Habitat—Foraging</i>							
Alkali Seasonal Wetland Complex	3,463	3,723	93.0	150	140		
Grassland	38,819	78,047	49.7	8,000	3,979		
Managed Wetland	6,991	70,798	9.9	8,100	800		
Other Natural Seasonal Wetland	188	842	22.3	0	0		
Tidal Brackish Emergent Wetland	773	8,501	9.1	0	0		
Vernal Pool Complex	7,940	12,132	65.4	600	393		
<i>Breeding Habitat—Nesting</i>							
Managed Wetland	57	70,798	0.1	8,100	7		
Nontidal Freshwater Perennial Emergent Wetland	279	1,509	18.5	0	0		50
Valley/Foothill Riparian	1,405	17,966	7.8	750	59		
<i>Nonbreeding Habitat—Foraging Ag</i>							
Cultivated land	194,251	487,106	39.9	48,600	19,381		26,300
<i>Nonbreeding Habitat—Roosting</i>							
Managed Wetland	9,889	70,798	14.0	8,100	1,131		
Nontidal Freshwater Perennial Emergent Wetland	935	1,509	61.9	0	0		
Tidal Brackish Emergent Wetland	4,880	8,501	57.4	0	0		
Tidal Freshwater Emergent Wetland	8,413	8,856	95.0	0	0		
Valley/Foothill Riparian	3,805	17,966	21.2	750	159		
<i>Nonbreeding Habitat—Foraging</i>							
Alkali Seasonal Wetland Complex	122	3,723	3.3	150	5		
Grassland	32,213	78,047	41.3	8,000	3,302		
Managed Wetland	1,588	70,798	2.2	8,100	182		
Tidal Brackish Emergent Wetland	145	8,501	1.7	0	0		
Vernal Pool Complex	228	12,132	1.9	600	11		
Western burrowing owl							
<i>High-Value Habitat</i>							
Cultivated land	68,761	487,106	14.1	48,600	6,860		1,000
Alkali Seasonal Wetland Complex	3,081	3,723	82.8	150	124		
Grassland	59,437	78,047	76.2	8,000	6,092		
Managed Wetland	7,365	70,798	10.4	8,100	843		
Vernal Pool Complex	10,706	12,132	88.2	600	529		
Other Natural Seasonal Wetland	0	842	0.0	0	0		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
<i>Low-Value Habitat</i>							
Cultivated land	235,559	487,106	48.4	48,600	23,502		
Alkali Seasonal Wetland Complex	122	3,723	3.3	150	5		
Grassland	28	78,047	0.0	8,000	3		
Managed Wetland	14,567	70,798	20.6	8,100	1,667		
Other Natural Seasonal Wetland	242	842	28.7	0	0		
Western Yellow-billed Cuckoo							
<i>Breeding Habitat</i>							
Valley/Foothill Riparian	1,970	17,966	11.0	750	82		
<i>Migratory Habitat</i>							
Valley/Foothill Riparian	10,409	17,966	57.9	750	435		
White-tailed kite							
<i>Breeding/Roosting</i>							
Valley/Foothill Riparian	13,655	17,966	76.0	750	570		
<i>Foraging</i>							
Cultivated land	357,626	487,106	73.4	48,600	35,681		
Alkali Seasonal Wetland Complex	3,450	3,723	92.7	150	139		
Grassland	74,961	78,047	96.0	8,000	7,684		
Managed Wetland	50,808	70,798	71.8	8,100	5,813		
Other Natural Seasonal Wetland	261	842	31.0	0	0		
Vernal Pool Complex	11,282	12,132	93.0	600	558		
Yellow-breasted chat							
<i>Primary Nesting and Migratory Habitat</i>							
Valley/Foothill Riparian	8,178	17,966	45.5	750	341		
<i>Secondary Nesting and Migratory Habitat</i>							
Valley/Foothill Riparian	5,528	17,966	30.8	750	231		
<i>Suisun Marsh/Upper Yolo Bypass Nest and Migratory Habitat</i>							
Valley/Foothill Riparian	520	17,966	2.9	750	22		
Reptiles							
Giant Garter Snake							
<i>Aquatic—Tidal</i>							
Tidal Perennial Aquatic	6,430	86,263	7.5	0	0		
Tidal Freshwater Emergent Wetland	5,667	8,856	64.0	0	0		
<i>Aquatic—Nontidal</i>							
Cultivated land	12,337	487,106	2.5	48,600	1,231		1,500 ^f

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
Nontidal Freshwater Perennial Emergent Wetland	1,359	1,509	90.0	25 ^g	23		
Nontidal Perennial Aquatic	5,331	5,567	95.8	25 ^g	24		
<i>Upland—High</i>							
Cultivated land	5,071	487,106	1.0	48,600	506		200 ^h
Alkali Seasonal Wetland Complex	644	3,723	17.3	150	26		
Grassland	14,490	78,047	18.6	8,000	1,485		700 ⁱ
Managed Wetland	923	70,798	1.3	0 ^j	0		
Vernal Pool Complex	454	12,132	3.7	600	22		
<i>Upland—Moderate</i>							
Cultivated land	3,406	487,106	0.7	48,600	340		
Alkali Seasonal Wetland Complex	230	3,723	6.2	150	9		
Grassland	8,375	78,047	10.7	8,000	858		
Managed Wetland	5,113	70,798	7.2	0 ^j	0		
Vernal Pool Complex	609	12,132	5.0	600	30		
<i>Upland—Low</i>							
Managed Wetland	31	70,798	0.0	0 ^j	0		
Vernal Pool Complex	1	12,132	0.0	600	0		
Western pond turtle							
<i>Aquatic Habitat</i>							
Cultivated land	15	487,106	0.0	48,600	2		
Grassland	0	78,047	0.0	8,000	0		
Managed Wetland	10,820	70,798	15.3	8,100	1,238		
Nontidal Freshwater Perennial Emergent Wetland	864	1,509	57.3	25	14		
Nontidal Perennial Aquatic	5,489	5,567	98.6	25	25		
Tidal Brackish Emergent Wetland	5,768	8,501	67.9	0	0		
Tidal Freshwater Emergent Wetland	8,855	8,856	100.0	0	0		
Tidal Perennial Aquatic	49,759	86,263	57.7	0	0		
<i>Upland Nesting and Overwintering Habitat</i>							
Cultivated land	150	487,106	0.0	48,600	15		
Grassland	13,983	78,047	17.9	8,000	1,433		
Managed Wetland	1,159	70,798	1.6	0	0		
Tidal Brackish Emergent Wetland	160	8,501	1.9	0	0		
Tidal Perennial Aquatic	1	86,263	0.0	0	0		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
Valley/Foothill Riparian	2	17,966	0.0	750	0		
<i>Upland Nesting and Overwintering Habitat—NHD^k</i>							
Cultivated land	114	487,106	0.0	48,600	11 ^k		
Grassland	31,186	78,047	40.0	8,000	3,197 ^k		
Managed Wetland	2,923	70,798	4.1	8,100	334 ^k		
Tidal Brackish Emergent Wetland	141	8,501	1.7	0	0 ^k		
Valley/Foothill Riparian	74	17,966	0.4	750	3 ^k		
Amphibians							
California red-legged frog							
<i>Aquatic Habitat</i>							
Managed Wetland	23	70,798	0.0	8,100	3		
Nontidal Freshwater Perennial Emergent Wetland	34	1,509	2.3	25 ^g	0		
Nontidal Perennial Aquatic	84	5,567	1.5	25 ^g			
Tidal Freshwater Emergent Wetland	6	8,856	0.1	0	0		
<i>Upland Cover and Dispersal Habitat</i>							
Grassland	6,729	78,047	8.6	8,000	690	1000 ^l	
Valley/Foothill Riparian	636	17,966	3.5	750	27		
Vernal Pool Complex	402	12,132	3.3	600	20		
California tiger salamander							
<i>Aquatic Breeding Habitat</i>							
Vernal Pool Complex	7,845	12,132	64.7	600	388	600	
<i>Terrestrial Cover and Aestivation*</i>							
Alkali Seasonal Wetland Complex	2,352	3,723	63.2	150	95	150	
Grassland	23,342	78,047	29.9	8,000	2,393	5,000 ^m	
Invertebrates							
Valley elderberry longhorn beetle							
<i>Riparian Vegetation</i>							
Valley/Foothill Riparian	17,451	17,966	97.1	750	729		
<i>Non-Riparian Channels and Grasslands</i>							
Grassland	15,943	78,047	20.4	8,000	1,634		
California linderiella							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
Vernal Pool Complex	8,571	12,132	70.6	600	424	600	

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Conservancy fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
Vernal Pool Complex	8,571	12,132	70.6	600	424	600	
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Longhorn fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
Vernal Pool Complex	8,571	12,132	70.6	600	424	600	
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Midvalley fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
Vernal Pool Complex	8,571	12,132	70.6	600	424	600	
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Vernal pool fairy shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
Vernal Pool Complex	8,571	12,132	70.6	600	424	600 ^e	
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		
Vernal pool tadpole shrimp							
<i>High Quality Habitat</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
Vernal Pool Complex	8,571	12,132	70.6	600	424	600 ^e	
<i>Low Quality Habitat</i>							
Vernal Pool Complex	2,713	12,132	22.4	0	0		

BGOs = Biological Goals and Objectives

^a Managed wetland benefit attributed to Managed Wetland—Wetland primary Low, Long-Term Conservation Value

^b All cultivated land natural community protection benefit applied to foraging model type.

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
<p>^c Grassland, vernal pool, and alkali seasonal wetland protection likely to occur outside the range of greater sandhill crane.</p> <p>^d Nontidal freshwater emergent wetland and aquatic habitat protected for tricolored blackbird unlikely to overlap with greater sandhill crane range.</p> <p>^e Given uncertainty of benefits on managed wetlands managed for waterfowl and shorebird foraging, nesting, and brooding, only 1,500 acres of managed wetland protection to benefit the salt marsh harvest mouse is applied.</p> <p>^f Objectives GGS1.4 and GGS2.3 provide for the conservation of 4,240 acres of rice or equivalent; assume 1,250 acres of muted tidal restoration (part of 65,00-acre commitment), 1,000 acres as nontidal restoration, 1,000 acres of rice protection, and 1,000 acres as upland protection. Additionally, 500 of 700 acres of cultivated land protection required under Objective GGS2.3 assumed to be rice.</p> <p>^g 25 acres of emergent wetland and 25 acres nontidal perennial aquatic assumed for 50-acre tricolored blackbird nontidal emergent wetland protection commitment under Objective TRBL1.1.</p> <p>^h 200 acres of the 700-acre commitment under Objective GGS2.3 assumed to be a non-rice crop type.</p> <p>ⁱ Of the 400 acres of grassland created or protected under Objectives GGS1.2 and 2.3, assume 200 acres protected and 200 acres restored. Additionally, for the 1,000 acres of grassland protected or created as "rice or equivalent" under Objectives GGS1.4 and GGS3.1 assume 500 acres are protected and 500 are restored.</p> <p>^j Managed wetland protection will be in Suisun Marsh, outside the range of the giant garter snake.</p> <p>^k 35% of total benefit calculated here will be carried forward to Table 5.6-7 <i>Net Effects, Wildlife</i>, see Appendix 2A.29 <i>Western Pond Turtle Species Account</i> for details.</p> <p>^l Under Objective GNC1.1, minimum grassland protection commitment in CZ8 where California red-legged frog habitat overlaps 100% with areas targeted for grassland protection.</p> <p>^m Under Objective GNC1.1, total minimum commitments for CZs 1, 8, and 11 where grassland protection overlaps with California tiger salamander habitat.</p>							

1
2

1 Table 5J.B-3. Natural Community Restoration Contributing to Covered Species Conservation—Plants

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Plants							
Brittlescale							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	23	3,723	0.6	72	0		
Grassland	174	78,047	0.2	2,000	4		
Vernal Pool Complex	182	12,132	1.5	67	1		
Heartscale							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	541	3,723	14.5	72	10		
Grassland	3,189	78,047	4.1	2,000	82		
Vernal Pool Complex	2,721	12,132	22.4	67	15		
San Joaquin spearscale							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	2,561	3,723	68.8	72	50		
Grassland	7,126	78,047	9.1	2,000	183		
Vernal Pool Complex	4,790	12,132	39.5	67	26		
Carquinez goldenbush							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	19	3,723	0.5	72	0		
Grassland	536	78,047	0.7	2,000	14		
Vernal Pool Complex	616	12,132	5.1	67	3		
Delta button celery							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	94	3,723	2.5	72	2		
Grassland	1,547	78,047	2.0	2,000	40		
Valley/Foothill Riparian	768	17,966	4.3	5,000	214		
Vernal Pool Complex	370	12,132	3.1	67	2		
Delta mudwort							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	364	8,501	4.3	6,000	257		
Tidal Freshwater Emergent	762	8,856	8.6	24,000	2,065		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
Wetland							
Valley/Foothill Riparian	954	17,966	5.3	5,000	265		
Mason's lilaeopsis							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	364	8,501	4.3	6,000	257		
Tidal Freshwater Emergent Wetland	762	8,856	8.6	24,000	2,065		
Valley/Foothill Riparian	954	17,966	5.3	5,000	265		
Delta tule pea							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	5,185	8,501	61.0	6,000	3,659		
Valley/Foothill Riparian	477	17,966	2.7	5,000	133		
Suisun marsh aster							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	5,185	8,501	61.0	6,000	3,659		
Valley/Foothill Riparian	477	17,966	2.7	5,000	133		
Side-flowering skullcap							
<i>Habitat</i>							
Valley/Foothill Riparian	2,497	17,966	13.9	5,000	695		
Slough thistle							
<i>Habitat</i>							
Valley/Foothill Riparian	768	17,966	4.3	5,000	214		
Soft bird's-beak							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	1,129	8,501	13.3	1,500	199	1,500	
Suisun thistle							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	1,281	8,501	15.1	1,500	226	1,500	
Vernal Pool Plants							
Alkali milk-vetch							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	67	14		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Restoration	Estimated Contribution of Natural Community Restoration to Species Habitat Restoration	Minimum Restoration Commitment from Natural Community BGOs	Minimum Restoration Commitment from Species-Specific BGOs
<i>Vernal Pool Complex</i>		3					
Vernal Pool Complex	8,709	12,132	71.8	67	48		
Legenere							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	67	14		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	67	48		
Heckard's peppergrass							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	67	14		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	67	48		
Boggs lake hedge-hyssop							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	67	14		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	67	48		
Dwarf downingia							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	72	4		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	67	14		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	67	48		

BGOs = Biological Goals and Objectives.

1 Table 5J.B-4. Natural Community Protection Contributing to Covered Species Conservation—Plants

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
Plants							
Brittlescale^a							
<i>Habitat</i>							75
Alkali Seasonal Wetland Complex	23	3,723	0.6	150	1		
Grassland	174	78,047	0.2	8,000	18		
Vernal Pool Complex	182	12,132	1.5	600	9		
Heartscale^a							
<i>Habitat</i>							75
Alkali Seasonal Wetland Complex	541	3,723	14.5	150	22		
Grassland	3,189	78,047	4.1	8,000	327		
Vernal Pool Complex	2,721	12,132	22.4	600	135		
San Joaquin spearscale							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	2,561	3,723	68.8	150	103		
Grassland	7,126	78,047	9.1	8,000	730		
Vernal Pool Complex	4,790	12,132	39.5	600	237		
Carquinez goldenbush							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	19	3,723	0.5	150	1		
Grassland	536	78,047	0.7	8,000	55		
Vernal Pool Complex	616	12,132	5.1	600	30		
Delta button celery							
<i>Habitat</i>							
Alkali Seasonal Wetland Complex	94	3,723	2.5	150	4		
Grassland	1,547	78,047	2.0	8,000	159		
Valley/Foothill Riparian	768	17,966	4.3	750	32		
Vernal Pool Complex	370	12,132	3.1	600	18		
Delta mudwort							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	364	8,501	4.3	0	0		
Tidal Freshwater Emergent Wetland	762	8,856	8.6	0	0		
Valley/Foothill Riparian	954	17,966	5.3	0 ^a	0		

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
Mason's lilaeopsis							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	364	8,501	4.3	0	0		
Tidal Freshwater Emergent Wetland	762	8,856	8.6	0	0		
Valley/Foothill Riparian	954	17,966	5.3	0 ^a	0		
Delta tule pea							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	5,185	8,501	61.0	0	0		
Valley/Foothill Riparian	477	17,966	2.7	0 ^a	0		
Suisun marsh aster							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	5,185	8,501	61.0	0	0		
Valley/Foothill Riparian	477	17,966	2.7	0 ^a	0		
Side-flowering skullcap							
<i>Habitat</i>							
Valley/Foothill Riparian	2,497	17,966	13.9	0 ^a	0		
Slough thistle							
<i>Habitat</i>							
Valley/Foothill Riparian	768	17,966	4.3	750	32	750	
Soft bird's-beak							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	1,129	8,501	13.3	0	0		
Suisun thistle							
<i>Habitat</i>							
Tidal Brackish Emergent Wetland	1,281	8,501	15.1	0	0		
Vernal Pool Plants							
Alkali milk-vetch							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	0	0		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	600	431	600	

Resource	Acres of Modeled Habitat Comprising the Natural Community	Total Acres of Natural Community in the Plan Area	Percentage of Modeled Habitat Comprising the Natural Community (%)	Total Acres of Natural Community Protection	Estimated Contribution of Natural Community Protection to Species Habitat Protection	Minimum Protection Commitment from Natural Community BGOs	Minimum Protection Commitment from Species-Specific BGOs
Legenere							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	0	0		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	600	431	600	
Heckard's peppergrass							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	0	0		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	600	431	600	
Boggs lake hedge-hyssop							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	0	0		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	600	431	600	
Dwarf downingia							
<i>Alkali Seasonal Wetland</i>							
Alkali Seasonal Wetland Complex	188	3,723	5.0	150	8		
<i>Degraded Vernal Pool Complex</i>							
Vernal Pool Complex	2,576	12,132	21.2	0	0		
<i>Vernal Pool Complex</i>							
Vernal Pool Complex	8,709	12,132	71.8	600	431	600	

BGOs = Biological Goals and Objectives.

^a Riparian protection under Objective 2.4 unlikely to overlap with the range of Delta mudwort, Mason's lilaeopsis, Delta tule pea, Suisun Marsh aster, or side-flowering skullcap.

1
2
3

Attachment 5J.C
**Analysis of Potential Bird Collisions at
Proposed BDCP Powerlines**

Date:	September 3, 2013
To:	Laura King Moon, Project Manager, BDCP California Department of Water Resources
Cc:	
From:	Paola Bernazzani Senior Conservation Biologist, ICF International Gary L. Ivey Research Associate, International Crane Foundation
Subject:	Analysis of Potential Bird Collisions at Proposed BDCP Powerlines

1

2

3

4

5

This memo describes the potential risk to avian species from collision with electrical powerlines that would be installed as part of the Bay Delta Conservation Plan (BDCP) and provides additional analysis of risk and mitigation for the greater sandhill crane (*Grus canadensis tabida*). The following specific factors are addressed.

6

- Assessment of vulnerability for covered birds.

7

- Mortality estimates and population-level effects for greater sandhill crane.

8

- Minimization and mitigation measures for greater sandhill crane based on anticipated levels of take.

9

10 1.0 Introduction

11 1.1 Definitions

12

13

14

Powerlines are rated and categorized by the voltage carried and the purpose served (Avian Power Line Interaction Committee 2006). Because voltages carried by powerlines are typically large, voltage is specified by the kilovolt (kV).

15

16

17

18

- **Distribution lines:** Electrical lines that are energized at lower voltages (60 kV or below). Up to 3.3 miles of temporary, 34.5-kV distribution lines would be installed under the BDCP; additional distribution lines could be used for mitigation. Typically, distribution lines range in height from 35 to 40 feet (11 to 12 meters) (Figure 1) (Avian Power Line Interaction Committee 2006).

19


20

21

- **Transmission lines:** Electrical lines that are energized at higher voltages (60 kV or above). Under the BDCP, 69-kV and 230-kV transmission lines would be installed. Typically, the higher-voltage (230-kV) lines vary in height from 90 to 110 feet (27 to 34 meters), while the “sub”

1 transmission (69-kV) lines vary from 50 to 70 feet (15 to 21 meters). (Figure 1) (Avian Power
2 Line Interaction Committee 2006).

- 3
- 4 • **Ground wire:** An overhead static wire that is installed for protection from lightning (Avian
5 Power Line Interaction Committee 2006).
 - 6 • **Powerlines:** Electrical lines that include both distribution and transmission lines. Overhead
7 powerlines are often equipped with a ground wire. For purposes of assessing risk from covered
activities in the discussion below, ground wires are included as “powerlines.”



8
9

Figure 1. Typical Powerline Structures and Heights

10 1.2 Background

11 Implementation of the BDCP would require installation of powerlines to provide temporary power
12 for construction of new tunnels and pumping facilities. Permanent power is also needed to operate
13 three new intakes on the Sacramento River. Risks to birds from powerlines are described in this
14 memo.

15 Millions of birds are thought to be injured or killed by powerline interactions each year (Erickson et
16 al. 2005; Hunting 2002a). Two main sources of powerline mortality are collision and electrocution.
17 Electrocution occurs when a bird, usually one with a large wingspan, touches two conductors of

1 different phases or a conductor and a ground at the same time (Avian Power Line Interaction
2 Committee 2006). This typically happens when a bird attempts to perch on a structure with
3 insufficient clearance between these elements, often on distribution lines with voltages less than
4 60 kV (Avian Power Line Interaction Committee 2006). Because the majority of the lines that BDCP
5 is constructing are higher-voltage transmission lines (no permanent lines below 69 kV are
6 proposed) and because adequate clearance would be provided between conductors or between
7 conductors and ground wires (e.g., 60 inches [1.5 meters] of horizontal separation and 40 inches [1
8 meter] of vertical separation), electrocution is anticipated to be a negligible source of mortality and
9 therefore is not analyzed further here. Covers on phases or grounds will be installed where
10 adequate separation is not feasible (Avian Power Line Interaction Committee 2006).

11 Bird mortality is also caused by direct collision with powerlines that can be difficult for birds to see,
12 particularly in bad weather. Collision mortality is commonly associated with ground wires, which
13 are found above transmission lines and are thinner and less visible. Ground wires would be
14 installed, under the BDCP, and risks associated with ground wires are included as part of the risk
15 analysis described below. Over 80% of collision fatalities at transmission lines occur through
16 collision with the ground wire (James and Haak 1979; Hunting 2002b). Collision risk at powerlines
17 can be exacerbated by factors that are biological (e.g., age and sex of birds), physical (e.g.,
18 topography), meteorological (e.g., winds, fog), and structural (e.g., line location and design) (Avian
19 Power Line Interaction Committee 1994; Bevanger 1994). Cranes, bustards, flamingos, waterfowls,
20 shorebirds, game birds, and some falcons are the bird groups most frequently affected by
21 transmission line collisions (Jenkins et al. 2010).

22 Despite the fact that several studies have established a strong correlation between powerlines,
23 including ground wires, and collisions risk (e.g., Avian Power Line Interaction Committee 1994;
24 Bevanger 1994, 1998; Janss and Ferrer 2000; Erickson et al. 2005), few estimates of collision
25 mortality exist, and most are based on extrapolations from individual or small-scale studies. A
26 quantitative estimate of powerline mortality requires dedicated surveys (Bevanger 1998), which are
27 time-consuming and costly to undertake. Absent specific information on the mortality rates of
28 covered bird species at transmission lines, this memorandum provides a qualitative discussion of
29 the relative vulnerability of each covered bird species to assess the potential for significant effects
30 from transmission line strikes. Subsequently, this memorandum provides a species-specific risk
31 assessment for greater sandhill crane, the species identified by the vulnerability analysis as at high
32 risk from collision mortality. Powerline collision is thought to be an influential factor in ongoing
33 population declines in several species of cranes (Jenkins et al. 2010), which have large body size, fast
34 flight, flocking behavior, long appendages, and low maneuverability—all risk factors for powerline
35 collision (Bevanger 1998; Hunting 2002b). This memorandum provides a collision risk map,
36 mortality assessment for individuals and populations of cranes, and a mitigation strategy.

37 **1.3 Location and Extent of Facilities**

38 Additional powerlines would provide permanent electric power for new intakes, pumping plants,
39 operable barriers, and gate control structures constructed as part of the BDCP. Also, temporary
40 powerlines would provide power during construction of water conveyance facilities. All proposed

1 permanent lines within the Plan Area are transmission lines (230- and 69-kV and associated ground
2 wires). Temporary lines are both transmission (230-kV) and distribution (34.5-kV).

3 Under the proposed powerline alignment, power would be delivered to the water conveyance
4 facilities using a “split” system that connects to the existing grid at two different locations. The
5 northern point of interconnection would be located north of Lambert Road and west of Highway 99.
6 From that location, a 230-kV transmission line would run west along Lambert Road for
7 approximately 5 miles, at which point one segment would run south to the intermediate forebay on
8 Glannville Tract and then on to tunnel shaft locations on Staten Island. Those segments extending
9 south of the intermediate forebay on McCormack-Williamson Tract and Staten Island are temporary
10 and would be removed following construction of associated tunnel facilities. The other segment
11 would run north to a substation, where permanent, 69-kV lines would connect to the intake
12 pumping plants.

13 In the south, the interconnection would be either southeast of Brentwood near Brentwood
14 Boulevard or adjacent to the Jones Pumping Plant. A 230-kV line would stretch from one of these
15 locations to a tunnel shaft northwest of Clifton Court Forebay and continue north following tunnel
16 shaft locations to Bouldin Island, where a 34.5-kV line would continue to the southern end of Staten
17 Island. All of the power lines extending from the southern point of interconnection would be
18 temporary, limited to the construction schedule for the relevant tunnel reaches and features
19 associated with Clifton Court Forebay.

20 The proposed alignment requires the installation of approximately 20 miles (32 kilometers) of
21 permanent transmission line (14 miles [23 kilometers] of 230-kV lines and 6 miles [10 kilometers]
22 of 69-kV lines) (Table 1).

23 **Table 1. Proposed Powerlines in the Plan Area**

Powerlines	Voltage (kV)	Length (Miles)
Permanent	230	14
	69	6
Total Permanent		20
Temporary	230	35
	34.5	3
Total Temporary		38
Total		58

24
25 The length of temporary lines is approximately 38 miles (61 kilometers) (3 miles [56 kilometers] of
26 34.5-kV line and 35 miles [5 kilometers] of 230-kV line). Temporary lines will be removed after
27 construction of the water conveyance facilities, within 10 years.

2.0 Vulnerability Analysis

Covered bird species were individually assessed to determine the relative risk of collision with the proposed BDCP powerlines and to evaluate whether this risk should be further analyzed, quantified, and mitigated for in the BDCP. The BDCP covers 12 bird species (information on the habitat and distribution of these species in the Plan Area is summarized in Exhibit 1). To assess the risk of powerline collision for each species, a brief analysis of physiological and behavior characteristics is provided. This information is synthesized and, using best professional judgment, a recommendation is made regarding the need for additional analyses.

As discussed above, many factors contribute to the risk of bird collisions with powerlines, including characteristics of the facility. However, all non-biological factors being equal, the relative vulnerability of a bird species to collision mortality depends primarily on its level of exposure (or proximity of the bird's habitat and resources to the powerline) and its sensitivity (morphological and behavioral characteristics that influence the bird's propensity to collide with a line).

For all species, exposure was determined by overlaying occurrences and modeled habitat with the proposed powerline alignment (Exhibit 2, Figures 2-1 through 2-12), using geographic information systems (GIS) (California Department of Fish and Game 2010). Results indicate that 8 of the 11 covered species have been observed within 6 kilometers of the proposed alignment. Species that were at farther distances were the California clapper rail, Suisun song sparrow, and least Bell's vireo. However, all species are discussed below, because covered birds may become more abundant in the Plan Area as the result of enhancement activities, and occurrence data may not accurately reflect species presence.

In addition, factors such as maneuverability, flight altitude, flight times, foraging, flocking, eyesight, and migration behavior were considered, to the extent that this information was available for each species.

Maneuverability

A bird's maneuverability is influenced by wing morphology and size. Maneuverability is one of the most important factors influencing the risk of powerline collision, because it determines a bird's ability to negotiate an obstacle while in flight (Rayner 1988; Bevanger 1994, 1998; Savereno et al. 1996). Different wing shapes correspond to different tradeoffs between speed, energy use, and agility (Bevanger 1998). Wing shape can be described in terms of wing loading—the ratio of bird weight to wing area (a small bird with large wings has low wing loading, while a large bird with small wings has high loading) and wing aspect ratio—the ratio of wing length to wing breadth.

The particular combination of wing loading and aspect ratio determines the type of flight that is possible, as discussed in detail in Rayner (1988) and Bevanger (1998). In general, birds with *low wing loading* and *high aspect-ratio wings* can maneuver relatively quickly around an obstacle. These wings allow rapid flight and quick, evasive actions. Birds with a *high wing loading* and *low aspect-ratio wings* have limited maneuverability and are therefore more susceptible to collision. Body size, in combination with wing morphology, influences a bird's maneuverability, with larger body sizes corresponding to reduced maneuverability, especially in species with relatively small wings.

1 Rails, followed by cranes, display the greatest vulnerability based on wing-shape morphology, with
2 low-aspect and high- or moderate-loading wings, respectively. The remaining covered bird species
3 show a range of low (owls, hawks), medium (cuckoo), and high (kites, falcons, terns, some
4 passerines) wing aspect ratios, but all have relatively low loading, which decreases their general risk
5 of collision (Bevanger 1998). Maneuverability is discussed for each species.

6 **Flight Altitude**

7 Collision risk associated with flight altitude depends on the heights of the lines and ground wires
8 and the flight behavior of a given species. (Meyer 1978; James and Haak 1979; Beaulaurier 1981). As
9 mentioned above, the powerlines that will be installed by the BDCP range from 50 to 110 feet.

10 For discussion purposes, the risk of collision is higher if birds commute to foraging areas within the
11 range of the anticipated height of BDCP distribution and transmission lines. Migration altitudes are
12 typically higher than 110 feet (33.5 meters) as noted in the descriptions below.

13 **Flight Times**

14 Species that are active at dawn or dusk and nocturnally active species are more susceptible to
15 collision because of low light conditions and reduced visibility (Bevanger 1994; Crowder and
16 Rhodes 2001). In addition, in the Central Valley, the collision risk is elevated for overwintering birds
17 because visibility is greatly reduced during the frequent dense fog and rains that occur in winter.

18 **Foraging**

19 Collisions are more likely where powerlines transect or parallel areas used for foraging (Scott et al.
20 1972; Brown et al. 1987; Morkill and Anderson 1991; Brown and Drewien 1995; Murphy et al.
21 2009).

22 **Flocking**

23 Birds in large flocks have less maneuverability and visibility when at the back of the flock
24 (Scott et al. 1972). Daily flock movements between feeding, breeding, and roosting areas place
25 flocking species at high risk of collision compared to species that do not flock (Avian Power Line
26 Interaction Committee 1994).

27 **Vision**

28 Raptors, and other birds of prey, have excellent eyesight and tend not to fly in low-visibility
29 conditions, making them less likely to collide with powerlines (Olendorff and Lehman 1986). Vision
30 is discussed as it pertains to reducing collision risk for relevant species below.

31 **Migration**

32 During migration, birds may collide with overhead wires; however, collisions are more likely
33 associated with taller structures such as communications towers or smoke stacks (Avian Power Line
34 Interaction Committee 1994). Nocturnal migration is the most common contributing factors to these
35 collisions (Avian Power Line Interaction Committee 1994). In general, daytime migrations do not

1 create major collision risk with overhead wires for birds. Except for landing and taking off, most
2 migrants fly well above powerlines (Avian Power Line Interaction Committee 1994). Rather, most
3 powerline collisions occur during flights in daily use areas associated with commuting or foraging
4 (Avian Power Line Interaction Committee 1994). However, during migration, migratory species
5 cross numerous powerlines on the way to and from their wintering grounds and, in general, may be
6 expected to experience greater risk of collisions than resident species (Bevanger 1994).

7 **2.1 California Black Rail**

8 The California black rail is found in the Plan Area year-round. Unlike other subspecies of black rail,
9 the California black rail is largely sedentary and is either nonmigratory or only locally migratory
10 (Eddleman et al. 1994). Migratory and juvenile dispersal movements tend to be localized (Trulio and
11 Evens 2000) with seasonal migratory and dispersal movement occurring within the breeding range
12 of the species. For example, black rails that nest in the north San Francisco Bay area have been
13 reported to winter in the south San Francisco Bay area (Trulio and Evens 2000).

14 In the Bay-Delta region, California black rail populations are restricted primarily to the remaining
15 tidal marshlands of the northern San Francisco Bay estuary, the vicinity of Suisun and Napa
16 Marshes, and the midchannel islands in the Delta. In Suisun Marsh, a high abundance of black rails
17 has been found at east Mallard Island, with moderate abundances at South Joice Island, Pacheco
18 Creek, East Peyton Slough, Cutoff Island, Peytonia Slough, and Southampton Bay. Spautz et al. (2005)
19 estimate a population of 12,000 black rails in the Suisun Bay region. Surveys conducted by the
20 California Department of Water Resources (DWR) from 2010 and 2011 document California black
21 rail occurrences in 21 discrete habitat patches located in the central Delta portion of the Plan Area.

22 This elusive species spends the majority of its life on the ground and hidden in the wetland and
23 adjacent upland canopy, where it forages, breeds, and winters (Evens et al. 1991). The species is not
24 particularly social and does not congregate or flock. While little information is available on its
25 foraging behavior, it is assumed to be an opportunistic daytime feeder that forages exclusively in
26 wetland habitat, presumably on or near the ground at the edges of emergent vegetation (Evens et al.
27 1991). Daily movements are apparently restricted to the breeding or wintering territory and thus
28 are highly localized and below the wetland and adjacent upland canopy. Movement above the
29 wetland canopy occurs primarily during local, seasonal migration and juvenile dispersal, which
30 occurs from August to October (Trulio and Evens 2000).

31 The Suisun Marsh population is at least 15 miles (24 kilometers) from the north-south powerline
32 right-of-way and unlikely to be affected by its presence. While the proposed north-south powerline
33 right-of-way does not intersect or is immediately adjacent to any known California black rail
34 occurrences, many of the small populations found in the central Delta are within 4 miles
35 (6 kilometers) of the proposed right-of-way (Exhibit 2, Figure 2-1). These sites represent a relatively
36 small proportion of the population in the Bay-Delta region; however, these populations may
37 contribute to the overall range and dispersal capabilities of the species.

38 As a taxon, rails are known to suffer mortality from powerline collision, likely associated with transit
39 between foraging areas and/or local, seasonal migration (Eddleman et al. 1994). Due to their wing
40 shape and body size, rails also have low to moderate flight maneuverability (Rayner 1988; Bevanger

1 1998), increasing susceptibility to collision mortality. However, there are relatively few occurrences
2 of California black rail collisions with overhead wires. Several factors contribute to the relatively low
3 collision susceptibility in this subspecies. Most important among these are daytime site fidelity and a
4 lack of long-distance night migration, considered a principal factor contributing to collision
5 mortality of the species (Eddleman et al. 1994). Movements within the Plan Area are likely short,
6 seasonal, and at low altitudes, typically less than 16 feet (5 meters) (Eddleman et al. 1994).
7 Therefore, while the species may have low to moderate flight maneuverability, its behavior
8 (e.g., sedentary, nonmigratory, ground-nesting and foraging, solitary, no flocking, secretive) reduces
9 potential exposure to overhead wires and vulnerability to collision mortality. No further analysis of
10 California black rail is recommended.

11 **2.2 California Clapper Rail**

12 There are very few occurrences of California clapper rails in the Plan Area. Surveys in Suisun Marsh
13 between 2005 and 2008 found rails only at First Mallard Branch, Rush Ranch, and Goodyear Slough.
14 These surveys estimated the California clapper rail population at less than 13 individuals. The
15 closest occurrence is 20 miles (32 kilometers) from the proposed powerline location. The closest
16 modeled habitat is a little over 11 miles (18 kilometers) from the proposed powerlines, with 1,493
17 acres of modeled habitat within 20 miles (32 kilometers) (Exhibit 2, Figure 2-2). Isolated patches of
18 suitable habitat may occur in the Plan Area as far east as (but not including) Sherman Island. Home
19 range and territory of the California clapper rail is not known, but in locations outside of California,
20 clapper rail territory ranges 0.3 acre to 8 acres (0.1 to 3.2 hectares) (Rush et al. 2012), indicating
21 that known occurrences are not likely to intersect with the proposed lines. The California clapper
22 rail is nonmigratory; however, some local, seasonal movements occur (e.g., between the north San
23 Francisco Bay area and the south San Francisco Bay area), probably in response to seasonal
24 hydrologic changes and their effect on habitat availability and quality. The location of the current
25 population and suitable habitat for the species make collision with the proposed powerlines highly
26 unlikely. No further analysis of California clapper rail is recommended.

27 **2.3 Greater Sandhill Crane**

28 Greater sandhill cranes overwinter in the Plan Area, including large roost sites on Staten, Bouldin
29 and Tyler Islands, Stone Lakes National Wildlife Refuge (NWR), and Brack and Canal Ranch Tracts
30 (Pogson and Lindstedt 1991; Littlefield and Ivey 2000; Ivey and Herziger 2003). Most of the Delta
31 winter range of the species occurs in the Plan Area. During the winter months (October through
32 March) approximately 2,000 greater sandhill cranes forage and roost in proximity to the proposed
33 powerlines (Exhibit 2, Figure 2-4). Ivey and Herziger (2003) estimated average winter home range
34 sizes of greater sandhill cranes in the Delta to be 0.66 square mile (1.7 square kilometers), varying
35 from 0.07 to 2.12 square miles (0.18 to 5.5 square kilometers). Average distance between roost sites
36 and feeding areas was estimated by Pogson (1990) to be 1.74 miles (2.8 kilometers) and by Ivey and
37 Herziger (2003) to be 0.88 mile (1.4 kilometer) (range 0.17 to 1.89 miles [0.27 to 3 kilometers]).
38 Active during the day, sandhill cranes fly frequently between roost and foraging areas, after which
39 they settle down at traditional roost sites for the night.

1 Several aspects of the species' behavior and morphology make greater sandhill cranes particularly
2 susceptible to collisions with overhead wires. Most importantly, flight altitudes during daytime
3 movements are within the range of heights for the proposed lines (50 to 110 feet [15 to
4 33.5 meters]). Therefore, the species is frequently in the risk zone, which increases collision
5 potential.

6 Because most crane movement occurs within 2 miles (3.2 kilometers) of their primary roost, the
7 proximity of the powerlines is a key issue in evaluating collision risk for cranes. Several known
8 roosting sites are less than 2 miles (3.2 kilometers) from the proposed alignment (Exhibit 2, Figure
9 2-4) and are known to intersect with traditional flight patterns (Ivey pers. comm.). Delta wintering
10 cranes are also regularly exposed to dense fog, which limits visibility and increases mortality risk
11 from collision with powerlines. While overall movement may decrease during foggy conditions,
12 greater sandhill cranes are known to fly in the fog, increasing their susceptibility to collision with
13 overhead wires. In addition, this species flies in flocks moving several times a day between feeding
14 and roosting areas. Flocking behavior increases collision risk compared to non-flocking species due
15 to decreased visibility toward the end of the flock. Lastly, the crane's large body size, with high wing
16 loading/low aspect ratio, limits maneuverability making cranes vulnerable to collision relative to
17 more agile species.

18 In addition to collision as a result of daytime travel between roosts and foraging areas, cranes also
19 experience nighttime mortality when flushed from their roosts (e.g., by coyotes), further
20 contributing to an increased risk of collision when powerlines are located near roost sites.

21 Migration flight could cause limited risks for cranes. Cranes arrive in the Delta region beginning in
22 early September, where they reside until late February to early March, when they begin their
23 northward migration back to the breeding grounds. Migration flights usually begin after mid-
24 morning, when thermals develop and finish before or just after sunset. During migration, birds fly at
25 altitudes of up to 11,800 feet. (3,600 meters), with most flights between 490 and 2,500 feet.
26 (150 and 760 meters), far above the height of proposed powerlines (Tacha et al. 1992). However,
27 cranes are exposed to collision risk during takeoff and landing associated with migration.

28 Collectively, the species' foraging and flocking behavior, its presence during winter months of
29 reduced visibility, and its lack of maneuverability make this species highly vulnerable to powerline
30 collision. This assessment concurs with findings in the published literature describing crane
31 mortality as a result of powerline collision (Avian Power Line Interaction Committee 1994;
32 Bevanger 1994; Bevanger 1998; Brown et al. 1987; Brown and Drewien 1995; Hunting 2002a; Yee
33 2008). Because of the crane's high level of vulnerability to powerline collision, an additional
34 assessment of mortality, minimization, and mitigation is provided below.

35 **2.4 Least Bell's Vireo**

36 Least bell's vireo is not currently found in the Plan Area and there are no records of least Bell's
37 vireos breeding in the Plan Area since at least the 1970s. Two singing males were detected in the
38 Yolo Bypass Wildlife Area in mid-April 2010, and again in 2011; no least Bell's vireos were detected
39 in the Yolo Bypass Wildlife Area in 2012 (California Department of Fish and Game 2012). The
40 species typically occurs in early to mid-successional riparian habitat, which is used to meet all of its

1 life requisites. Least Bell's vireo are rarely observed in open habitats away from riparian vegetation.
2 The species does not form flocks and generally remains at or below the riparian canopy. Other than
3 narrow and sparse patches along watercourses, suitable early-to mid-successional riparian habitat
4 is relatively uncommon the Plan Area and particularly in the vicinity of the proposed powerlines.
5 While the species is expected to recolonize the Plan Area during the permit term, this is expected to
6 occur primarily in response to BDCP riparian restoration, which will occur largely in Conservation
7 Zone 7, outside the 6-km buffer zone for the new powerlines (Exhibit 2, Figure 2-5). Territory size
8 ranges from 0.5 to 7.5 acres (0.2 to 3 hectares), but on average are between 1.5 and 2.5 acres (0.6
9 and 1 hectare) in California (U.S. Fish and Wildlife Service 1998). The lack of occurrences in the Plan
10 Area, the lack of current and future higher value habitat patches in the vicinity of the proposed
11 powerlines, and the behavior and habitat requirements of the species make collision with the
12 proposed powerlines highly unlikely. No further analysis of least Bell's vireo is recommended.

13 **2.5 Suisun Song Sparrow**

14 The range of the Suisun song sparrow extends eastward into the Plan Area to approximately Kimball
15 Island. There are several reported occurrences from Kimball Island, Browns Island, and in the
16 Suisun Marsh in the western portion of the Plan Area. These known occurrences, along with areas of
17 suitable habitat, are far from both of the proposed North-South and East-West powerline routes
18 (Exhibit 2, Figure 2-6). During the breeding season, the Suisun song sparrow occupies small
19 territories (approximately 0.1 acre [0.04 hectares] in optimal habitat), usually adjacent to the
20 territories of other Suisun song sparrows in a single linear arrangement along the edges of sloughs
21 and bays. During the fall and winter, adults and young may range up to 600 feet (183 meters) from
22 the territory and occupy adjacent seasonal marshes or grasslands, while continuing to occupy the
23 same general area and return to the same breeding territory each year (Marshall 1948; Walton
24 1975), indicating that known occurrences are not likely to intersect with the proposed lines.
25 Location of the current population, behavior, range, and suitable habitat in the plan area make
26 collision with the proposed powerlines highly unlikely. No further analysis of Suisun song sparrow
27 is recommended.

28 **2.6 Swainson's Hawk**

29 Swainson's hawks are found in the Plan Area from early March through mid-September. A small
30 number, from approximately 16 to 30 individuals, is also known to overwinter in the Delta
31 (Exhibit 2, Figure 2-7) (Herzog 1996). A relatively common breeding raptor in the Plan Area, the
32 nesting distribution extends throughout most of the Plan Area, and foraging is likely to occur in
33 agricultural and grassland habitats. At least 85 nests were documented throughout the Delta during
34 limited surveys in 2009, and a total nesting population of at least 300 nesting pairs is estimated
35 within the Plan Area, including occurrences near proposed powerline facilities. A very dense nesting
36 population occurs immediately west of the Plan Area boundary in Yolo and Solano Counties (Estep
37 2008; LSA 2004). The species is territorial during the breeding season, particularly near the nest site
38 but will also forage communally with other Swainson's hawks away from its nest. During migratory
39 and wintering periods, the species is more social, foraging and migrating in groups (Estep 1989;
40 Babcock 1995; England et al. 1997). However, while the species does congregate in foraging and

1 premigratory groups, individual movements are independent of the group, thereby minimizing
2 collision risk for groups of Swainson's hawks, as opposed to typical flocking behavior where
3 individual movements are more interdependent and thus may increase collision risk for birds
4 within the flock.

5 The species is an aerial predator that hunts primarily from the wing typically at altitudes ranging
6 from 98 to 295 feet (30 to 90 meters) (Estep 1989; England et al 1997), although higher altitudes
7 have been reported (Fitzner 1980). Circling above grassland and farmland foraging habitats, prey
8 are captured by rapidly diving or stooping toward the ground. Other typical flight behaviors include
9 high-elevation courtship flight and high-elevation, midday soaring. While Swainson's hawks hunt
10 within the range of heights proposed for the new powerlines (50 to 110 feet [15 to 33.5 meters]),
11 their keen vision and high maneuverability substantially reduce powerline collision risk for the
12 species. Like other diurnal raptors, Swainson's hawks have highly developed eyesight (Jones et al.
13 2007), allowing them to detect small prey while hunting from relatively high altitudes. Keen
14 eyesight also allows for detection and avoidance of other aerial objects, including aboveground
15 utility lines. Like many other Falconiformes, Swainson's hawk has a long, narrow, tapered wings and
16 body size that allow for efficient soaring flight and highly developed aerial maneuverability. In
17 addition, Swainson's hawks are rarely active during inclement weather and are not typically
18 observed in flight during rainy or foggy conditions (Fitzner 1980).

19 The species' general maneuverability, its keen eyesight, and fair-weather flight behavior, make it a
20 low relative risk for powerline collision mortality. Mortality associated with powerline collision is
21 not anticipated to affect the Plan Area population, and no further analysis of Swainson's hawk is
22 recommended.

23 **2.7 Tricolored Blackbird**

24 Historical records indicate breeding colonies of the tricolored blackbird have occurred within the
25 Plan area along the eastern edge of the Suisun Marsh in Bird's Landing, west of French Camp along
26 the south eastern edge of the plan area, and locations outside of the Plan Area including areas near
27 Davis, Napa, Elk Grove, Vernalis, and two occurrences just north of the Plan Area boundary
28 (California Department of Fish and Game 2012). More recent surveys conducted in the last 15 years
29 have documented tricolored blackbird breeding colonies throughout the Plan Area at sites near Yolo
30 Bypass; near Stockton, Manteca, and Tracy in the southeastern corner of the Plan Area; north of
31 Bradford Island; and along the eastern edge of Suisun Marsh; and along the Sacramento River Deep
32 Water Ship Channel (Information Center for the Environment 2011; Meese 2011). Breeding colonies
33 have also been recorded just outside of the plan area within the past 15 years south of the Plan Area
34 along the San Joaquin River, just outside of the southwest Plan Area boundary, near Vallejo, and east
35 of Woodland outside of the northwest Plan Area boundary (Information Center for the Environment
36 2011; Meese 2011).

37 A single nesting colony of about 1,000 breeding adults was recorded during the 2011 statewide
38 survey in the Plan Area along the northern edge of Suisun Marsh (Information Center for the
39 Environment 2011). Between 2009 and 2011, DWR biologists surveyed several thousand acres of
40 potentially suitable tricolored blackbird nesting habitat in the Plan Area (excluding Suisun Marsh

1 and the Potrero Hills area) during the optimal breeding period and detected no nesting colonies
2 (Delta Habitat Conservation and Conveyance Program 2011). There are 31 occurrences within 5
3 miles (8 kilometers) (Exhibit 2, Figure 2-8). All observations appeared to be foraging birds; no
4 nesting by tricolored blackbirds was confirmed. Although recent nesting colonies in the Plan Area
5 have generally been small (comprising less than 2,000 breeding adults), several larger colonies have
6 been reported from just outside the Plan Area, including colonies of 35,000, 57,000 and
7 18,900 breeding adults on the Conaway Ranch in the Yolo Bypass north of Interstate 80 in 2007,
8 2009, and 2010, respectively.

9 In the Central Valley some tricolored blackbird populations are resident, residing all year in the
10 Central Valley, while some migrate, moving in large flocks from inland breeding locations to
11 wintering habitats in the Delta and coastal areas. Generally, overwintering birds roost in areas
12 dominated by emergent wetland vegetation in and around Bird's Landing in southern Solano County
13 and forage primarily in association with cultivated lands (e.g., irrigated and non-irrigated pasture,
14 rice, corn) between Sacramento and Stockton.

15 Tricolored blackbirds exhibit different flight behaviors during the nesting and wintering seasons.
16 When nesting, tricolored blackbirds are likely to travel shorter distances between the nesting site
17 and foraging grounds. In order to transport food items back to the nest, they make multiple trips a
18 day between the nest site and foraging grounds. The nature of foraging behavior during the nesting
19 season naturally results in lower flight heights, more direct flight patterns, and smaller, more loosely
20 formed flocks. Lower flight heights means most breeding birds are flying beneath the height of most
21 wires and, where lower wires exist, individuals or small flocks of birds can maneuver to avoid them
22 without issue (Meese pers. comm.).

23 During the winter, tricolored blackbirds migrate into the Plan Area in large flocks. Altitude during
24 migration is not known, but it is likely that birds have greater potential to strike the proposed
25 powerlines (50 to 110 feet [15 to 33.5 meters]) during migration than during nesting. Wintering
26 birds make daily flights between roosting sites, which are located primarily near Bird's Landing in
27 southern Solano County, and foraging grounds, which are cultivated land types found throughout
28 the Plan Area. Although tricolored blackbirds leave from and return to wintering roost sites in very
29 large flocks, they forage throughout the day in smaller flocks. These smaller flocks move between
30 foraging locations primarily through low-altitude flights. While tricolored blackbirds are likely more
31 vulnerable during migration and overwintering due to larger flock size, likely increased flight
32 altitudes, and dense fog that is common to the area, there has been no evidence of mortality due to
33 collision with overhead wires (Meese pers. comm.).

34 In summary, tricolored blackbirds have the potential to intersect the proposed powerline routes
35 largely due to winter movements throughout the Plan Area. While migratory flight behavior may
36 increase the risk of strike hazard, daily movements associated with winter foraging likely occur
37 below the height of the lines. In addition, tricolored blackbirds are considered strong and agile flyers
38 with moderately maneuverability (i.e., low wing loading/low aspect ratio) (Beedy et al. 1999) and
39 therefore physically equipped to avoid collision with powerlines. Current scientific evidence and
40 best professional judgment suggest that powerlines are not a significant cause of mortality for
41 tricolored blackbirds (Meese pers. comm.). Mortality associated with powerline collision is not

1 anticipated to affect the Plan Area population, and no further analysis of tricolored blackbird is
2 recommended.

3 **2.8 Western Burrowing Owl**

4 While nesting and wintering burrowing owls could occur in grassland, pastureland, and agricultural
5 habitats throughout most of the Plan Area, the majority of reported occurrences indicate that the
6 species is concentrated in grassland and pasturelands west of the Sacramento Deep Water Ship
7 Channel in Yolo and Solano Counties and in the grassland habitats along the western edge of the
8 Plan Area (roughly between Brentwood/Antioch and Tracy). The species is also found in lower
9 densities elsewhere in the Plan Area, with documented occurrences on Brannan Island and near
10 Suisun Bay and Clifton Court Forebay, and the species may occur elsewhere where habitat, such as
11 grassland and pastureland, is available. Burrowing owls persist in some cultivated or ruderal
12 habitats, such as near Stockton where they are typically found along levees, canals, field edges, and
13 some ruderal habitats or idle fields. However, few burrowing owls have been reported from the
14 central portion of the Delta and the northern Delta east of the Sacramento Deep Water Ship Channel,
15 probably due to regular cultivation, lack of undisturbed habitats, and lack of ground-squirrel
16 populations. The few active sites in this area are generally restricted to levee embankments and
17 along irrigation canals. The species is a year-round resident in the Plan Area; however, local
18 migratory patterns and the extent to which migrants occupy the Plan Area during the non-breeding
19 season are unclear.

20 Twenty five occurrences are within 1 mile of the southern end of the proposed powerline alignment,
21 and 115 known occurrences are within 5 miles (8 kilometers) of the east-west segment of the
22 northern end (Exhibit 2, Figure 2-9). Potential habitat consisting of high- and low-value grassland is
23 mapped along both the northern and southern portions of the line.

24 Western burrowing owls forage throughout the day but are largely crepuscular, hunting mostly at
25 dusk and dawn. Hunting in low light can be a risk factor for powerline collision. However owls have
26 acute eyesight adapted to low-light conditions and a wide range of vision. In addition, the species
27 feeds primarily on the ground where it catches insects by walking and hopping or catching from
28 burrow mound or perch (Haug et al. 1993). Burrowing owls may hunt vertebrates from both perch
29 and by hovering low to the ground. Hunting typically occurs at about 33 feet (10 meters) above
30 ground, while direct flights back to the nest (prey delivery) were 3 to 6 feet (1 to 2 meters) above
31 ground and at a flight speed of 33 miles per hour (53 kilometers per hour), keeping the owl out of
32 the range of proposed powerlines (Poulin et al. 2011).

33 The species is large-bodied but with relatively long and rounded wings, making it moderately
34 maneuverable. While burrowing owls may nest in loose colonies, they do not flock or congregate in
35 roosts or foraging groups. Collectively, the species' keen eyesight and largely ground-based hunting
36 behavior make it a relatively low-risk species for powerline collision. While the species is not
37 widespread in the Plan Area, it may become more widely distributed as grassland enhancement
38 improves habitat for the species. Even so, the risk of effects on the population are low, given its
39 physical and behavioral characteristics. No further analysis of western burrowing owl is
40 recommended.

2.9 Western Yellow-Billed Cuckoo

The western yellow-billed cuckoo is a rare summer resident in California with a disjunct breeding distribution extending through the interior of the Central Valley. While the Plan Area is within the species' breeding range, there have been no confirmed breeding records for the Plan Area or vicinity for several decades (Exhibit 2, Figure 2-10). Studies conducted since the 1970s indicate that there may be fewer than 50 breeding pairs in California (Gaines 1974; Halterman 1991; Laymon et al. 1997). While a few occurrences have been detected elsewhere recently, the only locations in California that currently sustain breeding populations include the Colorado River system in southern California, the South Fork Kern River east of Bakersfield, and isolated sites in remnant riparian patches along the Sacramento River in Glenn, Butte, and Tehama Counties (Laymon and Halterman 1989; Laymon 1998).

While there are few historical records from the Plan Area, presumably the species nested along the Sacramento, San Joaquin, and Mokelumne Rivers and along smaller tributary drainages, including Lost Slough, White Slough, and Disappointment Slough. In 2009, DWR detected one and possibly two western yellow-billed cuckoos in a remnant patch of riparian forest near Mandeville Island. However, breeding status was not confirmed. The Plan Area supports several remnant riparian patches in the vicinity of Mandeville and Medford Islands that provide suitable riparian vegetation for cuckoos but may not provide sufficiently large patch size to support breeding cuckoos.

Portions of both the Sacramento and Mokelumne Rivers are very near to the proposed powerline, with several sections occurring less than 1 mile from these rivers. One occurrence is within one mile (2 kilometers) of the proposed powerline alignment and another is within five miles (8 kilometers). However, based on the species' current status and distribution in the Plan Area, risk of collision with proposed powerlines is very low. Habitat in the Plan Area will be enhanced and the status of the western yellow-billed cuckoo may improve.

Because the western yellow-billed cuckoo uses riparian forests to meet all of its breeding and wintering life requisites, the species remains primarily within the canopy of riparian forests and rarely ventures into open spaces except during migration, limiting its opportunity to encounter the proposed powerlines. As a summer resident, the species occurs in the Plan Area during periods of relatively high visibility and clear weather conditions, thus further reducing collision risk from daily use patterns or seasonal migration flights. Finally, western yellow-billed cuckoo wing shape is characterized by low wing loading and a moderate aspect ratio, making the species moderately maneuverable (Bevanger 1998) and presumably able to avoid collisions, especially during high-visibility conditions.

Because of its rarity in the Plan Area, its proclivity to remain in the riparian canopy, its presence during periods of relative high visibility, and its overall ability to successfully negotiate around overhead wires that it may encounter, the western yellow-billed cuckoo is considered to have a very low susceptibility to collision with overhead wires. No further analysis of the western yellow-billed cuckoo is recommended.

2.10 White-Tailed Kite

The white-tailed kite is a year-round resident in the Plan Area, although relatively few nesting locations have been documented. The California Natural Diversity Database (CNDDDB) reports only five locations within the Plan Area (California Department of Fish and Game 2011). Nesting occurrences have been reported in the Delta, along the Sacramento River west of Stone Lake, and in the north-central and east-central Delta. Recent surveys in the Yolo and Sacramento County portions of the Plan Area have documented active nests sites in riparian habitats in the Yolo Bypass and along Steamboat and Georgiana sloughs, and the Sacramento River (Estep 2007, 2008). Surveys from 2009 to 2011 documented 10 active white-tailed kite nest sites (Delta Habitat Conservation and Conveyance Program 2011).

Several of the known occurrences are within 5 miles of both the proposed powerline North-South and East-West routes. Along the north-south route, known occurrence locations have been recorded within 1 mile of the proposed powerline (Exhibit 2, Figure 2-11). Nesting distribution is limited by the dearth of suitable trees in much of the central Delta, and nesting density in that area is likely significantly lower than that found in the northern and southern portions of the Plan Area. The species is territorial, defending relatively small home ranges ranging from approximately 4 to 296 acres (1.6 to 120 hectare) (Dunk and Cooper 1994; Waian 1973; Henry 1983). While tolerant of conspecifics, the species does not flock or typically engage in communal foraging except during the winter when communal roosts will form.

The white-tailed kite is an aerial predator that hunts primarily from the wing at altitudes ranging from 5 to 25 meters. Hovering, or kiting, the kite captures prey by dropping or stooping vertically toward the ground. Other flight behaviors include aerial courtship displays and territorial defense, which the kite engages in near the nest. While white-tailed kite flight behavior puts them regularly within the range of heights proposed for the new transmission lines (50 to 110 feet), their keen vision and high maneuverability substantially reduce powerline collision risk for the species. Like other diurnal raptors, white-tailed kites have highly developed eyesight (Jones et al. 2007), allowing them to detect small prey while hunting from relatively high altitudes. Keen eyesight also allows for detection and avoidance of other aerial objects, including above-ground utility lines. Like many other Falconiformes, the white-tailed kite has long, narrow, tapered wings and body size that allow for efficient soaring flight and highly developed aerial maneuverability. While kites occur in the Central Valley during the winter months when dense fog can reduce visibility, the species is not usually active during inclement weather and not typically observed in flight during rainy or foggy conditions.

Therefore, while the species may be frequently within the risk zone of the proposed powerlines, its general maneuverability, its keen eyesight, and lack of flocking behavior make it a low relative risk for powerline collision mortality. Mortality associated with the proposed powerlines is not anticipated to affect the Plan Area population.

2.11 Yellow-Breasted Chat

The yellow-breasted chat is a neotropical migrant songbird whose range extends from southern Canada to Mexico. Comrack (2008) includes the central Delta within the current breeding range of the yellow-breasted chat. There are few breeding records of the species in the Plan Area. Most occurrences are fall and winter migrants found along Putah Creek near the northern edge of the Plan Area in Yolo and Solano Counties or along the Cosumnes River in the Cosumnes River Preserve. In 2008, the National Audubon Society noted pairs of yellow-breasted chat at Liberty Island, Sherman Island, and Piper Slough in the central Delta. Recent field surveys have confirmed late spring and summer occurrences of chats in the Plan Area (Delta Habitat Conservation and Conveyance Program 2011). Ten occurrences are within one mile (2 kilometers) of the proposed powerlines and 18 are within 5 miles (Exhibit 2, Figure 2-12).

A total of 51 nest sites were identified from 2009 to 2011 (Delta Habitat Conservation and Conveyance Program 2011) within the Plan Area. Territory size ranges from 0.3 to 3.2 acres (0.1 to 1.3 hectares) (Zeiner et al. 1990). Territory sizes have not been measured in California, but in California riparian habitat, breeding densities ranged from 6.5 to 27 males per 247 acres (100 hectares) (Eckerle and Thompson 2001) and Gaines (1974) reported a breeding density from the Sacramento Valley of one chat per 10 acres (4 hectares).

Yellow-breasted chats nest and forage in dense riparian thickets of willows, vines, and brush associated with streams and other wetland habitats. With moderate wing loading and a moderate aspect ratio, the species usually flies through dense vegetation, starting from a high perch and ending on a higher perch or in low, dense vegetation and only occasionally crosses open fields, flying at altitudes of less than 3.2 feet (less than 1 meter), virtually eliminating the risk of collision with proposed powerlines. When foraging, the solitary species gleans prey from foliage of low, dense shrubs or from the ground.

Yellow-breasted chats are migratory and usually arrive at California breeding grounds in April from their wintering grounds in Mexico and Guatemala. Departure for wintering grounds occurs from August to September. These are periods of relative high visibility when the risk of powerline collisions will be low. The species' small, relatively maneuverable body; its foraging behavior; and its presence in the Plan Area during the summer contribute to a low risk of collision with the proposed powerlines. No further analysis of the yellow-breasted chat is recommended.

3.0 Greater Sandhill Crane Effects Analysis

Based on the vulnerability analysis developed above, the greater sandhill crane is the only covered species to exhibit a high risk for collision with proposed powerlines, using the criteria of exposure and sensitivity. This is consistent with the published literature and expert opinion. Therefore, additional efforts to contextualize and quantify risks were developed for the greater sandhill crane.

1 **3.1 Collision Risk Map**

2 A map of the distribution and risk of greater sandhill crane from powerline collisions in the Plan
3 Area was developed to represents the risk of collision spatially and to help identify powerline routes
4 that minimize risk to greater sandhill crane. Over time, the powerline alignment has been
5 significantly shortened to reduce the potential loss of greater sandhill crane due to strikes. DWR
6 engineers, greater sandhill crane experts, and conservation land managers continue to discuss
7 alignment optimization alternatives to further reduce impacts to individuals as well as to roosting
8 and foraging habitat.

9 Surveys of greater sandhill cranes were conducted during the winters of 2006–2007, 2007–2008,
10 and 2008–2009 by automobile, aircraft, and on foot (Ivey et al. in preparation [a]), and birds
11 outfitted with transmitters were tracked to identify roosting and foraging areas. These efforts
12 quantify the approximate number of night-roosting greater sandhill cranes, with estimates in a roost
13 site complex ranging from 10 to 1,500 birds (Staten Island).

14 Greater sandhill cranes outfitted with radio transmitters (n = 33) were used to determine the
15 distance between roost sites and foraging areas and the proportion of birds that foraged within
16 different distance intervals of the roost. In other words, studies determined the proportion of the
17 roosting population that can be expected to forage within 1, 2, and 3.7 miles (2, 4, and 6 kilometers)
18 of the roost. Results indicate that all greater sandhill cranes (100%) forage within 1.2 miles
19 (2 kilometers) of the roost site, 18% between 1.2 and 2.5 miles (2 and 4 kilometers) of the roost, 9%
20 between 4 and 5 kilometers, and 5% between 3 and 3.7 miles (5 and 6 kilometers) (Ivey et al. in
21 preparation [b]). In order to weight risk relative to the size of a given roosting site, the number of
22 birds at each roost was divided by 1,500 (the maximum number of greater sandhill crane at a roost-
23 site complex). Using this method, the largest roost site would be standardized to a value of 1 and the
24 smallest roost site (10 birds) would be assigned a value of 0.0067 (10/1500). This value was then
25 multiplied by the percentages derived above to determine the relative risk in a given area based on
26 roost size and distance from the roost. This final number is the collision risk index value. Results
27 were made spatially explicit in ArcGIS, where each cluster of roost sites was buffered by a radius of
28 1, 2, 3, and 3.7 miles (2, 4, 5, and 6 kilometers), and collision risk index values were mapped within
29 those distance categories. In cases where the roost-site buffers overlapped, the values were added
30 together (i.e., risk in that area increased). The final collision risk index values were grouped in the
31 following ranges: 0.001 to 0.01, 0.01 to 0.1, 0.1 to 0.2, 0.2 to 0.4, 0.4 to 0.6, 1.0 to 1.2, and 1.2 to 1.4
32 (no values between 0.6 and 1.0 were found) and are color coded in Figure 2, which visually
33 represents collision risk within the Plan Area.

34 **3.2 Estimated Collision Mortality of Greater Sandhill Crane**


35 To calculate mortality in the Plan Area, the collision risk index numbers for polygons associated with
36 a particular roost were used to estimate “crossings” where proposed powerlines intersect mapped
37 polygons (Figure 2). Some of these risk polygons overlap and have higher collision risk values
38 because birds from adjacent roosts use the same areas to forage. The values for polygons that
39 intersect a potential line segment associated with a particular roost were averaged, weighted by

1 length of line crossing them, to estimate the number of cranes expected to cross those lines on a
2 daily basis.

3 Using this approach, an average population size was determined for each line segment, which was
4 then multiplied by 130 days (the mean number of days that greater sandhill crane spend in the Delta
5 wintering area) and by four flights per day (birds going between foraging areas and roost sites twice
6 a day, crossing the lines twice in the morning and twice in the evening). Based on the assumption
7 that the probability of flying out of the roost in a given cardinal direction is 25%, this number was
8 then divided by four, resulting in a crossing estimate for each segment and for the total line (Table
9 2.). The number of crossings was then multiplied by collision mortality rates that were calculated for
10 greater sandhill crane in the Rocky Mountains of Colorado (Brown and Drewien 1995). These data
11 were used because local or regional data are not available. Brown and Drewien (1995) estimated
12 that annual collision mortality of greater sandhill crane at unmarked lines was between 2.5×10^{-5}
13 (low estimate) and 30.4×10^{-5} collisions per crossing (high estimate). For the purposes of this
14 analysis, the high estimate was used to ensure that all potential impacts were captured.

15 Because lack of visibility is one of the most commonly implicated causes of collision mortality, live
16 or ground wires can be marked to increase their visibility. While it hasn't been studied, the efficacy
17 of bird flight diverters are likely diminished with reduced visibility associated with the new moon or
18 fog. However, it is reasonable to assume that bird flight diverters still reduce mortality. Other
19 markers also include dampers, hanging plates, and spheres. Marking lines has been shown to
20 decrease collision risk substantially. Brown and Drewien (1995) estimated that annual collision
21 mortality rates of birds at marked lines were reduced by 62 and 66% for two types of markers, and
22 it is likely that birds found dead in these studies were also flying at night. Morkill and Anderson
23 (1991) indicated a 54% reduction in crane mortality at marked lines. In addition to the risk map
24 derived above, collision risk and mortality in the Plan Area were estimated relative to the proposed
25 powerline locations. This was done for both marked and unmarked lines.

26 Absent line marking, which increases visibility and reduces collision risk (i.e., without minimization
27 measures), the potential annual take of greater sandhill crane is estimated at 18 per year at
28 permanent lines and 120 per year at temporary lines. Assuming a reduction of 66% (Brown and
29 Drewien 1995), potential mortality at marked lines is estimated at 7 per year at permanent lines and
30 41 per year at temporary lines.



1

2

Figure 2. Collision Risk Index Map for Greater Sandhill Crane

1 **Table 2. Estimated Collision Mortality of Greater Sandhill Crane at BDCP Marked and Unmarked**
 2 **Powerlines**

Powerline Type	Crossings/Year ^a	Deaths/Year ^b (unrounded)	
		Unmarked Lines	Marked Lines ^c
69-kV line (permanent)	749,949	16 (15.18)	6 (5.16)
230-kV line (permanent)	6,586	2 (2.00)	1 (0.68)
230-kV line (temporary)	321,120	96 (95.89)	33 (32.60)
34.5-kV line (temporary)	76,862	24 (23.37)	8 (7.95)

^a Baseline mortality = $30.4 \times 10^{-5} \times$ crossings/year.
^b Values have been rounded up to the nearest integer unless otherwise specified.
^c 66% reduction based on Brown and Drewien (1995) for sandhill cranes in Colorado.

3
 4 Based on the analysis above, the cumulative mortality associated with marked temporary lines is
 5 estimated to be 410 birds over a 10-year period. While it is possible to calculate cumulative impacts
 6 from permanent lines over the permit term, mortality will continue at these lines as long as they are
 7 present. Therefore, deaths per year is a better metric for describing mortality at permanent lines.
 8 Note that mitigation is also calculated on an annual, ongoing basis.

9 4.0 Population Impacts

10 Greater sandhill cranes that winter in the Plan Area are designated as the Central Valley population
 11 (Pacific Flyway Council 1997). Although there is no current estimate for the Central Valley
 12 population, recent counts of summering cranes in California, Oregon, and Washington total
 13 approximately 4,200 (Ivey and Herziger 2000, 2001), and a recent estimate of summering cranes in
 14 interior British Columbia totaled an additional 4,000 (Breault pers. comm.). These birds are all
 15 within the same regional population; resulting in a total population of approximately 8,200 birds
 16 (also see Littlefield 2002).

17 Assuming a population of 500 birds in 1945 (based on literature reporting less than 200 pairs in
 18 Oregon and California) (Gabrielson and Jewett 1940; Walkinshaw 1949) and 8,200 birds in 2012
 19 (Littlefield 2002), the overall annual rate of increase is 1.4% per year. Because cranes are long-lived
 20 with relatively low recruitment rates and high annual survival rates (usually greater than 90%)
 21 (Tacha et al. 1992; Drewien et al. 1995), additional mortality is unlikely to be compensated by
 22 population growth, and losses could directly affect population dynamics. Also, greater sandhill
 23 cranes are highly faithful to wintering sites and are primarily sedentary during winter, so birds that
 24 roost close to proposed powerlines are particularly vulnerable. Note that the current rate of growth
 25 accounts for existing sources of mortality for greater sandhill crane, such as collision at existing

1 lines. We do not make predictions about future changes in other sources of mortality outside the
2 Plan Area other than covered activities.

3 Table 3 summarizes the impacts of the estimated annual take (Table 2) on the Central Valley greater
4 sandhill crane population as a percentage of the total population. A population decline is expected if
5 the impact exceeds the estimated rate of population increase (1.4%). Table 2 provides the percent
6 impact for marked and unmarked lines using the high estimated collision mortality rates derived by
7 Brown and Drewien (1995). The table displays the effect of proposed powerlines during project
8 initiation, when only the temporary lines affect cranes, and subsequently during operations, after
9 the temporary lines are removed and the permanent lines are in place. There may be a period of
10 time during project construction when both temporary and permanent lines impact cranes. In this
11 case, the impacts from temporary and permanent lines are additive for the period of time that both
12 temporary and permanent lines exist. Using the higher collision mortality rate, the level of take from
13 temporary lines has the potential to exceed the growth rate of the Central Valley population if lines
14 are unmarked.

15 **Table 3. Estimated Impacts on the Central Valley Population of Greater Sandhill Cranes from**
16 **Collisions with Proposed BDCP Power Lines**

Line Type	Annual Impact (%)	
	Unmarked Lines	Marked Lines ^a
Temporary	1.46	0.50
Permanent	0.21	0.07

A population decline is expected if the annual impact is greater than the assumed average rate of population increase (1.4%), marked in dark grey.
^a 66% reduction based on Brown and Drewien (1995) for sandhill cranes in Colorado

17
18 An estimated 2,000 to 3,000 cranes wintered in the Delta in 2008–2009 (Ivey et al. in preparation
19 [a]). Assuming a population of 2,500, the impacts on this subpopulation of greater sandhill cranes
20 will be proportionally greater than impacts on the larger Central Valley population.

21 Based on the same annual growth rate used above (1.4%), absent line marking, the temporary lines
22 will result in a net decline of this subpopulation (losses greater than 1.4%) (Table 4).

23 The most important roost site area in the Delta is Staten Island, where approximately 1,500 greater
24 sandhill cranes have been counted. Therefore, the losses will come largely from this group of birds.
25 The cranes at Staten Island will only be affected by the temporary lines. Other roost sites along the
26 proposed lines support 10 to 300 birds. The second-most important roosts are the Stone Lakes NWR
27 roost sites near the north end of the proposed permanent lines, which support approximately 300
28 birds. The new permanent lines as proposed will affect birds using Stone Lake NWR and Cosumnes
29 River Preserve roost sites.

1 **Table 4. Estimated Impacts on the Delta Wintering Population of Greater Sandhill Cranes from**
 2 **Collisions with Proposed BDCP Power Lines**

Line Type	Annual Impact (%)	
	Unmarked Lines	Marked Lines ^a
Temporary	4.78	1.62
Permanent	0.69	0.24

A population decline is expected if the annual impact is greater than the assumed average rate of population increase (1.4%), marked in dark grey.
^a 66% reduction based on Brown and Drewien (1995) for sandhill cranes in Colorado.

4 5.0 Minimization and Mitigation

5 The analysis above documents potential impacts on greater sandhill crane from the installation of
 6 new temporary and permanent powerlines as part of the BDCP. However, the proposed lines are a
 7 small portion of the existing lines in the Plan Area. Collectively, 4,491 miles of distribution, sub-
 8 transmission, and transmission lines currently exist in the Plan Area (Table 5). New, permanent
 9 lines proposed by the BDCP represent less than 0.5% of the amount of existing lines in the Plan Area.

10 **Table 5. Existing Powerlines in the Plan Area**

Line Type	Voltage (kV)	Length (Miles)
Distribution	<1	35
	4	57
	11	1,655
	12	131
	17	120
	21	1,309
	22	504
	60	170
Sub Total		3,981
Transmission	69	43
	70	2
	115	209
	230	156
	500	100
Sub Total		510
Plan Area Total		4,491

11
 12 Although the risk posed by new lines is small relative to existing lines, any additional impacts to
 13 cranes could be detrimental, as described above. There are several options for minimizing impacts,
 14 including the placement of the proposed lines (which has been revised iteratively reducing impacts),

1 removal of the ground wire, and fitting the ground wire with markers—brightly colored “aviation”
2 balls, thickened wire coils, or luminescent, shiny, or hinged flashing or flapping devices. All of these
3 marker options have the potential to reduce bird collision frequency by as much as 89% (Avian
4 Power Line Interaction Committee 1994).

5 In order to minimize impacts on cranes, the Implementation Office will install bird diverters on all
6 new lines erected as part of the BDCP. Line marking with bird diverters will follow Avian Power Line
7 Interaction Committee protocols.

8 While marking lines substantially decreases collision risk for cranes, it does not eliminate it. Based
9 on our estimates, using the higher collision risk estimate of 30.4×10^{-5} collisions per crossing, a loss
10 of 7 birds per year for permanent lines and 41 birds per year for temporary lines will need to be
11 mitigated to maintain no net loss of greater sandhill cranes (Table 2). In order to compensate for
12 this loss, bird diverters may be placed on existing lines within 2 kilometers of existing roost sites,
13 with priority given to those lines adjacent to larger roost sites.

14 The mitigation value of a given length of powerline can be determined using the same methods
15 described in Section 3.2, *Estimated Collision Mortality of Greater Sandhill Crane*. Instead of using the
16 location of the proposed lines to estimate mortality, as was done above, the location of existing lines
17 is used to quantify the benefit of installing bird flight diverters (BFDs) at a given location. For
18 purposes of analyzing the feasibility and cost of this option, we assume that the mitigation value of
19 retrofitting an existing line with BFDs increases with proximity to a roost site. This effect of the
20 mitigation is scaled to the size of the roost (i.e., the larger the roost population, the greater the
21 mitigation value).

22 To inform feasibility discussions and costing, several potential mitigation sites were identified. This
23 analysis should be rerun at the time that mitigation is implemented. If roost sites, available line
24 segments, collision rates, or other factors differ at that time, the values used below can and should
25 be replaced with improved estimates. The methodology outlined herein and in Box 1 provides the
26 information necessary to adjust mitigation at the time of project implementation.

27 These sites selected below consist of currently unmarked distribution lines near two of the largest
28 greater sandhill crane roosts in the Plan Area: Staten Island and Isenberg Reserve (or Woodbridge
29 Ecological Reserve), along Staten Island and Woodbridge Roads, respectively. However, these
30 mitigation sites may not be available at project implementation, in which case the needed mitigation
31 may be acquired at other lines, using the methods developed in this assessment.

32 Table 6 summarizes the location of these lines and the calculated mitigation value of each, and Box 1
33 provides an example mitigation calculation. The mitigation approach, like the impact approach
34 described in Section 3.2, *Estimated Collision Mortality of Greater Sandhill Crane*, assumes a 66%
35 reduction in mortality based on the installation of BFDs (Brown and Drewien 1995). For temporary
36 lines, the proposed approach significantly over-mitigates impacts to greater sandhill crane because
37 new BFDs on existing lines will be retained long after the temporary lines are removed, and risk of
38 collision is removed.

Box 1: Example Calculation

Assume a line adjacent to the roost sites at Staten Island spans 12 kilometers within 2 kilometers of the roost complex and supports an estimated 1,500 greater sandhill cranes. Then:

- 1,500 cranes x 130 days = 195,000 crossings/year, and
- 195,000 crossings/year x 0.000304 = estimated 59.28 (60) deaths/year at the unmarked line segment.

If the annual reduction in these losses is 66% (Brown and Drewien 1995), then the number of greater sandhill crane deaths avoided is:

- 59.28 mortalities/year x 0.66 = 39.12 (40) mortalities avoided/year.

1
2
3
4
5
6
7
8
9
10

11 **Table 6. Mitigation for Greater Sandhill Crane Mortality on BDCP Powerlines**

Line Segment	Maximum Allowable Take (Per Year for New Marked Lines)	Proposed Mitigation: Mark Existing Lines		Mitigation Example*
		Miles of Line	Reduced Mortality (Cranes/Year)	
Permanent	7	1.5	7	4.3 miles (6.8 kilometers) of line on Staten Island Road, adjacent to the primary crane roost site = 20 cranes/year
Temporary	41	4.4	38	5.5 miles (8.9 kilometers) of line on Staten Island Road, adjacent to the primary crane roost sites= 23 cranes/year. 1 mile of the line on Woodbridge Road, beginning at the entrance road to North Isenberg Reserve, (east for 1 mile; 1.6 kilometers) = 10 cranes. 0.4 miles (0.6 kilometers) of the line that runs east-west along Hog Slough, north of North Isenberg Reserve = 5 cranes.
Total		5.9		

* Mitigation example only to demonstrate feasibility. Actual mitigation would be determined during implementation using this approach to calculating an equivalent reduction of mortality probability.

12
13
14
15
16
17
18
19
20
21
22
23

Total minimization and mitigation costs are based on the types of lines on which BFDs are placed. BFDs cost approximately \$40 per unit (Pleiss pers. comm.). For optimum results, the recommended spacing distance for BFDs is 15 to 16.5 feet (4.5 to 5 meters) (Avian Power Line Interaction Committee 1994). Installation of BFDs in 15-foot (4.5-meter) intervals requires 325 units per mile (222 units per kilometer), or \$13,000 per mile (\$8,880 per kilometer). For distribution lines, installation of BFDs requires a lineman who can install approximately 1 mile of BFDs per day. For transmission lines a helicopter and crew are required (price TBD). The existing high-risk lines proposed for mitigation are all distribution lines. The total cost of mitigation is \$84,180, based on 6.1 miles of distribution. The total cost of minimizing and mitigating the distribution lines through placement of diverters on existing lines is \$122,820 (Table 7). The cost to minimize the transmission lines is to be determined (Table 7). As mentioned above, the mitigation sites proposed in Table 6

1 were chosen to inform feasibility discussions and costing. While impacts from new lines will be
 2 minimized by the addition of BFDs, the location and required length of mitigation lines will need to
 3 be determined at the time of implementation based on field-verified information.

4 **Table 7. Costs Associated with Minimization and Mitigation of Temporary and Permanent Lines**

	Miles of Existing Lines (Mitigation)		Costs (\$/mile)		Total Costs		
	Miles of Proposed Lines	Mitigation	Materials	Installation	Materials	Installation	Total
	Minimization						
Distribution	3	5.9	\$13,000	\$800	\$115,700	\$7,120	\$122,820
Transmission	55	NA	\$13,000	TBD	\$715,000	TBD	TBD

5
 6 Placement of BFDs on existing lines is one of several options that may be implemented to meet a
 7 performance standard of no net increase in bird strike risk for greater sandhill cranes in the Plan
 8 Area. Other options include designing the transmission line alignment to further minimize risk;
 9 removing, relocating, or undergrounding existing lines; and managing cultivating land roost sites to
 10 shift roosting areas away from high risk areas. A combination of options may be implemented to
 11 achieve the standard of no net increase in bird strike risk for greater sandhill cranes in the Plan
 12 Area.

13 6.0 Summary

14 New powerlines proposed by the BDCP have the potential to affect birds in the Plan Area. For all
 15 species except greater sandhill crane, this effect is unlikely to pose high levels of potential risk.
 16 Because of the physical and behavioral characteristics of greater sandhill crane, the species'
 17 propensity to collide with and suffer mortality from powerlines is high. Mortality estimates vary
 18 with the location of the proposed lines relative to roost sites and on the use of line markers, which
 19 reduce collisions. To minimize mortality from the proposed powerlines, the Implementation Office
 20 will install line markers on all BDCP powerlines as they are established, thereby reducing future risk
 21 of mortality by approximately 66%. To compensate for remaining risks and achieve a performance
 22 standard of no net increase in bird strike risk for greater sandhill cranes in the Plan Area, a
 23 combination of options may be implemented. These options may include siting new lines to
 24 minimize risk; removing, relocating, or undergrounding existing lines; managing cultivating land
 25 roost sites to shift roosting areas away from high risk areas; and installing BFDs on existing lines
 26 using the methods described here.

1 7.0 References Cited

2 7.1 Literature Cited

- 3 Avian Power Line Interaction Committee. 1994. *Mitigating Bird Collisions with Power Lines: State of*
4 *the Art in 1994*. Edison Electric Institute. Washington, D.C.
- 5 Avian Power Line Interaction Committee. 2006. *Suggested Practices for Avian Protection on Power*
6 *Lines: State of the Art in 2006*. Edison Electric Institute, APLIC, and the California Energy
7 Commission. Washington, DC and Sacramento, CA.
- 8 Babcock, K. W. 1995. Home Range and Habitat Use of Breeding Swainson's Hawks in the Sacramento
9 Valley of California. *Journal of Raptor Research* 29:193-197.
- 10 Beaulaurier, D. L. 1981. *Mitigation of Bird Collisions with Transmission Lines*. Bonneville Power
11 Administration, Portland, OR.
- 12 Beedy, E. C. and W. J. Hamilton. 1999. Tricolored Blackbird (*Agelaius tricolor*). In: *The Birds of North*
13 *America, No. 423*. American Ornithologists' Union.
- 14 Bevanger, K. 1994. Bird Interactions with Utility Structures: Collision and Electrocution, Causes and
15 Mitigating Measures. *Ibis* 136:412-425.
- 16 Bevanger, K. 1998. Biological and Conservation Aspects of Bird Mortality Caused by Electricity
17 Power Lines: A Review. *Biological Conservation* 86: 67-76.
- 18 Brown, W. M., and R. C. Drewien. 1995. Evaluation of Two Power Line Markers to Reduce Crane and
19 Waterfowl Collision Mortality. *Wildlife Society Bulletin* 23:217-227.
- 20 Brown, W. M., R. C. Drewien, and E. G. Bizeau. 1987. Mortality of Cranes and Waterfowl from
21 Powerline Collisions in the San Luis Valley, Colorado. In: J. C. Lewis (ed.). *Proceedings of the*
22 *Crane Workshop, 1985*. Platte River Whooping Crane Maintenance Trust, Grand Island, Nebraska.
23 Pages 128-136.
- 24 California Department of Fish and Game. 2010. *California Natural Diversity Database (CNDDDB)*.
25 Available: <<http://dfg.ca.gov/biogeodata/cnddb/>>.
- 26 California Department of Fish and Game. 2012. *California Natural Diversity Database (CNDDDB)*.
27 Available: <<http://dfg.ca.gov/biogeodata/cnddb/>>.
- 28 Comrack, L. A. 2008. Yellow-Breasted Chat (*Icteria virens*), California Bird Species of Special
29 Concern: A Ranked Assessment of Species, Subspecies, and Distinct Populations of Birds of
30 Immediate Conservation Concern in California. In: W. D. Shuford, and T. Gardali (eds.). *Studies of*
31 *Western Birds 1*. Western Field Ornithologists, Camarillo, California, and California Department
32 of Fish and Game, Sacramento.

- 1 Crowder, M. R., and O. E. Rhodes. 2001. Avian Collisions with Power Lines: A Review. In: R. G. Carlton
2 (ed). *Proceedings of a Workshop on Avian Interactions with Utility and Communication Structures*,
3 *Charleston, SC*. Electric Power Research Institute, Palo Alto, California. Pages 139–168
- 4 Delta Habitat Conservation and Conveyance Program. 2011. *2009 to 2011 Bay Delta Conservation*
5 *Plan EIR/EIS Environmental Data Report*. Review Draft 1. December. Prepared for consideration
6 by the Lead Agencies.
- 7 Drewien, R. C., W. M. Brown, and W. L. Kendall. 1995. Recruitment in Rocky Mountain GSCs and
8 Comparison with Other Crane Populations. *Journal of Wildlife Management* 59:339–356.
- 9 Eckerle, K. P. and C. F. Thompson. 2001. Yellow-Breasted Chat (*Icteria virens*). A. Poole (ed.). *The*
10 *Birds of North America Online*. Ithaca: Cornell Lab of Ornithology. Available:
11 <<http://bna.birds.cornell.edu/bna/species/575>>.
- 12 Eddleman, W. R., R. E. Flores, and M. Legare. 1994. Black Rail (*Latterallus jamaicensis*). In: A. Poole
13 (ed.). *The Birds of North America*. Cornell Lab of Ornithology, Ithaca, NY. Retrieved from the
14 Birds of North American Online: <<http://bna.birds.cornell.edu/bna/species/123>>.
- 15 England, A. S., M. J. Bechard, and C. S. Houston. 1997. Swainson's Hawk (*Buteo swainsoni*). In *Birds of*
16 *North America* no. 265. A. Poole and F. Gill (eds.). The Academy of Natural Sciences, Philadelphia,
17 PA, and The American Ornithologists' Union, Washington, D.C.
- 18 Erickson, W. P., G. D. Johnson, and D.P. Young Jr. 2005. *A Summary and Comparison of Bird Mortality*
19 *from Anthropogenic Causes with an Emphasis On Bird Collision*. USDA Forest Service Technical
20 Report. PSW-GTR-191. 2005.
- 21 Estep, J. A. 1989. *Biology, Movements, and Habitat Relationships of the Swainson's Hawk in the Central*
22 *Valley of California, 1986-87*. California Department of Fish and Game. Unnumbered Report.
- 23 Estep, J. A. 2008. *The Distribution, Abundance, and Habitat Associations of the Swainson's Hawk*
24 *(Buteo swainsoni) in Yolo County*. Prepared by Estep Environmental Consulting for Technology
25 Associates International Corporation and the Yolo County Habitat/Natural Community
26 Conservation Plan JPA.
- 27 Evens, J., G. W. Page, S.A. Layman and R. W. Stallcup. 1991. Distribution, relative abundance and
28 status of the California black rail in western North America. *Condor* 93: 952–966.
- 29 Fitzner, R. E. 1980. Behavioral ecology of the Swainson's hawk (*Buteo swainsoni*) in southeastern
30 Washington. Pac. NW Lab PLN-2754.
- 31 Gabrielson, I. N., and S. G. Jewett. 1940. *Birds of Oregon*. Oregon State College, Corvallis, OR.
- 32 Gaines, D. 1974. A New Look at the Nesting Riparian Avifauna of the Sacramento Valley, California.
33 *Western Birds* 5: 61–79.
- 34 Halterman, M. D. 1991. *Distribution and Habitat Use of the Yellow-Billed Cuckoo (Coccyzus*
35 *americanus occidentalis) on the Sacramento River, California, 1987-90*, MS Thesis: California
36 State University: Chico, CA.

- 1 Haug, E. A., B. A. Millsap, and M. S. Martell. 1993. Burrowing Owl (*Speotyto cunicularia*). In: *The Birds*
2 *of North America, No. 61*. A. Poole and F. Gill (eds.). Philadelphia: The Academy of Natural
3 Sciences; Washington D.C.: The American Ornithologist's Union.
- 4 Herzog, S. K. 1996. Wintering Swainson's Hawks in California's Sacramento-San Joaquin River Delta.
5 *Condor* 98:876-879.
- 6 Hunting, K. 2002a. *Avian Power Line Collisions: Potential Impact on Central Valley Bird Populations*.
7 Available: <http://www.cvbirds.org/CVBC_Bull/V.5no.4/V5no.4pp62-65.pdf>.
- 8 Hunting, K. 2002b. *A Roadmap for PIER Research on Avian Collisions with Power Lines in California*.
9 PIER 500-03=2-071F.
- 10 Information Center for the Environment. December 12, 2011. University of California, Davis.
11 Tricolored Blackbird Portal. Available: <<http://tricolor.ice.ucdavis.edu/>>.
- 12 Ivey, G. L., and C. P. Herziger. 2000. *Distribution of GSC Pairs in Oregon, 1999/00*. Oregon Department
13 of Fish and Wildlife Nongame Technical Report #03-01-00. Salem, OR.
- 14 Ivey, G. L., and C. P. Herziger. 2001. *Distribution of GSC Pairs in California, 2000*. California Dept. Fish
15 and Game, Sacramento, CA.
- 16 Ivey, G. L., and C. P. Herziger. 2003. Sandhill Crane Monitoring at Staten Island, San Joaquin County,
17 California 2002-03. A survey for The Nature Conservancy, Galt, CA.
- 18 Ivey, G. L., B. E. Dugger, C. P. Herziger, M. L. Casazza, and J. P. Fleskes. In Preparation [a]. *Distribution,*
19 *Abundance, and Timing of Arrival and Departure of Greater and Lesser Sandhill Cranes Wintering*
20 *in the Sacramento-San Joaquin River Delta Region Of California*. Proceedings of the 12th North
21 American Crane Workshop.
- 22 Ivey, G. L., B. E. Dugger, C. P. Herziger, M. L. Casazza, and J. P. Fleskes. In Preparation [b]. *Survival,*
23 *Movements, and Home Range Size of Greater and Lesser Sandhill Cranes Wintering the*
24 *Sacramento-San Joaquin Delta Region of California*.
- 25 James, B. W., and B. A. Haak. 1979. *Factors Affecting Avian Flight Behavior and Collision Mortality at*
26 *Transmission Lines*. Final Report of a Study for the Bonneville Power Administration, Portland,
27 OR.
- 28 Janss, G. F. E., and M. Ferrer. 2000. Common Crane and Great Bustard Collision with Power Lines:
29 Collision Rate and Risk Exposure. *Wildlife Society Bulletin* 28:675-680.
- 30 Jenkins, A. R., J. J. Smallie, and M. Diamond. 2010. Avian Collisions with Power Lines: A Global Review
31 of Causes and Mitigation with a South African Perspective. *Bird Conservation International*.
- 32 Jones, K. A., Krebs J. R., Whittingham, M. J. 2007. Vigilance in the third dimension: head movement
33 not scan duration varies in response to different predator models. *Animal Behavior* 74: 1181-
34 1187.
- 35 Laymon, S. A. 1998. Yellow-Billed Cuckoo (*Coccyzus americanus*). In *The Riparian Bird Conservation*
36 *Plan: A Strategy for Reversing the Decline of Riparian-Associated Birds in California*. California

- 1 Partners in Flight. Available: <[http://www.prbo.org/calpif/htmldocs/species/riparian/yellow-](http://www.prbo.org/calpif/htmldocs/species/riparian/yellow-billed_cuckoo.htm)
2 billed_cuckoo.htm>.
- 3 Laymon, S. A. and M. D. Halterman. 1989. *A Proposed Habitat Management Plan for Yellow-Billed*
4 *Cuckoos in California*. U.S. Department of Agriculture, Forest Service General Technical Report
5 PSW-110:272–277.
- 6 Laymon, S. A., P. L. Williams, and M. D. Halterman. 1997. *Breeding Status of the Yellow-billed Cuckoo*
7 *in the South Fork Kern River Valley, Kern County, California: Summary Report 1985 - 1996*.
8 Administrative Rep. U.S.D.A Forest Service, Sequoia National Forest, Cannell Meadow Ranger
9 District, Challenge Cost-Share Grant #92-5-13.
- 10 Littlefield, C. D. 2002. Winter Foraging Habitat of Greater Sandhill Cranes In Northern California.
11 *Western Birds* 33:51–60.
- 12 Littlefield, C.D., and G. L. Ivey. 2000. Conservation Assessment for Greater Sandhill Cranes Wintering
13 on the Cosumnes River Floodplain and Delta Regions of California. The Nature Conservancy,
14 Galt, CA.
- 15 LSA Associates, Inc. 2004a. Swainson’s hawk population and habitat use assessment. Prepared for
16 the Solano County Water Agency. Vacaville, California.
- 17 Marshall, J. T., Jr. 1948. Ecologic Races of Song Sparrows in the San Francisco Bay Region. Part 1:
18 Habitat and Abundance. *Condor* 50:193–215.
- 19 Meyer, J. R. 1978. *Effects of Transmission Lines on Bird Flight Behavior and Collision Mortality*.
20 Prepared for Bonneville Power Administration, Portland, OR.
- 21 Morkill, A. E., and S. H. Anderson. 1991. Effectiveness of Marking Powerlines to Reduce Sandhill
22 Crane Collisions. *Wildlife Society Bulletin* 19:442–229.
- 23 Murphy, R. K., S. M. McPherron, G. D. Wright, and K. L. Serbousek. 2009. *Effectiveness of Avian*
24 *Collision Averters in Preventing Migratory Bird Mortality from Powerline Strikes in the Central*
25 *Platte River, Nebraska*. Nebraska Game and Parks Commission, U.S. Fish and Wildlife Service, and
26 University of Nebraska, Kearney.
- 27 Olendorff, R.R and R.N. Lehman. 1986. *Raptor Collisions with Utility Lines: an Analysis using*
28 *Subjective Field Observations*. Pacific Gas and Electric. San Ramon, CA.
- 29 Pacific Flyway Council. 1997. *Pacific Flyway Management Plan for the Central Valley Population of*
30 *Greater Sandhill Cranes*. Pacific Flyway Study Committee (c/o Pacific Flyway Representative
31 USFWS), Portland, Oregon, USA.
- 32 Pogson, T. H. 1990. *Distribution, Abundance and Behavior of Greater Sandhill Cranes (Grus*
33 *Canadensis Tabida) Wintering in California’s Central Valley*. Ms. Thesis, University of Alaska,
34 September 1990.
- 35 Pogson, T. H. and S. M. Lindstedt. 1991. Distribution and Abundance of Large Sandhill Cranes, *Grus*
36 *Canadensis*, Wintering in California’s Central Valley. *Condor* 93:266–278

- 1 Poulin, R., L. D. Todd, E. A. Haug, B. A. Millsap, and M. S. Martell. 2011. Burrowing Owl (*Athene*
2 *cunicularia*). In: A. Poole (ed.). *The Birds of North America*. Cornell Lab of Ornithology, Ithaca, NY.
3 Retrieved from the *Birds of North American Online*:
4 <<http://bna.birds.cornell.edu/bna/species/061>>.
- 5 Rayner, J. M. V., 1988. Form and Function in Avian Flight. *Current Ornithology* 5:1–66.
- 6 Rush, Scott A., Karen F. Gaines, William R. Eddleman and Courtney J. Conway. 2012. Clapper Rail
7 (*Rallus longirostris*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of
8 Ornithology; Retrieved from the Birds of North America.
- 9 Savereno, A. J., L. A. Savereno, R. Boettcher, and S. M. Haig. 1996. Avian Behavior and Mortality at
10 Power Lines in Coastal South Carolina. *Wildlife Society Bulletin* 24:636–648.
- 11 Scott, R. E., L. J. Roberts, and C. J. Cadbury. 1972. Bird Deaths from Power Lines at Dungeness. *British*
12 *Birds* 65(7):273–286.
- 13 Spautz, H., N. Nur, and D. Stralberg. 2005. California black rail (*Laterallus jamaicensis coturniculus*)
14 distribution and abundance in relation to habitat and landscape features in the San Francisco
15 Bay estuary. USDA Forest Service General Technical Report PWS-GTR-191.
- 16 Tacha, T. C., S. A. Nesbit, and P. A. Vohs. 1992. Sandhill Crane (*Grus canadensis*). In: A. Poole, P.
17 Stettenheim, and F. Gill (eds.). *The Birds of North America, No. 31* (The Academy of Natural
18 Sciences of Philadelphia; The American Ornithologist's Union, Washington, DC.
- 19 Trulio, L. A. and J. G. Evens. 2000. California black rail. Pp. 341-345 In P. R. Olofson (ed.). Goals
20 Project. Baylands Ecosystem Species and Community Profiles: Life histories and environmental
21 requirements of key plants, fish and wildlife. Prepared by the San Francisco Bay Area Wetlands
22 Ecosystem Goals Project. San Francisco Bay Regional Water Quality Control Board, Oakland,
23 California.
- 24 U.S. Fish and Wildlife Service. 1998. *Draft Recovery Plan for the Least Bell's Vireo*. Portland, OR.
- 25 Walkinshaw, L. H. 1949. *The Sandhill Cranes*. Cranbrook Institute of Science Bulletin No. 29,
26 Bloomfield Hills, MI.
- 27 Walton, B. J. 1975. *San Francisco Bay Region Salt Marsh Song Sparrow Survey, 1974*. Report to
28 California Department of Fish and Game, Sacramento, CA.
- 29 Yee, M. L. 2008. *Testing the Effectiveness of an Avian Flight Diverter for Reducing Avian Collisions with*
30 *Distribution Power Lines in the Sacramento Valley, California*. Pier Final Project Report. Prepared
31 for the California Energy Commission Public Interest Energy Research Program. Available:
32 <<http://65.74.139.74/2007publications/CEC-500-2007-122/CEC-500-2007-122.PDF>>.
- 33 Zeiner, D. C., W. F. Laudenslayer, Jr., K. E. Mayer, and M. White (eds). 1990. California's Wildlife.
34 Volume 2. Birds. State of California, Department of Fish and Game. Sacramento, California.

1 **7.2 Personal Communications**

2 Best, Mike. Pacific Gas and Electric Avian Protection Program Manager. November 9, 2012—Email to
3 Paola Bernazzani. ICF International.

4 Breault, A. Waterfowl and Habitat Biologist, Canadian Wildlife Service, Delta BC. March 2012—
5 Presentation to the Pacific Flyway Study Committee.

6 Estep, Jim. September 11, 2012—Email to Elizabeth Strange and Rebecca Sloan, ICF International.
7 Regarding: Bird species presence in Plan Area and collision vulnerability.

8 Ivey, G.L. Wildlife Biologist. Greater Sandhill Crane Specialist. 2010 assessment of greater sandhill
9 crane use areas in the Sacramento-San Joaquin Delta.

10 Meese, R. J. Staff Research Associate, Department of Environmental Science and Policy (SEP) and
11 Information Center for the Environment (ICE), University of California, Davis. December 20,
12 2011—GIS data transfer, in Access database format, detailing known, historic breeding colony
13 sites in or near the Plan Area that had not yet been entered into ICE.

14 Pleiss, John. Sabre Industries. November 5, 2012—Phone conversation with Todd Jones, ICF
15 International. Regarding bird flight diverters. See also
16 <<http://www.tessco.com/products/displayProducts.do?bgCategoryId=803926>>.

17

1
2
3

Exhibit 1

Summary of Habitat, Distribution, and Occurrence of Covered Bird Species in the Plan Area

Common and Scientific Names	Status	Habitat and Distribution	Potential for Occurrence in the Plan Area
California black rail <i>Laterallus jamaicensis coturniculus</i>	BCC/T, FP	Nests and forages in saline, freshwater, or brackish emergent marshes with gently grading slopes and upland refugia with vegetative cover beyond the high-water line. Year-round range includes Suisun Marsh, San Pablo Bay, Morro Bay, a few patches in the Sierra Nevada foothills, and portions of southern California; winter range expands to include San Francisco Bay and the Marin County coast.	Several historic nesting occurrences documented in the southern half of the Plan Area. Survey in 2009 found one nest at White Slough and one in an instream island west of Stockton.
California clapper rail <i>Rallus longirostris obsoletus</i>	E/E, FP	Nests and forages in dense cordgrass and cattail marshes with vegetated refugia during the highest tides. Year-round near coastal range, surrounds San Francisco and San Pablo Bays, and documented at several locations in Suisun Bay.	Range does not include the Plan Area with the exception of Suisun Marsh.
Greater sandhill crane <i>Grus canadensis tabida</i>	-/T, FP (nesting, wintering)	Forages primarily in croplands with waste grain; also frequents grasslands and emergent wetlands. Winter range includes the Central Valley and Delta, Carrizo Plain, southern California south of the Salton Sea, and Colorado River. Breeds in northeastern California.	May forage during winter throughout the crane use area.
Least Bell's vireo <i>Vireo bellii pusillus</i>	E/E (nesting)	Nests and roosts in low riparian thickets of willows and shrubs, usually near water but sometimes along dry, intermittent streams. Formerly a common and widespread summer resident throughout Sacramento and San Joaquin Valleys, and in the coastal valleys and foothills from Santa Clara County south, but its numbers have drastically declined, and the species has vanished from much of its California range.	Does not occur in the Plan Area, but potentially could expand range with riparian restoration.

Common and Scientific Names	Status	Habitat and Distribution	Potential for Occurrence in the Plan Area
Suisun song sparrow <i>Melospiza melodia maxillaris</i>	BCC/SSC	Nests and forages in brackish water marshes dominated by cattails, tules, and pickleweed. Year-round range includes the marshes surrounding Suisun Bay, from the confluence of the Sacramento and San Joaquin Rivers to the Carquinez Strait.	Present in Suisun Marsh. However, not expected in the remainder of the Delta or Plan Area.
Swainson's hawk <i>Buteo swainsoni</i>	BCC/T (nesting)	Nests in isolated trees, open woodlands, and woodland margins; forages in grasslands and agricultural fields. Breeding range spans the Central Valley and Delta west of Suisun Marsh, northeastern California, and a few additional scattered sites. Most of the population migrates south of California in fall/winter, although a small number winters in the Delta.	A minimum of 85 nests were documented throughout the Delta during limited surveys in 2009; estimated total is over 300 pairs (Estep pers. comm.).
Tricolored blackbird <i>Agelaius tricolor</i>	BCC/SSC (nesting)	Nests colonially in large, dense stands of freshwater marsh, riparian scrub, and other shrubs and herbs; forages in grasslands and agricultural fields. Year-round resident throughout the Central Valley and the central and southern coasts, with additional scattered locations throughout California.	High potential to occur throughout the Plan Area.
Western burrowing owl <i>Athene cunicularia</i>	BCC/SSC (nesting)	Nests and forages in grasslands, agricultural fields, and low scrub habitats, especially where ground squirrel burrows are present; occasionally inhabits artificial structures and small patches of disturbed habitat. Year-round range includes the Central Valley and Delta and portions of the central coast, eastern California, and southern California.	May occur throughout the Plan Area where habitat is suitable; documented on Brannan Island and near Suisun Bay and Clifton Court Forebay.
Western yellow-billed cuckoo <i>Coccyzus americanus occidentalis</i>	FC, BCC/SE	Nests in valley, foothill, and desert riparian forest with densely foliated deciduous trees and shrubs, especially willows. Historically common but now a rare summer resident at isolated sites in Sacramento Valley in northern California and along Kern and Colorado River systems in southern California; occasionally documented in Colusa, Glenn, Butte, Sutter, and Yolo Counties within the last 20 years.	One occurrence of unconfirmed breeding within the Plan Area during 2009 BDCP surveys at a location north of Walnut Grove, California.

Common and Scientific Names	Status	Habitat and Distribution	Potential for Occurrence in the Plan Area
White-tailed kite <i>Elanus leucurus</i>	- /FP (nesting)	Forages in ponds, marshes, slow-moving streams, sloughs, and irrigation/drainage ditches; nests in nearby uplands with low, sparse vegetation. Year-round range spans the Central Valley, Coast Ranges and coast, Sierra Nevada foothills, and Colorado River.	May nest and forage throughout the Plan Area; documented in the Delta along the Sacramento River west of Stone Lake, and in the north-central and east-central Delta.
Yellow-breasted chat <i>Icteria virens</i>	- /SSC (nesting)	Nests and forages in riparian thickets of willow and other brushy tangles near water and thick understory in riparian woodland. Breeding range includes the northern Sacramento Valley, Cascade Range, Sierra Nevada foothills, northwestern California, most of the Coast Ranges, the Colorado River, and other scattered sites. Migrates south of California in fall/winter.	Nests in patches of the Plan Area where habitat is suitable; surveys in 2009 found more nests than expected, but not in all available habitat.

Exhibit 2

**Maps of Species Occurrences and Modeled Habitat
Relative to Proposed Powerlines**

1

2

3

4




Figure 2-1. Map of California Black Rail Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3
 4




Figure 2-2. Map of Clapper Rail Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3




Figure 2-3. Map of Greater Sandhill Crane Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3




Figure 2-4. Map of Least Bell's Vireo Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3
 4




Figure 2-5. Map of Suisun Song Sparrow Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
2
3
4




Figure 2-6. Map of Swainson's Hawk Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3
 4




Figure 2-7. Map of Tricolored Blackbird Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3
 4




Figure 2-8. Map of Western Burrowing Owl Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3
 4




Figure 2-9. Map of Western Yellow-Billed Cuckoo Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
 2
 3
 4

1





Figure 2-10 Map of White-Tailed Kite Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

2
3
4
5



Plan Area

- Plan Area
- Conservation Zone (CZ)
- Primary Nesting and Migratory Modeled Habitat
- Secondary Nesting and Migratory Modeled Habitat
- Suisun Marsh/Upper Yolo Bypass Nesting and Migratory Modeled Habitat

Occurrences

- CNDDB—Presumed Extant
- Non-CNDDB

Transmission Lines

- Proposed Permanent
- Proposed Temporary

0 5 10
Miles

1
2

Figure 2-11. Map of Yellow-Breasted Chat Occurrences and Modeled Habitat Relative to Proposed Transmission Lines

1
2
3

Attachment 5J.D
**Indirect Effects of the Construction of the BDCP
Conveyance Facility on Sandhill Crane**

Attachment 5J.D

Indirect Effects of the Construction of the BDCP Conveyance Facility on Sandhill Crane

Contents

		Page
6		
7	Attachment 5J.D Indirect Effects of the Construction of the BDCP Conveyance Facility on	
8	Sandhill Crane	5J.D-1
9	5J.D.1 Introduction	5J.D-1
10	5J.D.1.1 Sandhill Crane Habitat Use in the Plan Area	5J.D-2
11	5J.D.1.2 Noise Impacts on Sandhill Cranes	5J.D-2
12	5J.D.2 Existing Noise Environment Conditions	5J.D-3
13	5J.D.3 Methods and Assumptions for Noise Impact Analysis	5J.D-3
14	5J.D.3.1 Sensitivity to Noise and Thresholds for Mitigation	5J.D-3
15	5J.D.3.2 Construction Equipment Noise Estimates.....	5J.D-4
16	5J.D.3.3 Construction Traffic Noise Estimates	5J.D-6
17	5J.D.3.4 Impact Assessment Methods	5J.D-7
18	5J.D.4 Noise Impact Conclusions	5J.D-10
19	5J.D.5 Nighttime Lighting Impacts on Sandhill Cranes	5J.D-11
20	5J.D.6 Existing Artificial Lighting Conditions.....	5J.D-12
21	5J.D.7 Proposed Project-Related Artificial Light.....	5J.D-12
22	5J.D.8 Human Presence/Visual Disturbance Impacts on Sandhill Cranes	5J.D-13
23	5J.D.9 Existing Human Presence/Visual Disturbance Environment	5J.D-14
24	5J.D.10 Proposed Project-Related Human Presence/Visual Disturbance	5J.D-14
25	5J.D.11 Proposed Human Presence Avoidance and Mitigation Measures	5J.D-14
26	5J.D.12 References	5J.D-14

Tables

1
2
3
4
5
6
7
8
9
10
11
12
13

	Page
5J.D-1 Human Population Density and Associated Ambient Noise Levels	5J.D-3
5J.D-2 Commonly Used Construction Equipment Noise Emission Levels	5J.D-5
5J.D-3 Matrix of Construction Noise Sources at Each Construction Activity Area Type (at 50 feet).....	5J.D-6
5J.D-4 Calculated Distance to Noise Contours for Each Type of General Construction Noise Source	5J.D-7
5J.D-5 Construction Equipment Noise Emission Levels and Estimated Noise Levels in Foraging and Roosting Habitat.....	5J.D-9
5J.D-6 Acres of Sandhill Crane Habitat Affected by Increased Noise Levels from Project Construction	5J.D-10

Figures

1

2	5J.D-1	Greater Sandhill Crane and Stone Lakes NWR
3	5J.D-2	Greater Sandhill Crane Habitat
4	5J.D-3	Greater Sandhill Crane Indirect Impacts: General Construction and Truck
5		Traffic Noise (North)
6	5J.D-4	Greater Sandhill Crane Indirect Impacts: General Construction and Truck
7		Traffic (South)
8	5J.D-5	Greater Sandhill Crane Indirect Impacts: Pile Driver Noise (North)
9	5J.D-6	Greater Sandhill Crane Indirect Impacts: Pile Driver Construction (South)

10

1

Acronym and Abbreviations

dB	decibel
dBA	A-weighted decibel
BDCP	Bay Delta Conservation Plan
DWR	California Department of Water Resources
EIR/EIS	environmental impact report/environmental impact statement
GIS	geographic information system
IES	Illuminating Engineering Society
L _{dn}	day-night sound level
LOS	Level of Service
USFWS	U.S. Fish and Wildlife Service

2

3

Attachment 5J.D

Indirect Effects of the Construction of the BDCP Conveyance Facility on Sandhill Crane

5J.D.1 Introduction

This memo summarizes the research and analysis of potential indirect effects of the construction of the Bay Delta Conservation Plan (BDCP) conveyance facility on the greater and lesser sandhill cranes (*Grus canadensis tabida* and *Grus canadensis canadensis*, respectively), referred to collectively here as sandhill crane. The indirect effects that are the focus of this research are noise, lighting, and other visual disturbance. While each effect may act independently, the effects are often correlated (especially visual disturbance and noise). For this reason, indirect effects on sandhill cranes are often discussed in combination in the literature as “human disturbance.”

The construction of the conveyance facility will require a substantial amount of heavy equipment over prolonged periods, and is expected to generate noise, require nighttime lighting, and create visual disturbance.

Two studies addressing human disturbance effects to sandhill cranes (Armbruster and Farmer 1981; Norling et al. 1990) were highlighted in the Platte River Recovery Implementation Program (U.S. Fish and Wildlife Service 2006). The former study (Armbruster and Farmer 1981) summarizes guidelines based on input from a team of crane experts and is expert professional opinion. The latter study (Norling et al. 1990) is based on empirical measurements of distances of flock locations to various types of human disturbance. These studies both indicate an effect of human disturbance on sandhill crane habitat use in roosting and foraging habitat. These reports did not include noise level or lighting measurements, but looked at the overall effect of proximity to human disturbance, which could include the combined potential effects of noise, visual disturbance, and other direct and indirect habitat modification associated with the edge effects of the man-made features (e.g., habitat loss, change in microclimate, increased frequency of humans and/or domestic animals, changes in hydrology, increases in nonnative/invasive species).

For roost sites, buffers ranged from 100 meters for activities such as sand and gravel operations to 800 meters for commercial and urban land use. For cropland foraging habitat, buffers ranged from 10 meters for powerlines to 500 meters for commercial and urban land uses. However, the Platte River document acknowledges that “there is no consensus on the influence of human disturbances to potential crane habitat, or even how the concept of disturbance should be evaluated” (U.S. Fish and Wildlife Service 2006). As part of the process of developing their document, the U.S. Fish and Wildlife Service (USFWS) (2006) used a geographic information system (GIS) analysis to apply disturbance buffers to roost sites. They found that in several cases, known roost sites used by sandhill cranes were located well within the disturbance features’ described zones of influence. Conflicts in the body of research regarding sandhill cranes and human disturbance are further explored in Section 5J.D.8, *Human Presence/Visual Disturbance Impacts on Sandhill Cranes*.

1 **5J.D.1.1 Sandhill Crane Habitat Use in the Plan Area**

2 Sandhill cranes use the Plan Area primarily as winter habitat (September through March) and have
 3 many known habitat areas for roosting, foraging, and loafing behavior. These habitat areas occur in
 4 suitable croplands and wetlands, many of which are in close proximity to and directly within the
 5 proposed construction areas. Cranes spend the nighttime hours (dusk to dawn) at roost sites; the
 6 morning and evening hours in foraging habitat (generally, sunrise to 10:30 AM and 2:30 PM to
 7 sunset); and the midday (generally 10:30 AM to 2:30 PM) loafing in these areas and other areas
 8 without optimal foraging, but away from active human disturbances. Of particular interest are the
 9 habitat areas on the Stone Lakes National Wildlife Refuge and on Staten Island. Figures 5J.D-1 and
 10 5J.D-2 show the location of known permanent and temporary roosts in croplands and wetlands
 11 along with modeled potential roosting and foraging habitat.

12 **5J.D.1.2 Noise Impacts on Sandhill Cranes**

13 The evaluation of noise impacts on birds and their behavior is difficult. A summary of the effects of
 14 highway noise on birds in a Caltrans report (Caltrans 2007) provides a useful list of variables that
 15 could affect how noise is perceived by birds, resulting in the outcome of any noise-related indirect
 16 effects. As described in the Caltrans report, there are many complications in assessing the effects of
 17 noise independent of several confounding variables, many of which are relevant to this analysis.

18 Without taking each of these potential variables (and others) into consideration, appropriate
 19 correlations between road noise and bird behavior cannot be made. These variables include, but are
 20 not limited to:

- 21 1) Bird species and their style of acoustic communication;
- 22 2) Bird species and their behavior in the presence of adverse stimuli;
- 23 3) Age and experience of the birds;
- 24 4) Hearing capabilities of a species in quiet;
- 25 5) Hearing capabilities of a species in noise; and
- 26 6) Other kinds of stimuli associated with highways that might include (among others);
 - 27 a. Visual signals (vehicle movement);
 - 28 b. Vehicle-produced air pollution;
 - 29 c. Substrate vibrations resulting from the vehicles moving on the highway;
 - 30 d. The ecosystem near the roadway including substrate, vegetation, etc.; and
 - 31 e. Food supply near the highway.

32 While sandhill cranes do show some aversion to human disturbance (as described in Section 5J.D.1,
 33 *Introduction*), they are known to habituate to a certain degree to increased levels of background
 34 noise when the background noise level is relatively constant, such as roadway noise (Gary Ivey pers.
 35 comm.; Rod Drewien pers. comm.; David Brandt pers. comm.; Dwyer and Tanner 1992). However,
 36 less is known about the ability of sandhill cranes to habituate to intermittent noise such as that
 37 associated with the operation of heavy equipment at a scale construction site (e.g., pile drivers,
 38 construction cranes, compressors, heavy trucks). While the crane habitat use areas of concern in this
 39 analysis are generally in a rural setting, noises such as roadway traffic and agricultural operations
 40 can be heard within actively used areas, to which the cranes have apparently adapted.

5J.D.2 Existing Noise Environment Conditions

Primary noise sources in the project area are traffic traveling on surrounding freeways, highways, and rural roadways; agricultural operations; overhead commercial aircraft; and recreation related noise (e.g., fishing boats and waterski boats). Land uses near sandhill crane habitat are primarily rural and consist of agricultural use and low-density residential development. As such, existing noise levels are in the range of 40 to 50). Typical ambient sound levels as a function of human population density are presented in Table 5J.D-1, below.

Table 5J.D-1. Human Population Density and Associated Ambient Noise Levels

Human Population Density Type	dBA, L _{dn}
Rural	40-50
Small Town or quiet suburban residential	50
Normal suburban residential	55
Urban residential	60
Noisy urban residential	65
Very noisy urban residential	70
Downtown, major metropolis	75-80
Area adjoining freeway or near major airport	80-90
L _{dn} = Day-night sound level Source: Hoover and Keith 2000	

5J.D.3 Methods and Assumptions for Noise Impact Analysis

5J.D.3.1 Sensitivity to Noise and Thresholds for Mitigation

The general human response to changes in sound levels having similar frequency content (for example, comparing increases in continuous traffic sound levels) is summarized as follows.

- A 3 dB change in sound level is considered a barely noticeable difference.
- A 5 dB change in sound level will typically be noticeable.
- A 10 dB change in sound level is considered to be a doubling in loudness.

This may not be an appropriate metric for sandhill cranes. Because of the scarcity of data on unweighted intensities of source noise, for this analysis we assume that sandhill cranes, like most vertebrate animals, have a hearing sensitivity greater than that of humans, therefore, small changes in ambient noise (e.g., 3 dB) are assumed to be noticeable. Any errors this may introduce are compensated by use of a very conservative metric.

In this analysis we consider noise above 50 dBA to be potentially noticeable to sandhill crane, and thus to have a potential effect on foraging and roosting behavior. This very conservative approach is used in the absence of data on the effects of noise on the sandhill crane. USFWS uses 60 dBA as a

1 significance threshold for other sensitive bird species, including least Bell's vireo and California
2 gnatcatcher; this threshold is also supported by the California Department of Water Resources
3 (DWR) Specification 05-16 (California Department of Water Resources 2010) that suggests the
4 following guidelines for DWR construction projects:

5 Where ambient noise levels are less than 60 dBA and it is determined that construction related noise
6 will cause noise levels to exceed 60 dBA, or where the ambient noise levels are greater than 60 dBA
7 and it is determined that construction related noise will cause noise levels to exceed the ambient
8 level by 5 dBA, a temporary sound wall shall be constructed between the sensitive area and the
9 construction related noise source. The 60 dBA limit is not a regulatory requirement. Although the 60
10 dBA limit is not a regulatory requirement, it has been established as a threshold for establishing
11 noise impacts by consensus of experts, local and resource agencies, including the U.S. Fish and
12 Wildlife Service (USFWS). It is estimated that among other things, noise levels above 60 dBA may
13 interfere with communication among birds and other wildlife.

14 **5J.D.3.2 Construction Equipment Noise Estimates**

15 A wide variety of construction equipment will be used at each facility construction site and will vary
16 throughout the construction period. Impact pile driving was analyzed separately due to the unique
17 characteristics of noise produced from this noise source type (intermittent impact noise). Multiple
18 source construction noise was characterized by calculating the noise levels that would be produced
19 when the loudest six pieces of construction equipment were operating simultaneously, and noise
20 from heavy trucks was calculated assuming three heavy trucks operating in the same general area
21 simultaneously. Certain portions of the conveyance facility project area will have more limited
22 construction activity and construction noise sources, including borrow areas, spoils/muck areas,
23 and tunnel muck conveyor belt corridors. Table 5J.D-2 lists the typical noise levels from construction
24 equipment, and Table 5J.D-3 indicates which construction activity areas are likely to have each
25 general noise source type.

1

Table 5J.D-2. Commonly Used Construction Equipment Noise Emission Levels

Equipment	Typical Noise Level (dBA) 50 Feet from Source
Pile-driver (Impact)	101
Grader	85
Bulldozers	85
Heavy Truck	85
Loader	80
Air Compressor	80
Backhoe	80
Pneumatic Tool	85
Excavator	85
Auger Drill Rig (for drilled piles)	85
Crane, Derrick	88
Concrete Mixer Truck	79
Compactor (Ground)	83
Concrete mixer	85
Conveyor Belt Return/Load/Booster Drive	85
Conveyor Belt Mid-segment	75
Federal Highway Administration 2006, and conveyor belt equipment specifications. dBA = A-weighted decibel	

2

1 **Table 5J.D-3. Matrix of Construction Noise Sources at Each Construction Activity Area Type (at 50**
 2 **feet)**

Construction Activity Areas	Noise sources for analysis			
	Pile driver	Multiple Source Construction	Conveyor Belt	Heavy Trucks
<i>Noise level at 50 feet from Source</i>	<i>101 dBA</i>	<i>96 dBA</i>	<i>85/75 dBA</i>	<i>85 dBA</i>
Intake	<i>See detail</i>	X		
Coffer dam	X	X		
Waterside intake feature	X	X		
Sediment basins	X	X		
Intake forebay	X	X		
Electrical substation	X	X		
Forebay	<i>See detail</i>	X		
Outlet structure	X	X		
Inlet structure	X	X		
Electrical substation	X	X		
Siphons	X	X		
Barge Unloading Facility	X	X		
Shaft Location	X	X	X	
Permanent Surface Impact	X	X		
Temporary Surface Impact		X		
Operable Barrier		X		
Concrete Batch Plant		X		
Tunnel Muck Area			X	X
Intake Work Area				X
Pipeline Work Area				X
Tunnel Work Area				X
Control Structure Work Area				X
Safe Haven Work Area				X
Potential Borrow Area				X
Potential Spoil Area				X
Fuel Station				X
Road Work Area				X
Temporary Access Road Work Area				X

3

4 **5J.D.3.3 Construction Traffic Noise Estimates**

5 Construction traffic will be directed to many roads throughout the Plan Area, ranging from rural
 6 agricultural access roads to highways (e.g., State Route 12) to Interstate 5. Project related
 7 construction traffic will cause the largest increases in noise levels where high volume construction
 8 traffic is directed onto roads with low current traffic loads. Conversely, it will have minimal effect on
 9 existing noise levels on roads with existing high traffic loads (e.g., State Route 12 and Interstate 5).

1 **5J.D.3.4 Impact Assessment Methods**

2 To assess the potential effect of noise on sandhill cranes we calculated the noise level expected in
 3 known roosting/foraging habitat (at temporary and permanent roosts), and in modeled foraging
 4 habitat. Calculations assume direct line-of-sight (no intervening barriers) with an atmospheric noise
 5 attenuation rate of approximately 6 dB with each doubling of distance plus an additional attenuation
 6 of 1.5 dB noise absorption due to propagation over soft ground (e.g., agricultural land, open natural
 7 habitat). Therefore, total noise attenuation was calculated as 7.5 dB per doubling of distance from
 8 the source. For construction noise, distance to noise level contours was calculated from the edge of
 9 each identified construction area, giving a conservative worst-case estimate of noise levels since
 10 most of the construction activity won't take place on the perimeter of each site. Distance to traffic
 11 noise level contours were calculated from the centerline of each roadway. Traffic noise contours
 12 were calculated for all roadway segments included in the Level of Service (LOS) analysis in the
 13 traffic section of the environmental impact report/environmental impact statement (EIR/EIS) for
 14 the BDCP (California Department of Water Resources et al. 2012).

15 Noise propagation and attenuation can be affected by a variety of other factors including air
 16 temperature, atmospheric pressure, humidity, and wind speed and direction. These factors are
 17 highly variable over space and time and therefore are not typically included in standard
 18 environmental noise calculations. Because there are many highly variable factors, the assumption of
 19 a 7.5-dB attenuation per doubling of distance is a conservative estimate.

20 Table 5J.D-4 lists the calculated distances to noise contour lines from each type of general
 21 construction noise source. The noise contours were then overlaid on the sandhill crane modeled
 22 foraging habitat and known temporary and permanent roost habitat to determine the potential
 23 effects of construction noise on sandhill crane habitat.

24 **Table 5J.D-4. Calculated Distance to Noise Contours for Each Type of General Construction Noise**
 25 **Source**

Construction Site Noise Source Type ¹	Noise level at 50 ft	Noise Contours (feet from source)			
		Distance to 80 dBA	Distance to 70 dBA	Distance to 60 dBA	Distance to 50 dBA
Impact Pile Driver	101	350	850	2,100	5,250
General Construction ²	96	225	550	1,350	3,350
Heavy trucks ³	90	125	300	750	1,900
Conveyor Belt Return/Load (ends of conveyor) and Boosting Drives (inline at 1.5 mile intervals)	85	80	200	500	1,200
Conveyor Belt Mid-segment (along the length of belt between ends and boosting drives)	75		80	200	500
¹ Federal Highway Administration 2006, conveyor belt equipment specifications, and calculated as below.					
² Calculated assuming the six loudest pieces of construction equipment (except pile driver) operating simultaneously.					
³ Calculated assuming three heavy trucks operating simultaneously in same area of site.					

26
 27 The construction noise contours for general construction noise (all sources except pile driving) were
 28 combined with the construction traffic noise contours. Overlay of the noise contours on the modeled

1 foraging and known roost/forage areas depicts the expected worst-case noise levels to occur in
2 these areas during project construction based on the assumptions above. As previously mentioned,
3 pile driving noise was analyzed and displayed separately due to the unique characteristics of this
4 particular construction noise source (Figures 5J.D-3 and 5J.D-4 for all construction noise expect pile
5 driving; Figures 5J.D-5 and 5J.D-6 for pile driving).

6 Evaluation of the general project construction noise contours (all construction types except pile
7 driving) in relationship to the known roosting/foraging sites shows that there are nine areas where
8 general construction noise levels are expected to exceed 50 dBA (locations G1 through G9 on
9 Figures 5J.D-3 and 5J.D-4). Figures 5J.D-5 and 5J.D-6 show that noise levels for pile driving activities
10 are expected to exceed 50 dBA in five areas (locations P1 through P5 on Figure 5J.D-4 and 5J.D-5).
11 Modeled foraging habitat occurs adjacent to or in the near vicinity of much of the BDCP conveyance
12 facility construction area. Table 5J.D-5 shows the highest expected noise level for each construction
13 activity type at the nearest roost/forage site, and nearest modeled habitat, absent implementation of
14 minimization measures.

15 The traffic noise contours shown on Figures 5J.D-3 and 5J.D-4 are based on a combination of
16 construction and non-construction traffic. The noise contours are calculated for peak traffic loads,
17 therefore, they represent the loudest noise levels expected, which would typically be during daytime
18 and peak commuting hours. Based on the current project design and absent measures to minimize
19 noise in crane habitat, 50 dBA traffic noise contour will affect the following roost sites:

- 20 • temporary roost site north of Lambert Road between Franklin Boulevard and Bruceville Road;
- 21 • permanent roost site on Hood Franklin Road just below North Stone Lake;
- 22 • several permanent roosts along Interstate 5;
- 23 • edge of the temporary and permanent roost sites along Tyler Island Road;
- 24 • permanent roost sites south of State Route 12 on Bouldin Island; and
- 25 • permanent and temporary roost sites north and south of West 8 Mile Road.

1 **Table 5J.D-5. Construction Equipment Noise Emission Levels and Estimated Noise Levels in Foraging**
 2 **and Roosting Habitat**

Possible Construction Equipment	Typical Noise Level ¹ (dBA) at 50 ft from Source	Calculated Noise Level (dBA)	
		at Nearest Modeled Foraging Habitat (distance)	at Nearest Roost/Forage Site (distance)
Pile-driver (Impact)	101	101 (50 ft)	51 (5,000 ft)
Combined noise generation ³	96	96 (50 ft)	48 (4,000 ft)
Heavy Trucks ⁴	90	90 (50 ft)	55 (1,300 ft)
Muck Conveyor Belt Return/ Load and Boosting Drives	85	85 (50 ft)	55 (750 ft)
Conveyor Belt Mid-segment	75	75 (50 ft)	< 50 (750 ft)

¹ Federal Highway Administration 2006.
² Calculated based on assumed attenuation of 7.5 dB with each doubling of distance over soft ground.
³ Calculated assuming the six loudest pieces of construction equipment (except pile driver) operating simultaneously.
⁴ Calculated assuming three heavy trucks operating simultaneously in same area of site.
 dBA = A-weighted decibel

3

4 To quantify the total effect of the increase in construction noise on sandhill crane habitat, we
 5 calculated the acreage of each sandhill crane habitat type occurring within each 10 decibel range
 6 interval. Table 5J.D-6 summarizes those results showing that as much as 4,466 acres of habitat
 7 (3,868 acres modeled foraging, 120 acres permanent roosting, 477 acres temporary roosting) could
 8 be affected by noise levels above 60 dBA (not including pile driving), which would be noticeably
 9 above existing baseline noise levels (40–50 dBA) in most areas. Pile driving noise is expected to
 10 affect a smaller total acreage because pile driving is expected to occur at only a few project sites (see
 11 Table 5J.D-3 and Figure 5J.D-6). However, where pile driving does occur, the higher noise levels will
 12 increase the total acreage of habitat effects.

1 **Table 5J.D-6. Acres of Sandhill Crane Habitat Affected by Increased Noise Levels from Project**
 2 **Construction**

Noise Level Range	Habitat Types	Pile Driver (acres)	General Construction (acres)
>80 dBA	Modeled Foraging	16	624
	Roosting-Permanent	0	2
	Roosting-Temporary	0	64
	<i>Subtotal Habitat</i>	16	690
80-70 dBA	Modeled Foraging	73	913
	Roosting-Permanent	0	13
	Roosting-Temporary	3	107
	<i>Subtotal Habitat</i>	77	1,033
70-60 dBA	Modeled Foraging	661	2,332
	Roosting-Permanent	0	105
	Roosting-Temporary	75	306
	<i>Subtotal Habitat</i>	736	2,743
60-50 dBA	Modeled Foraging	5,491	8,013
	Roosting-Permanent	11	548
	Roosting-Temporary	755	1,085
	<i>Subtotal Habitat</i>	6,257	9,646

3

4 **5J.D.4 Noise Impact Conclusions**

5 Based on the assumptions and calculations in this analysis, in the absence of avoidance and
 6 minimization measures as much as 14,112 acres of crane habitat could experience noise levels
 7 above baseline levels as a result of general construction, and as much as 7,086 acres could
 8 experience noise levels above baseline levels as a result of pile driving activity.

9 Note that this analysis was conducted based on the assumption that there was direct line-of-sight
 10 from sandhill crane habitat areas to the construction site, and therefore is a worst-case estimate of
 11 effects. In many areas existing levees will partially or completely block the line-of-sight and will
 12 function as effective noise barriers substantially reducing noise transmission. Additionally, as
 13 described above, in the absence of data indicating the effect that noise levels above baseline would
 14 have on greater sandhill crane, a conservative approach was used by assessing noise levels above 50
 15 dBA even though the standard significance threshold for DWR is 60 dBA.

16 Sandhill cranes have been observed to habituate to increased levels of roadway noise (Gary Ivey,
 17 pers. comm.; Rod Drewien pers. comm.; David Brandt pers. comm.; Dwyer and Tanner 1992);
 18 however, little is known about their response to intermittent noise (Gary Ivey, pers. comm.; Rod
 19 Drewien pers. comm.; David Brandt pers. comm.). As stated in the Platte River Recovery
 20 Implementation Program Final Environmental Impact Statement, "At present, there is no consensus
 21 on the influence of human disturbances to potential crane habitat, or even how the concept of
 22 disturbance should be evaluated." (U.S. Fish and Wildlife Service 2006). Therefore, it is not possible
 23 at this stage to draw definitive conclusions regarding the sandhill crane response to the increased
 24 noise environment expected to be caused by this project. We can conclude that the noise

1 environment will be affected and noise levels will increase in sandhill crane habitat by moderate
 2 levels over larger areas (e.g., up to 20 decibel increase on approximately 17,000 acres), and by high
 3 levels over a more limited area (e.g., 20-30 decibel increase over approximately 1,000 acres).

4 Avoidance and minimization measures may be implemented to reduce noise related effects on
 5 cranes. Measures to reduce effects may include designing the project to avoid noise producing
 6 activities near high crane use areas, reducing noise producing activities during the winter when
 7 cranes are present, reducing night time activities in the vicinity of crane roost sites, and installing
 8 noise barriers between construction and traffic activities and crane roost sites.

9 **5J.D.5 Nighttime Lighting Impacts on Sandhill Cranes**

10 There has been little research into the impact of artificial lighting on roosting birds. Most discussion
 11 of birds and lighting concerns attraction, disorientation, and collisions of nocturnal migrators
 12 and/or foragers while in transit (Raine et al. 2007, Poot et al. 2008, Evans Ogden 1996, Kerlinger
 13 2000). In addition, lighting-induced disorientation of migrating birds can make it very difficult for
 14 them to find a suitable roost location and can lead to collision and/or exhaustion (Raine et. al 2007).

15 Artificial lighting can have a number of potential impacts on birds that are not in migration.
 16 Nighttime lighting can affect foraging timing and efficiency and interfere with breeding and
 17 migration (Navara et. al 2007, Titulaer et. al 2012, Santos et. al 2010, Hill 1992). A number of studies
 18 show effects of artificial lighting on timing of avian life cycles, influencing breeding behavior and
 19 sleep-wake cycles in passerines (Raine et. al 2007, Dominoni et. al 2013, and Nordt and Klenke
 20 2013). In a field study in the Netherlands, Titulaer et al. (2012) found that providing an artificial
 21 light source near nest boxes increased feeding rates of great tits (*Parus major*). This finding suggests
 22 that artificial light may affect nesting birds' energy expenditure on parental care, potentially
 23 impacting the parents' overall fitness. According to Dominoni et. al (2013), nighttime light exposure
 24 can affect a bird's metabolism by causing them to be more active during the daytime. In their study
 25 of European blackbirds (*Turdus merula*), wild-caught individuals exposed to night lighting in the lab
 26 developed their reproductive systems earlier, with earlier maturation of testes, earlier singing, and
 27 earlier molting. In a field study located in Germany, Nordt and Klenke (2013) found that urban
 28 blackbirds started singing up to 5 hours earlier than their rural counterparts.

29 No studies were identified that examined the effects of artificial lighting on roosting birds, including
 30 cranes. Direct light from automobile headlights has been observed to cause roosting cranes to flush
 31 and it is thought that they may avoid roosting in areas where lighting is bright (Ivey, pers comm).
 32 However, cranes exhibit high roost site fidelity and in some cases may still use artificially lit sites
 33 due to this loyalty. If the birds do use artificially lit roosting sites, they may be vulnerable to the
 34 sleep-wake cycle shifts and reproductive cycle shifts discussed above. Potential risks include a
 35 reduction in the cranes' quality of nocturnal rest, and changes to their sense of photoperiod which
 36 might cause them to shift their physiology towards earlier migration and breeding (Ivey, pers
 37 comm). Impacts such as these could prove detrimental to the cranes' overall fitness and
 38 reproductive success (which could in turn have population-level impacts). A change in photoperiod
 39 interpretation may also cause cranes to fly out earlier from roost sites to forage, and might increase
 40 their risk of power line collisions if they leave roosts before dawn (Ivey pers. comm.).

1 5J.D.6 Existing Artificial Lighting Conditions

2 Within the Plan Area, sandhill crane roosting sites are located near agricultural and rural residential
 3 land uses. Within agricultural areas in the Plan Area, artificial light is generally absent. The
 4 landscape is dark at night, except for occasional views of farmsteads dispersed through the
 5 landscape. Within rural portions of the Plan Area, lighting is related to the varied building sources
 6 (interior and exterior lighting and signage). Street lighting may be present but often is limited in
 7 extent.

8 5J.D.7 Proposed Project-Related Artificial Light

9 Construction of each intake structure would take up to 4 years to complete and would occur Monday
 10 through Friday for up to 24 hours per day. Dewatering near intakes, pumping plants, and certain
 11 pipeline construction areas and north of the intermediate forebay would take place 7 days per week
 12 and 24 hours per day. Evening and nighttime construction activities would require the use of bright
 13 lights. Nighttime construction could also result in headlights flashing into roost sites when
 14 construction vehicles are turning onto or off of construction access routes. Proposed surge towers
 15 would require the use of safety lights that would alert low-flying aircraft to the presence of these tall
 16 structures.

17 Completed BDCP facilities would require safety lighting. Lighting equipment associated with BDCP
 18 facilities would increase the amount of nighttime lighting in the Delta above existing ambient light
 19 levels. In particular, security lighting for the Sacramento River intakes and their associated pumping
 20 stations and facilities would create very noticeable effects relating to increased nighttime light at
 21 those locations. Lighting would be designed in accordance with guidance given by DWR's WREM No.
 22 30a, *Architectural Motif, State Water Project* and through coordination with local agencies through
 23 an architectural review process. This guidance is set forth as follows.

24 All artificial outdoor lighting is to be limited to safety and security requirements. All lighting is to
 25 provide minimum impact on the surrounding environment and is to be shielded to direct the light
 26 only towards objects requiring illumination. Lights shall be downcast, cut-off type fixtures with non-
 27 glare finishes set at a height that casts low-angle illumination to minimize incidental spillover of light
 28 onto adjacent properties, open spaces or backscatter into the nighttime sky. Lights shall provide good
 29 color rendering with natural light qualities with the minimum intensity feasible for security, safety
 30 and personnel access. All outdoor lighting will be high pressure sodium vapor with individual
 31 photocells. Lighting will be designed per the guidelines of the Illuminating Engineering Society (IES).
 32 Additionally, all lights shall be consistent with energy conservation and are to be aesthetically
 33 pleasing. Lights will have a timed on/off program or will have daylight sensors. Lights will be
 34 programmed to be on whether personnel are present or not.

35 Although the lighting would be designed to be shielded and oriented in such a manner minimize
 36 illumination of the immediate surroundings, these types of light generate an ambient nighttime
 37 luminescence that is visible for substantial distances from a large portion of the Delta.

38 Measures that may be implemented to reduce lighting effects on cranes include: limiting
 39 construction to daylight hours in the vicinity of crane roosts; locating nighttime construction
 40 lighting away from crane roost sites; avoiding nighttime construction activity in frequently used
 41 flight paths; routing truck traffic to reduce headlight impacts in roosting habitat; operating portable

1 lights at the lowest possible wattage and height; limiting the number of nighttime lights; and
2 installing visual barriers

3 **5J.D.8 Human Presence/Visual Disturbance Impacts** 4 **on Sandhill Cranes**

5 It is possible that the general presence and movement of humans, vehicles, and other equipment
6 could disturb sandhill cranes within the study area. The effect of human presence on cranes is at
7 least somewhat confounded with that of human-caused noise and light. As mentioned above, there is
8 conflicting information regarding the level of disturbance caused to sandhill cranes by human
9 activities (U.S. Fish and Wildlife Service 2006). The use of different types of behavioral response as
10 measures of disturbance (including roost site selection, vigilance, and flushing) may account for
11 some of the difference in disturbance response observed. Some studies have shown that, while
12 sandhill cranes do show a response to human presence, it does not appear to be a substantial
13 response in many cases (Wilkins 2012, Eldred 2009), and some degree of habituation does seem to
14 occur over time (Norling et al. 1992).

15 Studies show that roost site selection is impacted by a number of human activities. In their study
16 along the Platte River in Nebraska, Norling et al. (1992) found that human disturbance influenced
17 selection of roost sites by cranes. The cranes were most likely to avoid areas near paved and gravel
18 roads, bridges, and single dwellings with considerable human activity. Alternatively, roost site
19 selection did not seem to be impacted by human activities at urban dwellings, gravel pits, private
20 roads, railroads and powerlines.

21 Studies that used flushing and vigilance as measures of disturbance show less of an impact on crane
22 behavior from human activities. Increasing numbers of visitors to a bird-watching festival at a
23 sandhill crane staging site in Colorado did not result in increased vigilance or increased flushing on
24 the part of the cranes (Wilkins 2012). An increase in vigilance was observed in cranes at the refuge
25 hosting the festival when compared to cranes at an off-refuge site nearby, but this change was not
26 attributed to the presence of the birdwatchers. Similarly, in his study of sandhill cranes nesting and
27 staging in southern Michigan, Eldred (2009) found that increased traffic near roosting sites did not
28 result in increased vigilance on the part of cranes. In fact, Eldred reports that “even with heavy
29 disturbance, it appears that cranes will not increase their time in the alert investigative posture.”

30 However, research using vigilance to measure disturbance in red-crowned cranes does show a
31 disturbance response, indicating that other species of crane may be more susceptible to human
32 disturbance than sandhill cranes. A study of overwintering red-crowned cranes in China showed
33 that increased human disturbance does significantly increase vigilance behavior, thereby decreasing
34 time available for foraging (Wang 2011). In this study, birds’ movement to a less disturbed area did
35 not mitigate the impacts, as many birds moved to less disturbed areas; the concentration of birds in
36 less disturbed areas resulted in continued high levels of vigilance due to intraspecies competition
37 and, therefore, continued lost foraging time. This lost foraging time, if experienced by a large
38 number of individuals, has population-level implications for the cranes.

39 While cranes may be impacted to varying degrees by human presence, it also seems that they can
40 habituate to disturbance to some extent. Eldred (2009) points out that while a disturbance such as a
41 home construction site does seem to bother cranes, it appears they are capable of adapting to “low

1 levels of human presence.” In addition, Norling (1992) suggests that “some form of acclimation”
2 occurs in sandhill cranes subject to disturbance from commercial and urban development.
3 Therefore, it is possible that the cranes could make some degree of adaptation to the project-related
4 increase in general human presence in the plan area.

5 **5J.D.9 Existing Human Presence/Visual Disturbance** 6 **Environment**

7 Existing human presence in and near the crane roosting and foraging areas is generally limited to
8 levels consistent with a rural/agricultural environment. Moderate to high traffic volume currently
9 exist on major highways near crane roosting and foraging habitat (e.g., Interstate 5, State Route 12).

10 **5J.D.10 Proposed Project-Related Human** 11 **Presence/Visual Disturbance**

12 The increase in human presence and visual disturbance will generally be correlated with the
13 intensity of construction activity in the project area, and along roadways where construction-related
14 traffic will occur. As discussed, increased noise and lighting are directly linked to these activities and
15 therefore, it is not possible to determine if there is an additional disturbance effect from human
16 presence above what would occur from increased noise and lighting.

17 **5J.D.11 Proposed Human Presence Avoidance and** 18 **Mitigation Measures**

19 Since human presence within the study area is so heavily linked with noise and artificial light
20 sources in the study area, the noise and lighting mitigation measures above would also be expected
21 to reduce any added effect from human presence and related visual disturbance.

22 **5J.D.12 References**

23 **5J.D.12.1 Literature Cited**


- 24 Armbruster, M. J., and A. H. Farmer. 1981. Draft Sandhill Crane Habitat Suitability Model. In: J. C.
25 Lewis (ed.). *Proceedings 1981 Crane Workshop*. Tavernier, FL: National Audubon Society Pages
26 136–143.
- 27 California Department of Water Resources. 2010. *Supplementary Information for EIR/EIS: Bay Delta*
28 *Conservation Plan*. Sacramento, CA
- 29 California Department of Water Resources, Bureau of Reclamation, U.S. Fish and Wildlife Service,
30 and National Marine Fisheries Service. 2012. *Environmental Impact Report/Environmental*

- 1 *Impact Statement for the Bay Delta Conservation Plan*. Prepared by ICF International.
2 Sacramento, CA. February.
- 3 Caltrans. 2007. *The Effects of Highway Noise on Birds*. Prepared for Caltrans by Environmental
4 BioAcoustics LLC, Rockville, MD. September 30.
- 5 Dominoni, Davide, Michael Quetting, and Jesko Partecke. 2013. Artificial Light at Night Advances
6 Avian Reproductive Physiology. *Proceedings of the Royal Society B*. 280: 1756.
- 7 Dwyer, N. C. and Tanner, G. W. 1992. Nesting Success in Florida Sandhill Cranes. *Wilson Bulletin*
8 104:22–31.
- 9 Elred, Trevor. 2009. *Vigilance Behavior and Land Use by Sandhill Cranes (Grus canadensis)*. Master's
10 Theses and Doctoral Dissertations. Paper 238.
- 11 Evans Ogden, Lesley J., 1996. Collision Course: *The Hazards of Lighted Structures and Windows to*
12 *Migrating Birds*. Fatal Light Awareness Program (FLAP), Paper 3.
- 13 Federal Highway Administration. 2006. *FHWA Roadway Construction Noise Model User's Guide*.
14 Available: <<http://ntl.bts.gov/lib/49000/49100/49175/rcnm.pdf>>.
- 15 Hill, David. 1992. *The Impact of Noise and Artificial Light on Waterfowl Behaviour: A Review and*
16 *Synthesis of Available Literature*. BTO Research Report No 61. British Trust for Ornithology,
17 Norfolk, U.K.
- 18 Hoover & Keith, Inc.. 2000. *Noise Control for Buildings and Manufacturing Plants*. Houston, TX
- 19 Kerlinger, Paul. 2000. *Avian Mortality At Communication Towers: A Review Of Recent Literature,*
20 *Research, And Methodology*. Prepared for U.S. Fish and Wildlife Service Office of Migratory Bird
21 Management by Curry & Kerlinger LLC, Cape May Point, NJ. March.
- 22 Navara, Kristen J. and Randy J. Nelson. 2007. The Dark Side of Light at Night: Physiological,
23 Epidemiological, and Ecological Consequences. *Journal of Pineal Research*.
- 24 Nordt, Anja and Reinhard Klenke. 2013. Sleepless In Town—Drivers of the Temporal Shift in Dawn
25 Song in Urban European Blackbirds. *PLoS ONE* 8(8): e71476.
- 26 Norling, B. S., S. H. Anderson, and W. A. Hubert. 1990. *The Influence of Water Depth, Unobstructed*
27 *Area, and Disturbance Features on the Selection Of Roost Sites by Sandhill Cranes Along the Platte*
28 *River, Nebraska*. Wyoming Cooperative Wildlife Research Unit Project Report to Bureau of
29 Reclamation, Grand Island, Nebraska.
- 30 Norling, B. S., S. H. Anderson, and W. A. Hubert. 1992. Nocturnal Behavior of Sandhill Cranes
31 Roosting in the Platte River, Nebraska. *Prairie Naturalist*. 23:17–20.
- 32 Poot, H., B. J. Ens, H. de Vries, M. A. H. Donners, M. R. Wernand, and J. M. Marquenie. 2008. Green
33 Light for Nocturnally Migrating Birds. *Ecology and Society* 13(2): 47.
- 34 Raine, Helen, John J. Borg, Andre Raine, Suzanne Bairner, and Matthew Borg Cardona. 2007. *Light*
35 *Pollution and Its Effect on Yelkouan Shearwaters In Malta; Causes and Solutions*. Prepared for EU
36 LIFE Nature Programme, Brussels.
- 37 Santos, Carlos D., Ana C. Miranda, Jose P. Granadeiro, Pedro M. Lourenco, Sara Saraiva, and Jorge M.
38 Palmeirim. 2010. *Acta Oecologica*. 36(2): 166-172.

- 1 Titulaer, Mieke, Kamiel Spoelstra, Cynthia Y. M. J. G. Lange, and Marcel E. Visser. 2012. Activity
2 Patterns during Food Provisioning are Affected By Artificial Light in Free Living Great Tits
3 (*Parus major*). *PLoS ONE* 7(5): e37377.
- 4 U.S. Fish and Wildlife Service. 2006. Platte River Recovery Implementation Program—Final
5 Programmatic Environmental Impact Statement. Sandhill Cranes Appendix, Platte River FEIS.
6 *Sandhill Crane Spring Habitat in Nebraska, and Potential Effects to resources from Proposed*
7 *Changes in River Corridor Management*. March 10.
- 8 Wang, Zhi, Zhongqiu Li, Guy Beauchamp, and Zhigang Jiang. 2011. Flock Size and Human
9 Disturbance Affect Vigilance of Endangered Red-Crowned Cranes (*Grus japonensis*). *Biological*
10 *Conservation* 144(1): 101-105.
- 11 Wilkins 2012: Wilkins, K. 2012. Effects of Birdwatchers on Sandhill Crane (*Grus Canadensis*)
12 Behavior at Spring Stopover Sites in the San Luis Valley, Colorado. Masters Thesis. Colorado
13 State University, Fort Collins.

14 **5J.D.12.2 Personal Communications**

- 15 Brandt, David. Wildlife Biologist. Northern Prairie Wildlife Research Center, U.S. Geological Survey.
16 June 20, 2013—Phone conversation with Scott Fleury, ICR, regarding sandhill crane response to
17 noise.
- 18 Ivey, Gary. International Crane Foundation. Bend, Oregon. August 25, 2013—Email correspondence
19 with Heather White, ICF, regarding nighttime lighting effects on cranes.
- 20 Drewien, Rod. May 2013—Conversations with Scott Fleury, ICF< regarding sandhill crane response
21 to noise.




K:\Projects\1\DW\000000_00_BDCP\data\Working\Chapter 5\273_GSHC_Indirect_Noise_Impacts_20130625\Fig_1_GSHC_StoneLakesNWR_20130720.mxd Date: 8/7/2013 Time: 3:16:34 PM 29381

	Plan Area
	Conservation Zone (CZ)
	Stone Lakes NWR
Habitat Type	
	Foraging
	Roosting-Permanent
	Roosting-Temporary

0 0.5 1 Miles

GIS Data Source: Conservation Zones, SAIC 2012; Plan Area, ICF 2012; Hydrological Subregions, ICF 2012; Restoration Opportunity Area, SAIC 2011.


Figure 1
Greater Sandhill Crane and Stone Lakes NWR



K:\Projects\1\DW\000000_00_BDCP\data\Working\Chapter_5\273_GSHC_Indirect_Noise_Impacts_20130625\Fig_2_GSHC_South_ImpactArea_20130720.mxd Date: 8/7/2013 Time: 3:27:50 PM 29391

GIS Data Source: Conservation Zones, SAIC 2012; Plan Area, ICF 2012; Hydrological Subregions, ICF 2012; Restoration Opportunity Area, SAIC 2011.


Figure 2
Greater Sandhill Crane Habitat



K:\Projects_1\DW\000000_00_BDCP\data\Working\Chapter-5\301_Road_Impacts_GSHC_Noise_Contours_20130904\Fig_3_IndirectNoiseImpacts_GeneralConstruction_North_GSHC_Roads_20130904.mxd Date: 9/11/2013 Time: 12:49:15 PM 29391


GIS Data Source: Stone Lakes NWR, Stone Lakes 2010; Construction Activities, DHCCP 2013.

Figure 3
Greater Sandhill Crane Indirect Impacts:
General Construction and Truck Traffic Noise (North)



GIS Data Source: Construction Activities, DHCCP 2013.


Figure 4
 Greater Sandhill Crane Indirect Impacts
 General Construction and Truck Traffic (South)



K:\Projects\1\DW\000000_00_BDCP\data\Working\Chapter 5\273_GSHC_Indirect_Noise_Impacts_20130625\Fig 5_IndirectNoiseImpacts_PileDriver_North_GSHC_20130720.mxd Date: 8/7/2013 Time: 3:22:14 PM 29391

GIS Data Source: Stone Lakes NWR, Stone Lakes 2010; Construction Activities, DHCCP 2013.

Figure 5
Greater Sandhill Crane Indirect Impacts:
Pile Driver Noise (North)



K:\Projects\1\DW\000000_00_BDCP\data\Working\Chapter 5\273_GSHC Indirect Noise Impacts 20130625\Fig. 6 Indirect Noise Impacts 20130720.mxd Date: 8/7/2013 Time: 3:31:34 PM 293301

GIS Data Source: Construction Activities, DHCCP 2013.

Figure 6
Greater Sandhill Crane Indirect Impacts:
Pile Driver Construction (South)

1
2
3
4
5

Attachment 5J.E
**Estimation of BDCP Impact on Giant Garter Snake
Summer Foraging Habitat (Acreage of Rice)
in the Yolo Bypass**

Date:	June 20, 2013
To:	Laura King Moon, Project Manager, BDCP California Department of Water Resources
Cc:	Carl Wilcox California Department of Fish and Wildlife
From:	Rebecca Sloan and Ellen Berryman ICF International Neil Clipperton and Jason Roberts California Department of Fish and Wildlife
Subject:	Estimation of BDCP Impact on Giant Garter Snake Summer Foraging Habitat (Acreage of Rice) in the Yolo Bypass

1

2 The giant garter snake has been shown to use rice in the Yolo Bypass as aquatic foraging habitat
3 throughout the summer. The amount of rice grown annually in the Yolo Bypass depends on a
4 number of factors, including the degree to which late season flooding in the bypass precludes the
5 preparation and planting of rice fields. BDCP Conservation Measure 2 (*CM2 Yolo Bypass Fisheries*
6 *Enhancement*) allows for late-season inundation within the Yolo Bypass, which would potentially
7 preclude the planting of rice in some portions of the bypass in some years. To estimate the loss of
8 giant garter snake aquatic foraging habitat (rice) in the Yolo Bypass as a result of CM2
9 implementation, we used geographic information systems (GIS) to intersect spatial representations
10 of a modeled, late-season inundation footprint associated with Fremont Weir operations under CM2
11 and the giant garter snake habitat model.

12 MIKE-21, a two-dimensional, hydrodynamic model, was used to estimate the spatial extent of
13 inundation in the Yolo Bypass under representative flow scenarios (cbec 2010a). Two versions of
14 the MIKE-21 model have been developed to inform Yolo Bypass effects analyses: one that includes
15 west side tributaries as well as flows that pass over the Fremont Weir (cbec 2010b), and one that
16 does not include the west side tributary flows (cbec 2010a). The version without west side tributary
17 flows was chosen for use in this analysis, because it is assumed that late-season flooding under CM2
18 would likely result from flows entering the bypass through a modified Fremont Weir and that west
19 side tributary flows in the late season would likely be negligible.

20 The hydrologic model was run for a range of flow scenarios between 1,000 and 6,000 cfs at 1,000-
21 cubic-foot-per-second (cfs) increments. Each flow scenario produced a spatially explicit inundation
22 footprint. The 4,000 cfs flow scenario produced the largest inundation footprint at 7,700 acres (cbec
23 2010a). To be conservative for the purposes of estimating habitat loss, the 4,000 cfs inundation
24 footprint was used in the GIS intersect.

1 The giant garter snake habitat model uses agricultural data from the California Department of Water
2 Resources (DWR) (2008) to model aquatic foraging habitat (rice). The 2008 DWR agricultural data
3 is used for all agricultural-related effects analyses in the BDCP and represents a year with relatively
4 high acreage of rice in the Yolo Bypass.

5 When intersected in GIS, the 7,700-acre inundation footprint overlaps with 1,662 acres of rice in the
6 2008 DWR agriculture dataset. The 1,662 acres of aquatic giant garter snake foraging habitat loss is
7 assumed to be permanent; that is, the preclusion of 1,662 acres of rice is assumed to occur annually,
8 resulting in the permanent loss of aquatic foraging habitat.

9 To conservatively estimate habitat loss, this analysis assumes late-season flooding occurs every
10 year. However, the actual frequency of late-season flooding as a result of CM2 implementation is
11 expected to be significantly less than annually. This conservative approach is considered
12 appropriate for the purpose of setting take limits for the annual loss of aquatic, summer foraging
13 habitat for the giant garter snake within the Yolo Bypass.

14 **Datasets**

- 15 1. 2008 Yolo County Land Use Survey Data (California Department of Water Resources 2008).
- 16 2. MIKE-21 4000 cfs flow scenario without Westside tributaries (cbec 2010a).

17 **Steps Using GIS Tools**

- 18 1. Queried out the areas attributed as rice in the 2008 Yolo County Land Use Data (California
19 Department of Water Resources 2008).
- 20 2. Intersected the “DWR 2008 Rice” data with the “MIKE-21 4,000 cfs inundation footprint” data to
21 create the dataset—“Rice within the 4,000 cfs Inundation Zone.”
- 22 3. In the “Rice within the 4,000 cfs Inundation Zone” layer, manually digitized remaining portions
23 of inundated rice fields to capture the total impacted acreage of agriculture.

24 **Result**

25 Estimated loss of rice is 1,662 acres.

26 **References**

- 27 California Department of Water Resources. 2008. 2008 Yolo County Land Use Survey Data.
- 28 cbec. 2010a. *Yolo Bypass Two-Dimensional Hydrodynamic Modeling*. Prepared for the Metropolitan
29 Water District of Southern California. May 2010.
- 30 cbec. 2010b. *BDCP Effects Analysis: 2D Hydrodynamic Modeling of the Fremont Weir Diversion*
31 *Structure*. Prepared for SAIC and the California Department of Water Resources. November
32 2010.